

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

**Pakistan
Journal of Biological Sciences**

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

Dynamics of Huanglongbing-associated Bacterium *Candidatus Liberibacter asiaticus* in *Citrus aurantifolia* Swingle (Mexican Lime)

^{1,2}José Abel López-Buenfil, ^{1,2}José Abrahán Ramírez-Pool, ¹Roberto Ruiz-Medrano, ¹María del Carmen Montes-Horcasitas, ²Claudio Chavarín-Palacio, ³Jesús Moya-Hinojosa, ²Francisco Javier Trujillo-Arriaga, ⁴Rosalía Lira-Carmona and ¹Beatriz Xoconostle-Cázares

¹Departamento de Biotecnología y Bioingeniería, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Av. IPN 2508 San Pedro Zacatenco, C.P. 07360, Ciudad de México, Mexico

²Servicio Nacional de Sanidad, Inocuidad y Calidad Agroalimentaria, Centro Nacional de Referencia Fitosanitaria, Unidad Integral de Servicios, Diagnóstico y Constatación, Km 37.5, Carretera Federal México-Pachuca, Tecámac, C.P. 55740, Estado de México, Mexico

³Facultad de Ingeniería Química, Benemérita Universidad Autónoma de Puebla, Colonia San Manuel, Ciudad Universitaria, C.P. 72592 Puebla, México, Mexico

⁴Hospital de Pediatría, Hospital General Siglo XXI del Instituto Mexicano del Seguro Social, Avenida Cuauhtémoc 330, Col. Doctores, C.P. 06720, Ciudad de México, Mexico

Abstract

Background: The bacterial disease citrus huanglongbing (HLB), associated with "*Candidatus Liberibacter asiaticus*" (CLas) has severely impacted the citrus industry, causing a significant reduction in production and fruit quality. In the present study, it was monitored the CLas population dynamics in symptomatic, HLB-positive Mexican lime trees (*Citrus aurantifolia* Swingle) in a tropical, citrus-producing area of Mexico. The objective of this study was to identify the dynamics of the population of huanglongbing-associated bacterium *Candidatus Liberibacter asiaticus* and its insect vector in *Citrus aurantifolia* Swingle (Mexican lime). **Materials and Methods:** Leaf samples were collected every 2 months over a period of 26 months for quantification of bacterial titers and young and mature leaves were collected in each season to determine preferential sites of bacterial accumulation. The proportion of living and dead bacterial cells could be determined through the use of quantitative real-time PCR in the presence of ethidium monoazide (EMA-qPCR). **Results:** It was observed a lower bacterial titer at high temperatures in the infected trees relative to titers in mild weather, despite a higher accumulation of the insect vector *Diaphorina citri* in these conditions. This study also revealed seasonal fluctuations in the titers of bacteria in mature leaves when compared to young leaves. No statistically significant correlation between any meteorological variable, CLas concentration and *D. citri* population could be drawn. **Conclusion:** Although, HLB management strategies have focused on vector control, host tree phenology may be important. The evaluation of citrus phenology, CLas concentration, ACP population and environmental conditions provides insights into the cyclical, seasonal variations of both the HLB pathogen and its vector. These findings should help in the design of integrative HLB control strategies that take into account the accumulation of the pathogen and the presence of its vector.

Key words: *Candidatus Liberibacter asiaticus*, huanglongbing, *Citrus aurantifolia*, population dynamics, plant management, epidemiology, bacterial quantification, environment influence

Received: November 30, 2016

Accepted: January 13, 2017

Published: February 15, 2017

Citation: José Abel López-Buenfil, José Abrahán Ramírez-Pool, Roberto Ruiz-Medrano, María del Carmen Montes-Horcasitas, Claudio Chavarín-Palacio, Jesús Moya-Hinojosa, Francisco Javier Trujillo-Arriaga, Rosalía Lira-Carmona and Beatriz Xoconostle-Cázares, 2017. Dynamics of huanglongbing-associated bacterium *Candidatus Liberibacter asiaticus* in *Citrus aurantifolia* Swingle (Mexican lime). Pak. J. Biol. Sci., 20: 113-123.

Corresponding Author: Beatriz Xoconostle-Cázares, Departamento de Biotecnología y Bioingeniería, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Av. IPN 2508 San Pedro Zacatenco, C.P. 07360 Ciudad de México, Mexico Tel: +52-55-57473318

Copyright: © 2017 José Abel López-Buenfil *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Citrus huanglongbing (HLB) is one of the most devastating diseases of citrus worldwide. This disease is associated with four phloem-restricted α -proteobacteria, "*Candidatus Liberibacter*" spp.,¹ which prevent the translocation of photoassimilates from source to sink plant tissues. The four reported bacterial species associated with HLB are "*Candidatus Liberibacter asiaticus*" (C.Las), which is the most widely distributed, infecting citrus crops in Asia, Africa and the America, "*Candidatus Liberibacter africanus*" (C.Laf) found only in Africa, "*Candidatus Liberibacter americanus*" (C.Lam) present only in Brazil¹ and "*Candidatus Liberibacter caribbeanus*" (C.Lca) recently reported in Colombia^{2,3}. The bacteria are transmitted by at least two insect vectors. The Asian citrus psyllid (ACP), *Diaphorina citri*, transmits three of the four species: C.Las, C.Lam and C.Lca^{1,4}, while *Trioza erytreae* only transmits C.Laf¹.

Citrus fruits are a high value crop in Mexico. The Pacific region of Mexico is renowned for its production of Mexican limes, with the state of Colima as one of the principal producers of this particular citrus fruit. According to data from the Food and Agriculture Organization, Mexico produces over two million tons of limes and lemons, making it the third largest producer globally, preceded only by India and China⁵. The total contribution of the citrus industry to the Mexican economy is valued at 567 million dollars⁶. The first reported cases of HLB in Mexico were detected in Mexican lime trees from the Tizimín municipality in Yucatán⁷. The HLB has the potential to greatly disrupt the total production of not just Mexican lemons and limes but all Mexican citrus fruits. Unfortunately, as of yet, there is no cure for HLB. So far in Mexico, HLB management strategies are based on vector control with chemical insecticides. Monitoring the ACP population using yellow sticky traps is an important phytosanitary measure used to decide which strategies for insecticide application to pursue⁶. Thus far, this has been the most effective way to mitigate the rapid spread of HLB in orchards when used in conjunction with disposal of infected trees and the use of healthy bud wood.

Attempts to correlate seasonal fluctuations in C.Las population with ACP populations and their effects on the spread and severity of HLB have yielded mixed results. Measurement of bacterial viability by propidium monoazide qPCR (PMA-qPCR) in graft-infected, greenhouse grown sweet orange and *Severinia buxifolia*, a non-host plant, across 20 months (August, 2010-April, 2012) demonstrated that the populations of live C.Las were lower during the winter months^{8,9}. In Brazil, the population dynamics of C.Las in naturally infected sweet orange trees were monitored for

2 years (April, 2010-April, 2012) using qPCR, revealing that bacterial concentrations tended to be highest in autumn and lowest in spring and summer⁹. A similar study carried out in Florida reported large numbers of HLB-positive trees diagnosed between August and January, though the highest bacterial concentrations were found in December¹⁰. Conversely, in Pakistan, the number of trees diagnosed with HLB was largest during the summer, in spite of fewer C.Las circulating in the ACP vector population¹¹.

Epidemiological models developed for HLB have assumed a linear relationship between infection and manifestation of symptoms because as of yet, it has not been possible to precisely determine the point where infection begins¹². Attempts to characterize the epidemiology of HLB are also hampered by the lack of studies on the variation of C.Las concentration in both plant tissues and the ACP vector¹³, again stemming from the difficulty in defining a starting point of infection. Under controlled conditions, the incubation period of HLB is estimated to range from 3-12 months. However, the incubation periods seen in the field can last ≥ 5 years^{14,15}. This discrepancy illustrates the importance of understanding how a plant pathogen progresses throughout its pathology and how that relates to crop management practices. So far, information on the behavior of C.Las in the field has been scarce since the complexity of its pathology makes it a difficult disease to observe.

It is worth noting that various technical reports have indicated a weak correlation between the concentration of the bacteria and the severity of the HLB symptoms. It seems as though the expression of symptoms in an infected tree depends on host factors such as citrus variety, age and nutritional status, among others^{13,16}. As such, visual quantification of the characteristic diffuse mottle in individual trees is not indicative of C.Las concentration in the plant¹⁶.

In this study, it was monitored HLB-infected Mexican lime trees in Tecomán, Colima, Mexico for over 2 years in order to track bacterial titer fluctuations and potentially reveal any seasonal patterns. Both living and dead C.Las concentrations were quantified in leaf midrib tissues using EMA-qPCR. Young and mature leaves were also sampled and analyzed separately to determine how the changes in C.Las titer vary across tissues in distinct stages of development. Counts of adult ACP within the orchard and meteorological data were also collected throughout the experimental period. While there were some slight seasonal variations in C.Las concentration observed, statistical analyses failed to uncover any meaningful relationships between the bacterial titers and the environmental variables considered in our study. To our knowledge, the data herein constitutes the first study on C.Las population dynamics in Mexican lime in this country. It will

serve as a useful complement to existing ACP control strategies; this will, in turn, produce more effective, integrated disease management protocols and yield further insight into the epidemiology of HLB.

MATERIALS AND METHODS

Tree selection: The state of Colima was chosen for the experiment because HLB is endemic to the area and it is one of the largest producers of Mexican lime. The trial was run in a citrus orchard called "Rancho el Pandelo y Las Animas" in the town of Tecomán, Colima State with the following geographical reference coordinates: 18.93487°-103.92576°. In June, 2013, leaf samples were collected from 3 years old trees, exhibiting the diffuse mottling typical of HLB for molecular diagnosis of HLB in the laboratory.

Total DNA purification and HLB detection: Finely minced leaf midrib tissue (100 mg) was transferred to a 1.5 mL microcentrifuge tube containing a porcelain pellet. About 550 μ L of PBS (50 mM KH_2PO_4 , 150 mM NaCl, pH 7.2) was added and the tissue was macerated using a TissueLyser LT homogenizer (Qiagen; Hilden, Germany). The DNA purification was performed with the commercial AxyPrep Multisource Genomic DNA Miniprep kit (Axygen Biosciences) according to the manufacturer's instructions. The DNA was resuspended in 100 μ L of elution buffer before quantifying concentration and purity with a NanoDrop 2000 spectrophotometer (Thermo Fisher, Scientific; Waltham, MA). All samples were stored at -20°C for further use.

Molecular detection of *CLas* by PCR: End-point PCR was employed for *CLas* detection using a previously described primer pair¹⁷. A T100 Thermal Cycler (Bio-Rad; Hercules, CA) was used to carry out all the 25 μ L PCR reactions containing 2 μ L of sample DNA (50 ng μ L⁻¹) and reagents at the following concentrations: 1x buffer (Invitrogen), 2.5 mM MgCl_2 (Invitrogen), 0.16 mM of each dNTP (Invitrogen), 0.5 μ M forward primer OI1 (5'-GCGCAAATGCGTGAGCGGTACCA-3'), 0.5 μ M reverse primer OI2c (5'-TCGGCCGCCCTTCGAAACCAT-3') and 1 U of platinum Taq (Invitrogen). The amplification program consisted of an initial denaturing step at 94°C for 5 min followed by 35 cycles of 94°C for 30 sec, 62°C for 30 sec, 72°C for 60 sec with a final extension at 72°C for 10 min. Positive HLB infection was confirmed by resolution of the 1160 bp PCR product on a 1% (w/v) agarose gel.

Twenty four HLB-positive trees, averaging 1.8 m in height, were selected. The trees were planted with an 8 m spacing between rows and 4 m between columns. The trees were isolated from the orchard with anti-aphid mesh tunnels

measuring 6×36×3 m high (12 trees structure⁻¹). Crop management was performed according to the regional practices. Leaf samples were taken every 2 months, starting in October, 2013.

Quantification of live and dead bacteria: Leaf samples for the determination of live HLB titers were collected in March (spring), May (summer), September (autumn) and December (winter). Seven young and mature leaves from four branches per tree were collected; all seven leaves from the same branch. Non-infected trees were similarly sampled to serve as a control. Young leaves were identified by their apical position on the selected branch while mature, photosynthetically active leaves were collected from the basal part of the same branch. Young and mature leaves were processed separately. The plant material was prepared for DNA extraction as follows: Leaf midribs (100 mg) were finely minced using disposable blades, weighed and transferred to a 1.5 mL microcentrifuge tube for homogenization with the TissueLyser LT (Qiagen). Duplicate samples were prepared: One for standard genomic DNA extraction and the other for extraction in the presence of Ethidium MonoAzide (EMA) to assess live/dead bacterial concentration. The homogenized tissue samples were stored at -80°C pending further processing.

DNA extraction from live bacteria: The DNA extraction from live bacteria was performed according to the previously described EMA method¹⁸ with the following modifications: About 300 μ L of PBS buffer (50 mM KH_2PO_4 , 150 mM NaCl, pH 7.2) was added to a 1.5 mL microcentrifuge tube containing plant tissue and the sample was agitated for 5 min. The EMA was added to the samples to a final concentration of 100 μ g mL⁻¹ and gently shaken for 5 min in the dark. The EMA was bound to free DNA by exposing the samples to a 650 W halogen bulb for 2 min while the samples were on ice. The samples were centrifuged at 10,000 rpm for 5 min, the supernatant was discarded and finally the tissue was macerated with a TissueLyser LT (Qiagen). The DNA extraction proceeded using the previously mentioned method. The DNA concentration and purity was determined with a NanoDrop 2000 spectrophotometer (Thermo Scientific).

***CLas* 16S recombinant plasmid construct:** A *CLas* 16S rDNA fragment was amplified using previously described primers OI1/OI2¹⁹ yielding a PCR product of 1160 bp, which was then cloned into the pGEM®-T-Easy vector (Promega; Madison, WI) by following the manufacturer's instructions. Chemically competent Mach1-T1 *E. coli* cells were transformed by heat shock and plated on LB agar with 100 μ g mL⁻¹ ampicillin. The bacterial colonies containing the insert were identified by

α -complementation. Plasmid purification was performed using the PureLink Quick Miniprep Plasmid Kit (Invitrogen; Thermo Fisher Scientific) according to the manufacturer's instructions. The DNA concentration was determined with a NanoDrop 2000 spectrophotometer (Thermo Scientific). To calculate a quantification curve, the plasmid copy number was determined as follows:

$$\text{Copies of 16S plasmid} = \frac{\text{Concentration (ng)} \times \text{Avogadro's number}}{\text{Length (bp)} \times 1 \times 10^9 \times 650}$$

where, 650 is the mass of each base pair in daltons and Avogadro's number is 6.022×10^{23} . Ten-fold serial dilutions of plasmid DNA at 10^8 , 10^7 , 10^6 and 10^5 copies μL^{-1} were prepared.

Detection of *C.Las* by quantitative real time PCR: The real-time PCR reactions for detecting *C.Las* used previously reported primers and probes^{17,19}. All qPCR amplifications were run on the CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad). Each 25 μL reaction contained 2 μL of sample DNA and the following reagents: 1x PCR buffer (Invitrogen), 2.4 mM MgCl_2 (Invitrogen), 0.24 mM of each dNTP (Invitrogen), 0.12 mM forward HLBas (5'-AGCTCGGCGGCATATATACG-3'), 0.12 mM reverse HLB (5'-TCCGCGTTACGTAGAAGGAAATAG-3'), 0.12 μM HLB probe (5'-6-FAM-AGACGGAGTGTGGCCAAC-3'BHQ-1), 0.12 mM forward COX (5'-GTATGCGCACACGTCTTCCAGA-3'), 0.12 mM reverse COX (5'-GCCAAAAGACTGCTGGCATTG-3'), 0.12 μM COX probe (5'-TET-ATCATGCAGGCACTTCTGG-3' BHQ-2) and 1 U Taq Platinum (Invitrogen). Each DNA sample and the standard curve reactions were assayed in triplicate. The qPCR assays were run at 94°C for 20 sec, followed by 40 cycles of 94°C for 1 sec then 58°C for 40 sec. Fluorescent signal capture was set for each 58°C step.

***C.Las* quantification:** Cycle threshold values, Ct, obtained from the standard curve reactions were used to draw a linear regression plot, from which the resulting equation was calculated: $y = -3.4282x + 46.596$, with an $R^2 = 0.9997$. Based on that equation, following formula was obtained for determining bacterial titer:

$$\text{Concentration} = 10^{\left(\frac{\text{ct} - 46.596}{-3.4282}\right)}$$

The *C.Las* concentrations in the experimental samples were determined by interpolating their Ct values.

Asian citrus psyllid quantification: Two sticky 24×18 cm yellow traps were placed at the center of each tunnel of Mexican lime trees. The old traps were collected and new

yellow traps were installed on the same dates as leaf sampling. Positive ACP identification and count was performed at the Laboratory of Entomology and Acarology of the National Phytosanitary Reference Centre.

Weather information: Meteorological data were obtained from the National Agricultural Meteorological Station Network of the National Institute of Forestry, Agriculture and Fisheries (INIFAP), from the Tecomán station in Colima, located at 18.96683°-103.84227°. Average temperature, minimum temperature, maximum temperature, average relative humidity and rainfall were all monitored during the trial period.

Statistical analysis: The database generated from the quantification of *C.Las* in the various samples obtained from distinct places on 24 Mexican lime trees was compiled in Microsoft Excel and then transferred to SAS²⁰ for an analysis of variation and identification of statistically significant differences between aggregation factors (evaluation dates, trees, branches, young and mature leaves) using the Fisher 'F' ratio of variances (PROC GLM in SAS). Having identified the significant relationship of the above factors, a multiple comparison of averages was applied with the Tukey test ($\alpha = 0.05$). Average values grouped in the same letter are not statistically different. The identification of possible statistical correlations between atmospheric variables, ACP counts from the yellow traps and concentration of *C.Las* was performed using the SAS PROC CORR procedure to generate the estimations of Pearson's "r" with a probability of $p < 0.05$. Calculated standard deviation is represented in graphs as error bars.

RESULTS

***C.Las* population dynamics in Mexican lime trees:** During the 2 years trial period, the total *C.Las* titer peaked numerous times (Fig. 1). The first peak, in December, 2013 shows an increase in the concentration of living bacteria. The next peak in total bacterial titer, in May, 2014, actually has a greater proportion of dead cells. The ratio of live/dead *C.Las* skewed towards live cells again in November, 2014. Interestingly, the total bacterial titer after the winter of 2014 never returns to the same levels, though there are two peaks in 2015, during July and December. In July, 2015, there was a majority of dead *C.Las*, while in December, 2015, there was a return to a majority of live cells. The ratio of live/dead bacteria was largest during the autumn and winter peaks (December, 2013, November, 2014 and December, 2015), while a higher proportion of dead *C.Las* was seen during the hot summer

Differences in *C.Las* titers between young and mature leaves: The quantification of live and dead *C.Las* in both developing and photosynthetic leaves indicated that a higher

total concentration of bacteria is present in mature leaves (Fig. 2). No disease symptoms were observed in the young leaves sampled, even though they contained high




Fig. 2(a-b): Differences in *C.Las* titers between young and mature photosynthetic leaves. The quantification of live and dead *C.Las* in both developing and photosynthetic leaves showed a higher bacterial concentration is present in mature leaves. (a) Bacteria present in young leaves, grey: Dead *C.Las*, white: Live *C.Las* and (b) Bacteria present in mature leaves, grey: Dead *C.Las*, white: Live *C.Las*. A-1 to A-4 represents pooled samples from 6 trees, while the collect month and year are indicated

concentrations of *C.Las*, which strongly supports the idea that the bacterial titer is not proportional to the severity of HLB symptoms.

Significant variation in bacterial concentration was observed within the same tree, ranging from 8.79×10^7 to 1.76×10^4 *C.Las* g^{-1} of tissue. Despite the fact that this tendency was similar for all sampled trees, in some cases the differences in concentration was very drastic. It was noted that, generally, the mature leaves contained a higher total concentration of bacteria, as well as a greater proportion of live *C.Las*, this is probably due to their high photoassimilate content, which can be employed as a carbon source by *C.Las* ($F = 12.45$, $p < 0.0001$). Yet, in the last two evaluations (September and December, 2015), the titers of dead bacteria were also higher in mature leaves than in developing leaves (Table 2). Other studies reported that young leaves harbor greater numbers of bacteria compared to mature leaves. Supposedly, this distribution takes advantage of the ACP feeding preference for younger leaves²¹ to bolster *C.Las* transmission, when in actuality the ACP seeks out new growth for deposition of eggs. Consequently, it is possible that the high concentrations seen in mature leaves are indicative of the bacterium attempting to evade high ambient temperatures by mobilizing towards the basal part of the branches¹¹.

Dynamics of *C.Las* population in 2015: The dynamics of living and dead *C.Las* were monitored in both young and mature leaves sampled during four different evaluation periods in 2015 (spring, summer, autumn and winter) (Fig. 3). Highest concentration of total bacteria occurred in spring, with dead bacteria present in low concentrations and mature leaves exhibiting higher titers (Fig. 3a, b). In summer, a drastic increase was observed in the concentration of dead bacteria (Fig. 3c, d). The trend in autumn showed a pattern similar to the one in summer. The concentration of dead bacteria exhibited an increase compared to the dynamics seen in the spring. However, dead bacteria were present at low concentrations in young leaves (Fig. 3e, f). In winter, dead bacteria declined overall, while in young leaves the total concentration of bacteria was noticeably lower than in mature leaves (Fig. 3g, h). The total concentration of *C.Las* was greater in older leaves in all seasons (Fig. 3b, d, f, h).

It is pertinent to mention that the major periods of citrus shoot emergence in Tecomán take place during the summer and autumn months. That period overlaps with spring and summer: The seasons with the greatest total concentrations of *C.Las* (Table 3). Nevertheless, the results of the statistical analyses indicate that the largest numbers of dead bacteria are present in these seasons, while, in autumn and winter, there is a recovery of living bacteria.

Weather conditions during the study: Tecomán, Colima State has a dry tropical climate: The average annual maximum and minimum temperatures were 32.65 and 20.78°C, respectively, with an average annual temperature of 26°C (Fig. 4), this tropical municipality is considered an isothermal area since the temperatures do not vary much during the four seasons. It is known that in tropical regions, the low temperatures during the nights induce new shoot growth in citrus²². For the state of Colima there are two periods of shoot emergence reported. The first and most important, is from November to March, while the second is from June until September, when the rains begin. During this study, new shoot growth was observed when the temperatures increased and the rains arrived. In fact, some slight new growth was recorded between the months of August and December.

Associations between *C.Las* population and external variables: An inverse, highly significant relationship ($r = -0.934$ Pearson, $p < 0.0001$) between the concentrations of live and dead bacteria emerged from the analyses of the Mexican lime tissues during the experimental period (Table 4). Based on the strength of this inverse relationship, it is suggested that the concentration of live bacteria correlates with a concomitant reduction of dead bacteria in Mexican lime, this could be caused by a compromised defense response on behalf of the plant, which could be affected by the temperature or other environmental variables. As a

Table 2: Average log concentration of *C.Las* found in young and mature leaves of Mexican lime

Leaf age	Total No. of samples	Average log <i>C.Las</i>	Tukey ($p < 0.05$)
Mature	1329	5.109	A
Young	1137	4.804	B
DMS = 0.085			

Table 3: Comparison of the average concentrations of live and dead *C.Las* in Mexican lime, based on tissue type and season

Parameters	Total young	Live young	Dead young	Total mature	Live mature	Dead mature
Summer	5.8676 ^A	3.9311 ^B	1.9267 ^A	6.7149 ^A	4.7150 ^B	1.9999 ^A
Spring	5.8507 ^A	4.9189 ^A	0.9318 ^{BC}	6.0414 ^B	5.0503 ^B	0.9911 ^B
Autumn	5.6848 ^A	5.0035 ^A	0.6813 ^C	6.8318 ^A	6.2547 ^A	0.5771 ^B
Winter	5.5889 ^A	4.2477 ^B	1.3412 ^B	6.4861 ^A	4.5187 ^B	1.9788 ^A
Mean value	5.7480 (0.697)	4.5253 (0.6601)	1.2202 (0.680)	6.5185 (0.5312)	5.1346 (0.7157)	1.3867 (0.5604)
DMS	0.527	0.4987	0.5139	0.4013	0.5406	0.4233




Fig. 3(a-h): Dynamics of *C.Las* population in 2015. Representation of living and dead *C.Las* in both young and mature leaves sampled during four different evaluation periods in 2015. (a, c, e, g) Young leaves and (b, d, f, h) Mature, photosynthetic leaves. Grey bars are dead bacteria, while white are alive. A-1 to A-24 are measurements of each tested tree

speculative note, it is also possible that quorum sensing may regulate oscillations in the bacteria population²³. There was no correlation between live bacterial concentration and the citrus Asian psyllid density, according to statistical analysis [Corr test,

parameters ($r = -0.137$, $p = 0.686$)]. Hence, this represents the first evidence from the Tecomán region, Colima, Mexico. In addition, there was a period of elevated bacterial concentrations between July and November, 2014 (Fig. 1),




Fig. 4(a-d): Weather conditions during the study, (a) Precipitation (mmHg), (b) Relative humidity (%), (c) Minimum temperature (°C) and (d) Maximum temperature (°C)

Table 4: Statistical correlations between *C.Las* titers and environmental variables

Parameters	Live	Dead	Temperature (°C _{med})	Relative humidity	Temperature (°C _{min})	Precipitation	ACP count
Live	1.000	-0.909<0.0001**	0.0065 (0.938) ^{ns}	0.0945 (0.758) ^{ns}	0.138 (0.651) ^{ns}	-0.153 (0.616) ^{ns}	-0.137 (0.686) ^{ns}
Dead		1.000	0.061 (0.841) ^{ns}	0.0118 (0.969) ^{ns}	-0.028 (0.925) ^{ns}	0.085 (0.781) ^{ns}	0.231 (0.494) ^{ns}
Temperature (°C _{med})			1.000	0.439 (0.132) ^{ns}	0.922 <0.0001**	0.500 (0.081) ^{ns}	-0.094 (0.782) ^{ns}
Relative humidity				1.000	0.594 (0.032)**	0.5192 (0.073) ^{ns}	0.166 (0.625) ^{ns}
Temperature (°C _{min})					1.000	0.524 (0.065) ^{ns}	-0.076 (0.822) ^{ns}
Precipitation						1.000	0.500 (0.117) ^{ns}
ACP count							1.000

occurring simultaneously with the period of new shoot emergence reported for the region. These sustained levels of *C.Las* were somehow related to the emergence of new growth, although our statistical analyses failed to establish a concrete correlation. Interestingly, it appears that the months with low nighttime temperatures are the same in which the highest concentrations of live bacteria are detected; conversely, there is an increase in the number of dead bacteria in the warmest months. Moreover, no relationship between bacterial titer and severity of HLB symptoms were observed.

DISCUSSION

The population dynamics of *C.Las* in Mexican lime were elucidated by monitoring the concentrations of both live and

dead bacteria throughout 2014 and 2015, identifying a similar pattern of cyclical behavior as previously⁹. This variation coincides with reports from Florida, USA and Paraná, Brazil and supports the hypothesis that the bacterial distribution changes throughout the year, depending on the host tree phenology and the environmental conditions^{8,9}. This cyclic variation should now be factored into the rationale of HLB management and ACP control strategies, since diagnostic results from sampling both plant material and psyllid vector can be skewed, depending on whether the *C.Las* population is in decline or on the rise.

The trends in variation of *C.Las* population in Mexican lime were observed over a period of a full year (January to December, 2014), similar to the trial that has been reported for *C.Las* in sweet orange⁸. On a speculative note, this annual

variation in the concentration of *C.Las* could be more closely associated with the phenomenon of Quorum Sensing (QS), in addition to the meteorological variables or ACP populations found in infected orchards. Laboratory tests with *C.Las*, grown in culture media supplemented with citrus juices, found that the viability of the bacteria showed cyclical growth patterns, with increasing and decreasing phases throughout the experimental period. Cyclic or oscillatory development is a feature of bacterial populations attributable to internal or external fluctuations in the system, mainly related to the presence of nutrients. According to some reports, QS phenomena are related to the physiological and nutritional state of the host. By knowing the distribution of viable and non-viable bacteria in infected trees, it is possible to infer the periods when ACP transmission of *C.Las* is most probable. In turn, the probability of transmission is also most likely related to the distribution of photoassimilates within the trees throughout the year. Therefore, it is necessary to characterize these physiological aspects of Mexican lime varieties, since the presence of photoassimilates contributes to the carbon sources available for *C.Las* growth and development.

The finding of higher *C.Las* concentrations in mature leaves, compared to young leaves, runs contrary to the trend reported by several research groups²⁴. In fact, the presence of major carbohydrate stores in the more photosynthetically active mature leaves would make them a more favorable environment for *C.Las* and could explain the difference in reported distributions^{11,25}.

This study demonstrated a lack of statistically significant relationships between environmental variables and *C.Las* concentration. The correlation values between the concentration of live bacteria and rainfall demonstrated an inversely proportional relationship (Pearson $r = -0.153$, $p = 0.616$), indicating that the number of living bacteria is reduced when precipitation increases in the region, albeit not in a statistically significant manner. Alternatively, a similar phenomenon occurred with ACP count and rainfall (Pearson's $r = -0.137$, $p = 0.686$), where the amount of ACP found in the yellow traps decreased with increasing rainfall in the region, although it was also not statistically significant (Table 4). This lack of correlations or presence of weak correlations between atmospheric variables, *C.Las* concentration and ACP populations has been reported in different citrus-growing areas of the world^{16,26}. While the evidence gathered thus far supports the notion of a dynamic *C.Las* population within an infected citrus tree, the research community has been unable to decisively associate any one environmental variable to bacterial titer.

These were time points when the average log concentrations of dead bacteria greatly surpassed the average log concentrations of live *C.Las*. The first instance was October, 2013, when the titer of live bacteria only reached $\log 4.3$ *C.Las* g^{-1} of tissue, compared to the $\log 6.7$ *C.Las* g^{-1} of tissue recorded for the dead *C.Las*. This represents a 100 fold increase in dead bacterial cells over living *C.Las*. The same was seen again in February, 2014, May, 2015 and September, 2015. This distinct proportion of dead to live cells could be associated with the formation of a biofilm. The dead cells quantified during these periods would be those that make up the biofilm and cover their living relatives. As the behavior is seen during warmer months, this could be another defense mechanism used by *C.Las* to protect itself from high temperatures. There have been recent reports of genes which encode components of bacterial communication systems found via quorum sensing in the *C.Las* genome^{26,27}, which could be a strategy used by these bacteria to regulate its population.

CONCLUSION

The evaluation of citrus phenology, *C.Las* concentration, ACP population and environmental conditions provides insights into the cyclical, seasonal variations of both the HLB pathogen and its vector. Depending on the season, bacterial population fluctuates, representing the first report on the seasonal variation of *C.Las* concentration in Mexican lime. The understanding of the fluctuation of HLB-associated bacteria and its insect vector populations will help in the design of integrative control strategies. This information should be consolidated into existing HLB agronomic management practices and decision-making policies.

ACKNOWLEDGMENTS

This study was supported by SENASICA 2014-2015 grant to BX-C and RR-M. The authors acknowledge the critical opinion of the present manuscript to the members of the laboratory.

REFERENCES

1. Antunes, L.C.M., R.B.R. Ferreira, M.M.C. Buckner and B.B. Finlay, 2010. Quorum sensing in bacterial virulence. *Microbiology*, 156: 2271-2282.
2. Beloti, V.H., G.R. Rugno, M.R. Felipe, A.D. Carmo-Uehara, L.F. Garbim, W.A. Godoy and P.T. Yamamoto, 2013. Population dynamics of *Diaphorina citri* Kuwayama (Hemiptera: Liviidae) in orchards of Valencia orange, Ponkan mandarin and Murcott Tangor trees. *Florida Entomol.*, 96: 173-179.

3. Bove, J.M., 2006. Huanglongbing: A destructive, newly-emerging, century-old disease of citrus. *J. Plant Pathol.*, 88: 7-37.
4. Da Graca, J.V. and L. Kornsten, 2004. Citrus Huanglongbing: Review, Present Status and Feature Strategies. In: *Diseases of Fruits and Vegetables: Volume I Diagnosis and Management*, Naqvi, S.A.M.H. (Ed.). Kuwer Academic Publisher, The Netherlands, pp: 229-245.
5. Davenport, T.L., 2000. Processes influencing floral initiation and bloom: The role of phytohormones in a conceptual flowering model. *HortTechnology*, 10: 733-739.
6. FAO., 2012. *Statistics*. Food and Agriculture Organization (FAO), Rome, Italy.
7. Folimonova, S.Y., C.J. Robertson, S.M. Garnsey, S. Gowda and W.O. Dawson, 2009. Examination of the responses of different genotypes of citrus to huanglongbing (citrus greening) under different conditions. *Phytopathology*, 99: 1346-1354.
8. Folimonova, S.Y. and D.S. Achor, 2010. Early events of citrus greening (Huanglongbing) disease development at the ultrastructural level. *Phytopathology*, 100: 949-958.
9. Fraser, L.R. and D. Singh, 1966. Greening virus, a threat to citrus industry. *Indian Hortic.*, 10: 21-22.
10. Gottwald, T.R., J.V. da Graca and R.B. Bassanezi, 2007. Citrus huanglongbing: The pathogen and its impact. *Plant Health Progr.* 10.1094/PHP-2007-0906-01-RV
11. Hu, H., A. Roy and R.H. Brlansky, 2014. Live population dynamics of *Candidatus Liberibacter asiaticus*, the bacterial agent associated with citrus huanglongbing, in citrus and non-citrus hosts. *Plant Dis.*, 98: 876-884.
12. Irey, M.S., T. Gast, J. Cote, P. Gadea, O. Santiago, L. Briefman and J.H. Graham, 2011. Seasonal variability in HLB testing data in plant and psyllid samples in Florida. *Proceedings of the 2nd International Research Conference on Huanglongbing*, January 10-14, 2011, Orlando, FL., pp: 72.
13. Jagoueix, S., M.J. Bove and M. Garnier, 1996. PCR detection of the two *Candidatus liberobacter* species associated with greening disease of citrus. *Mol. Cell. Probes*, 10: 43-50.
14. Keremane, M.L., C. Ramadugu, A. Castaneda, J.E. Diaz and E.A. Penaranda *et al.*, 2015. Report of *Candidatus Liberibacter caribbeanus*, a new citrus-and psyllid-associated *Liberibacter* from Colombia, South America. *Proceedings of the APS Annual Meeting*, August 1-5, 2015, Pasadena, CA.
15. Lee, C., J. Kim, S.G. Shin and S. Hwang, 2006. Absolute and relative QPCR quantification of plasmid copy number in *Escherichia coli*. *J. Biotechnol.*, 123: 273-280.
16. Lopes, S.A., G.F. Frare, E. Bertolini, M. Cambra and N.G. Fernandes *et al.*, 2009. *Liberibacter* associated with citrus Huanglongbing in Brazil: *Candidatus Liberibacter asiaticus* is heat tolerant, *Ca. L. americanus* is heat sensitive. *Plant Dis.*, 93: 257-262.
17. Parker, J.K., S.R. Wisotsky, E.G. Johnson, F.M. Hijaz, N. Killiny, M.E. Hilf and L. De La Fuente, 2014. Viability of *Candidatus Liberibacter asiaticus* prolonged by addition of citrus juice to culture medium. *Phytopathology*, 104: 15-26.
18. Luo, X., A.L. Yen, K.S. Powell, F. Wu and Y. Wang *et al.*, 2015. Feeding behavior of *Diaphorina citri* (Hemiptera: Liviidae) and its acquisition of *Candidatus Liberibacter asiaticus*, on huanglongbing-infected *Citrus reticulata* leaves of several maturity stages. *Florida Entomol.*, 98: 186-192.
19. Manjunath, K.L., S.E. Halbert, C. Ramadugu, S. Webb and R.F. Lee, 2008. Detection of *Candidatus Liberibacter asiaticus* in *Diaphorina citri* and its importance in the management of citrus huanglongbing in Florida. *Phytopathology*, 98: 387-396.
20. SAS., 2009. *Statistical Analysis System for Windows*. Version 9, Statistical Analysis System, Cary, NC.
21. Razi, M.F., M.L. Keremane, C. Ramadugu, M. Roose, I.A. Khan and R.F. Lee, 2014. Detection of citrus huanglongbing-associated *Candidatus Liberibacter asiaticus* in citrus and *Diaphorina citri* in Pakistan, seasonal variability and implications for disease management. *Phytopathology*, 104: 257-268.
22. Robles, G.P., A.H. Sanchez and V.I. Delgadillo, 2009. Programa nacional para la deteccion del huanglongbing (*Candidatus Liberibacter spp.*) en Mexico. *Direccion General de Sanidad Vegetal*, SENASICA-SAGARPA.
23. Sauer, A.V., C.A. Zanutto, P.T.R. Nocchi, M.A. Machado, C.H. Bock and W.M. Nunes, 2015. Seasonal variation in populations of *Candidatus Liberibacter asiaticus* in citrus trees in Parana state, Brazil. *Plant Dis.*, 99: 1125-1132.
24. Tabachnick, W.J., 2015. *Diaphorina citri* (Hemiptera: Liviidae) vector competence for the citrus greening pathogen *Candidatus Liberibacter asiaticus*. *J. Econ. Entomol.*, 108: 839-848.
25. Trivedi, P., U.S. Sagaram, J.S. Kim, R.H. Brlansky and M.E. Rogers *et al.*, 2009. Quantification of viable *Candidatus Liberibacter asiaticus* in hosts using quantitative PCR with the aid of ethidium monoazide (EMA). *Eur. J. Plant Pathol.*, 124: 553-563.
26. Yamamoto, P.T., M.R. Felipe, L.F. Garbin, J.H.C. Coelho, E.C. Martins and A.P.R. Leite, 2006. *Diaphorina citri* (Kuwayama) (Hemiptera: Psyllidae): Vector of the bacterium *Candidatus Liberibacter americanus*. *Proceedings of the Huanglongbing-Greening International Workshop Fundecitrus*, July 16-20, 2006, Araguara, pp: 96.
27. Yan, Q., A. Sreedharan, S. Wei, J. Wang, K. Pelz Stelinski, S. Folimonova and N. Wang, 2013. Global gene expression changes in *Candidatus Liberibacter asiaticus* during the transmission in distinct hosts between plant and insect. *Mol. Plant Pathol.*, 14: 391-404.