

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

Pakistan Journal of Biological Sciences

ANSInet

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

Antimicrobial Activities of *Rhopalaea*-Associated Fungus *Aspergillus flavus* strain MFABU9

¹Deiske A. Sumilat, ¹Elvy L. Ginting, ²Gracia A.V. Pollo, ³Ahmad A. Adam and ⁴Trina E. Tallei

¹Department of Marine Science, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado, Indonesia

²Department of Biology, Faculty of Mathematics and Natural Sciences, Gadjah Mada University, Yogyakarta, Indonesia

³Department of Dental, Faculty of Medicine, Sam Ratulangi University, Manado, Indonesia

⁴Department of Biology, Faculty of Mathematics and Natural Sciences, Sam Ratulangi University, Manado, Indonesia

Abstract

Background and Objective: *Rhopalaea* is a genus of ascidian belonging to the family Diazonidae. Ascidians provide niches for various microorganisms including fungi. This present study describes the potential new source for natural bioactive compounds from *Rhopalaea*-associated fungi obtained from Bunaken marine park. **Materials and Methods:** As part of an on-going research program to explore the chemical diversity of marine derived fungi, we performed an antimicrobial bioactivity-guided screening of EtOAc extracts of the fungi isolated from ascidian *Rhopalaea* sp. **Results:** The study confirms that the ascidian obtained from Bunaken marine park was *Rhopalaea* sp. The fungus isolated from the ascidian was *Aspergillus flavus* which showed antimicrobial activity against bacteria *Escherichia coli*, *Staphylococcus aureus*, *Aeromonas hydrophila* and antifungal against the human pathogenic fungus *Candida albicans*. **Conclusion:** *Aspergillus flavus* isolated from ascidian *Rhopalaea* sp. has the potential as antibacterial and antifungal.

Key words: Ascidian, antimicrobial, *Aspergillus*, bioactive compound, marine derived-fungus, *Rhopalaea*

Citation: Deiske A. Sumilat, Elvy L. Ginting, Gracia A.V. Pollo, Ahmad A. Adam and Trina E. Tallei, 2020. Antimicrobial activities of *Rhopalaea*-associated fungus *Aspergillus flavus* strain MFABU9. Pak. J. Biol. Sci., 23: 911-916.

Corresponding Author: Deiske A. Sumilat, Department of Marine Science, Faculty of Fisheries and Marine Science, Sam Ratulangi University, Manado, Indonesia

Copyright: © 2020 Deiske A. Sumilat *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Marine-derived fungi are well recognised as a source of various novel metabolites, many of which possess valuable biological properties^{1,2} and pharmacological properties³. Marine-derived fungi which found in algae⁴, mangrove⁵, ascidians⁶ and sponges⁷ were shown to have antibacterial and anticancer activities.

Ascidians are found abundantly all over the world. They mostly live in shallow water with salinities over 2.5%⁸. However, they also can be found in the depths of the sea. Ascidian is a marine invertebrate animal and a member of the subphylum Tunicata. It productively produces a wide variety of secondary metabolites which are biologically and pharmacologically active⁹. These compounds have properties such as antibacterial¹⁰ and anticancer¹¹ activities which make them candidates for potential new drugs.

Correlations between ascidian microorganisms and ascidian metabolites are also being investigated, while the structures and synthetic pathways of a growing number of relevant compounds have been identified¹². Many of these metabolites are produced by the fungi isolated from ascidian^{9,13,14}. A number of new metabolites has been reported describing the fungal association with ascidians. As an example, an Indonesian ascidian is associated with *Penicillium verruculosum* which inhibited the activity of PTP 1B¹⁵ as well as *P. albobiverticillum*¹⁶. Menezes *et al.*¹⁷ analyzed in detail about fungal diversity in *Didemnum* spp.

Some compounds have been isolated from marine ascidians-derived fungi, which are taxonomically similar or identical to terrestrial fungi, such as *Aspergillus*¹⁸. The genus *Aspergillus*, which includes approximately 200 species, has been well studied and shown to produce many new metabolites¹⁹. Various fungal strains which predominated by *Aspergillus* and *Cladosporium* have been isolated from various Australian coral reefs²⁰. An ascidian-derived fungus *Aspergillus* sp. KMM 4676 exhibited cytotoxic activity as hormone therapy-sensitive human prostate cancer cells²¹. *Aspergillus candidus* isolated from colonial ascidian had a cytotoxic activity against hormone-sensitive line LNCaP²². Various kind of marine fungal strains has been reported to produce many kind of novel antimicrobial compounds. These compounds belong to alkaloids, macrolides, terpenoids, peptide derivatives as well as other types of structures²³.

Bunaken marine park in North Sulawesi (Indonesia) has been known for its various types of ascidians, among other is *Rhopalaea* sp. This type of ascidian is rarely investigated in association with the fungi and their antibacterial activities. In

this present study, *Rhopalaea* sp. from Bunaken marine park was studied to observe the possibility of its associated fungi for antibacterial activity.

MATERIALS AND METHODS

Sample preparation: Ascidian *Rhopalaea* sp. (Fig. 1) was collected at the Bunaken marine park in North Sulawesi, Indonesia, in March, 2019. The ascidian was rinsed with sterilized sea water and immersed in ethanol 70% for 1 min, then kept in the bottle vial and stored in the cool box and transported to the laboratory for further observation²⁴.

Isolation of ascidian-derived fungi from *Rhopalaea* sp.: A small piece of the sample was cut into cubes, washed with sterilized seawater and grown on PDA plate (BD, Franklin Lakes, NJ, the USA). The plate was incubated at 25°C for a week. The different appearing fungal colonies surrounding the samples that have different characteristics were isolated and grown on PDA. The isolation process was conducted for another round to obtain a pure single colony²⁴. One pure isolate designated as MFABU9 was chosen and grown on PDA. Afterwards, the isolate was inoculated into a 100-mL Erlenmeyer flask containing 50 mL sterilized sea water and 50 mg sterilized rice medium for 14 days.

EtOAc extraction of MFABU9 isolate: The 100 g of rice medium containing the selected growing fungus was extracted with EtOAc for 24 h. To complete the extraction process, the rice was incubated in 200 mL of EtOAc for 3 days at 25°C with constant shaking. The solvent was filtered using Whatman No.1 filter paper. The first filtrate was set aside in one clean Erlenmeyer and the remaining debris was soaked in 200 mL of EtOAc for 3 days at 25°C with constant shaking. The second and third filtrates were obtained using the same process. All of the filtrates were then combined and concentrated using a vacuum rotary evaporator at 40°C to obtain concentrated extract. The concentrated extract was then evaporated in the incubator to obtain a dry extract. The dry extract was weighted and stored in 20°C until used.

Screening for antimicrobial activity: The indicator pathogens used were bacterial strains *E. coli*, *S. aureus*, *Aeromonas hydrophila*, *Salmonella* sp., *Edwardsiella tarda* and *C. albicans*. Each of the bacterial inoculum *E. coli*, *S. aureus* and *A. Hydrophila* were cultured in liquid media B1 (peptone, meat extract, NaCl and aquadest) for 1×24 h, while *Salmonella* sp. and *E. tarda* were cultured in TSA media. Anti-



Fig. 1: Ascidian *Rhopalaea* sp. collected at Bunaken marine park in North Sulawesi, Indonesia

bacterial assay was carried out by the paper disc method using agar diffusion Kirby-Bauer methods, following the guidelines of Clinical and Laboratory Standard Institute (CLSI)²⁵.

The EtOAc extract of isolate MFABU9 was examined for the inhibitory activities against the indicator pathogenic bacteria and fungus with the concentrations of 20 µg/disk. The pathogenic bacteria were grown in nutrient agar (NA), while the indicator fungus was grown in sabouraud dextrose agar (SDA). Chloramphenicol was used as positive control for bacteria and Ketoconazole for fungus. For negative control, 40% CH₃OH was used. The plates were incubated at 37°C and the inhibitory activities were measured after 48 h of incubation. The results were interpreted by measuring the diameter of the inhibition zone using a caliper. The inhibition zone is a measure of the effectiveness of an active compound. Generally, a larger zone of inhibition means that the antimicrobial is more potent.

Molecular identification of isolate MFABU9: DNA extraction of the fungal isolate was carried out using Quick-DNA Fungal/Bacterial Miniprep Kit (Zymo Research, D6005). The ITS (internal transcribed spacer) region was amplified using primer pair ITS1 (F 5'-TCC GTA GGT GAA CCTGCG G-3') and ITS4 (R 5'-TCC TCC GCT TAT TGA TATGC-3') in the MyTaq HS Red Mix (Bioline, BIO-25047). The following PCR amplification conditions were used: one cycle of initial denaturation at 95°C for 5 min, followed by 35 cycles with a step of denaturation at 95°C for 30 sec, annealing at 55°C for 1 min and extension at 72°C for 1 min, followed by one cycle at 72°C for 6 min.

PCR products were purified using Zymoclean™ Gel DNA Recovery Kit (Zymo Research, D4001) and sent to the sequencing service provider. The sequencing result was

processed following the procedure performed by Tallei *et al.*²⁶ and subjected to BLAST (Basic Local Alignment Search Tool) search at NCBI (National Center for Biotechnology Information <https://www.ncbi.nlm.nih.gov/>), ISHAM (International Society for Human and Animal Mycology <https://www.isham.org/>) and BOLD (Barcode of Life Data System <http://www.boldsystems.org/>) and MycoBank (<http://mycobank.org>) for species identification.

RESULTS AND DISCUSSION

The fungal extract of isolate MFABU9 only showed antibacterial activity against *E. coli*, *S. aureus*, *A. hydrophila* and antifungal activity against *C. albicans* (Table 1). The extract, however, failed to suppress the growth of *E. tarda* and *Salmonella* sp. The isolate MFABU9 was molecularly identified as *Aspergillus flavus*, hence it is called *A. flavus* strain MFABU9. Inhibitory activity was defined by the diameter of inhibition zones. Inhibitory activity is defined as weak if the diameter is less than 10 mm. Inhibitory activity is defined as weak if the diameter is less than 10 mm, intermediate if the diameter ranges between 10-15 mm, and strong if the diameter is more than 16 mm. According to this, the extract of *A. flavus* strain MFABU9 showed a weak inhibitory activity against *S. aureus* and *C. albicans* and intermediate inhibitory activity against *E. coli* and *A. hydrophila*.

Only a little information about *A. flavus* isolated from ascidian that has been reported. Marine-derived *A. flavus* produced various kinds of secondary metabolites including mutagenic mycotoxins and other bioactive compounds²⁷. Ivanets *et al.*²¹ were able to isolate asperindoles A-D and a p-Terphenyl derivative from the ascidian-derived fungus *Aspergillus* sp. KMM 4676 which inhibited the growth of hormone therapy-resistant PC-3 and 22Rv1 and hormone therapy-sensitive human prostate cancer cells.

Some marine algal-derived *A. flavus* has been reported to produce bioactivity compounds. A cerebroside, an antibacterial cerebroside derivative isolated from *A. flavus* had antibacterial activity against *S. aureus*, methicillin-resistant *S. aureus* and multidrug-resistant *S. aureus*²⁸. Citrinadins A and B had been isolated from *A. flavus* which is associated with a green alga *Enteromorpha tubulosa*. This compound was cytotoxic to several tumor cell lines²⁹ HL-60, MOLT-4, A-549 and BEL-7402.

Ascidian has become a source of so many types of secondary metabolites. The resulting metabolites are used for physiological functions and specifically for defense mechanism against predators. Secondary metabolites which are synthesized by ascidians are not only synthesized by

Table 1: Antimicrobial assay of *A. flavus* strain MFABU9 extract

Indicator microbes	Inhibition zone (mm)
Indicator bacteria	
<i>Escherichia coli</i>	10.75
<i>Staphylococcus aureus</i>	9.00
<i>Edwardsiella tarda</i>	-
<i>Salmonella</i> sp.	-
<i>Aeromonas hydrophila</i>	11.45
Control+Chloramphenicol	25.00
Control	-
Indicator fungus	
<i>Candida albicans</i>	9.50
Control+Ketoconazole	25.00
Control	-

ascidians themselves, but also can be synthesized by the associated-microorganisms³⁰. Fungi are one of the microorganisms associated with ascidians. Most of the fungi can be found in the ascidian's tunic and several others can be found in the inside of the ascidian³¹.

Some types of fungi show specific relationships with ascidians. Fungi are involved in the ascidian's synthesis of bioactive secondary metabolites. However, the relationship between most fungi and specific ascidians is unclear. This indicates that the functional interactions between ascidians and fungi, especially in the interaction of *Rhopalaea* sp. and *A. flavus* are still unclear³².

Aspergillus is a genus of filamentous fungi. It has been widely used as a source of medicines³². Several species of the genus *Aspergillus* produce various types of compounds. Some of the compounds were aminobenzoic peptide seco-clavatustide B, clavatoic acid derivative 5-acetyl-2,4-dihydroxy-3-methyl-benzoic acid³³, clavatustide B³⁴, demethylsiderin, 3,7-dihydroxy-4-methylcoumarin and demethylkotanin³⁵.

Fungal species of the genus *Aspergillus* which could be found in water and associated with marine organisms, have many metabolites that have been proven to have the antibacterial and antifungal ability³⁶. The analysis of *A. flavus* extract showed that this fungus has antibacterial activity against *S. aureus*, *E. coli* and *A. hydrophila* and antifungal against *C. albicans*. This can be caused by the role of secondary metabolites of *A. flavus*. However, these antibacterial and antifungal activities did not only depend on certain types of synthesized compounds. These activities may be influenced by various metabolite compounds that play a synergistic or antagonistic role that support these activities³⁷. This is because the responses showed by the extract not only arises against fungi (*C. albicans*), but also against Gram-positive (*S. aureus*) and Gram-negative bacteria (*E. coli* and *A. hydrophila*).

Besides the synergistic and antagonistic roles, specific types of synthesized compounds may also perform different antibacterial mode of actions to specific types of bacteria. It is because of different types of bacterial cells, may only be affected by the specific mode of action³⁸. It can be implied that *A. flavus* may synthesize compounds with antibacterial activity against *S. aureus*, *E. coli* and *A. hydrophila* with specific reaction and mode of action.

CONCLUSION

The results obtained from the present study confirmed that *A. flavus* strain MFABU9 isolated from ascidian *Rhopalaea* sp. showed weak to intermediate inhibitory activities against selected indicator pathogenic microorganisms. Further studies are needed to be conducted to isolate each of bioactive compounds from this strain and test for their antibacterial and anticancer activities. The data obtained in this study have opened of the importance of a preliminary screening study. The molecular docking can be conducted after bioactive compounds from this strain have been elucidated.

SIGNIFICANCE STATEMENT

This study discovered that fungus *A. flavus* isolated from ascidian *Rhopalaea* sp. obtained from the Bunaken marine park showed antibacterial activities against *E. coli*, *S. aureus* and *A. hydrophila* and antifungal activity against *C. albicans*. However, it is advisable to explore further the bioactive compounds of this fungus which can be used as antimicrobial agents.

ACKNOWLEDGMENT

This study was financed by the Directorate of Research and Community Service, Ministry of Research, Technology and Higher Education, Republic of Indonesia, through Basic Research Sceme fiscal year 2019 (Grant Number:127/UN12.13/LT/2019).

REFERENCES

1. Deshmukh, S.K., V. Prakash and N. Ranjan, 2018. Marine fungi: A source of potential anticancer compounds. Front. Microbiol., Vol. 8. 10.3389/fmicb.2017.02536.
2. Butler, M.S., A.A.B. Robertson and M.A. Cooper, 2014. Natural product and natural product derived drugs in clinical trials. Nat. Prod. Rep., 31: 1612-1661.

3. Ebel, R., 2010. Terpenes from marine-derived fungi. Mar. Drugs, 8: 2340-2368.
4. Handayani, D., N. Ananda, M.A. Artasasta, R. Ruslan, O. Fadriyanti and T.E. Tallei, 2019. Antimicrobial activity screening of endophytic fungi extracts isolated from brown algae *Padina* sp. J. Applied Pharmaceut. Sci., 9: 9-13.
5. Handayani, D., N. Sandrawati, R. Ruslan, O. Nestianda, A. Fajrina and T.E. Tallei, 2019. Cytotoxic and antimicrobial activities of ethyl acetate extract of mangrove plant *Scyphiphora hydrophyllacea* C. F. Gaertn-associated fungi. J. Applied Pharmaceut. Sci., 9: 75-79.
6. Chen, L., J.S. Hu, J.L. Xu, C.L. Shao and G.Y. Wang, 2018. Biological and chemical diversity of ascidian-associated microorganisms. Mar. Drugs, Vol. 16, No. 10. 10.3390/md 16100362.
7. Handayani, D. and M.A. Artasasta, 2017. Antibacterial and cytotoxic activities screening of symbiotic fungi extract isolated from marine sponge *Neopetrosia chaliniformis* AR-01. J. Applied Pharmaceut. Sci., 7: 66-69.
8. Saha, A., R. Yadav and N. Rajendran, 2014. Biomaterials from sponges, ascidians and other marine organisms. Int. J. Pharmaceut. Sci. Rev. Res., 27: 100-109.
9. Watters, D.J., 2018. Ascidian toxins with potential for drug development. Mar. Drugs, Vol. 16, No. 5. 10.3390/md 16050162.
10. Adam, A.A., J. Posangi, E. Tumewu and T.E. Tallei, 2014. Aktivitas antibakteri ekstrak kasar tunikata *Polycarpa aurata* terhadap *Streptococcus mutans*. Dentire, 3: 118-122.
11. Chandramathi, V., R.D. Thilaga and V. Sivakumar, 2011. Antibacterial activity of methanol extract of colonial ascidian *Distaplia nathensis*. Global J. Biotechnol. Biochem., 6: 183-185.
12. Chen, L., C. Fu and G. Wang, 2017. Microbial diversity associated with ascidians: A review of research methods and application. Symbiosis, 71: 19-26.
13. Sumilat, D.A., D.S. Wewengkang, H. Rotinsulu, H. Yamazaki, T. Oda, K. Ukai and M. Namikoshi, 2018. Bioactivity of extracts from ascidians collected in North Sulawesi as seeds of marine-derived drugs. Aquacult. Aquarium Conserv. Legisl., 11: 516-524.
14. Sumilat, D.A., J.R. Rimper, E.T. Opa and D. Kurnia, 2019. The potential of marine ascidians as sources of natural antioxidant and antibacterial agents from Manado, North Sulawesi. Aquacult. Aquarium Conserv. Legisl., 12: 373-377.
15. Yamazaki, H., W. Nakayama, O. Takahashi, R. Kirikoshi and Y. Izumikawa *et al.*, 2015. Verruculides A and B, two new protein tyrosine phosphatase 1B inhibitors from an Indonesian ascidian-derived *Penicillium verruculosum*. Bioorg. Med. Chem. Lett., 25: 3067-3090.
16. Sumilat, D.A., H. Yamazaki, K. Endo, H. Rotinsulu, D.S. Wewengkang, K. Ukai and M. Namikoshi, 2017. A new biphenyl ether derivative produced by Indonesian ascidian-derived *Penicillium albobiverticillium*. J. Natural Med., 71: 776-779.
17. Menezes, C.B.A., R.C. Bonugli-Santos, P.B. Miqueletto, M.R.Z. Passarini and C.H.D. Silva *et al.*, 2010. Microbial diversity associated with algae, ascidians and sponges from the North Coast of Sao Paulo State, Brazil. Microbiol. Res., 165: 466-482.
18. Namikoshi, M. and J.Z. Xu, 2009. Filamentous Fungi in the Marine Environment: Chemical Ecology. In: Fungi from Different Environments, Misra, J.K. and S.K. Deshmukh (Eds.). Taylor and Francis, London, UK, ISBN-13: 9781578085781, pp: 81-112.
19. Carroll, A.R., B.R. Copp, R.A. Davis, R.A. Keyzers and M.R. Prinsep, 2019. Marine natural products. Nat. Prod. Rep., 36: 122-173.
20. Morrison-Gardiner, S., 2002. Dominant fungi from Australian coral reefs. Fungal Divers, 9: 105-121.
21. Ivanets, E.V., A.N. Yurchenko, O.F. Smetanina, A.B. Rasin and O.I. Zhuravleva *et al.*, 2018. Asperindoles A-D and a ρ -terphenyl derivative from the ascidian-derived fungus *Aspergillus* sp. KMM 4676. Mar. Drugs, Vol. 16, No. 7. 10.3390/ md16070232.
22. Yurchenko, A.N., E.V. Ivanets, O.F. Smetanina, M.V. Pivkin, S.A. Dyshlovoi, G. von Amsberg and S.S. Afiyatullov, 2017. Metabolites of the marine fungus *Aspergillus candidus* KMM 4676 associated with a kuril colonial ascidian. Chem. Nat. Compd., 53: 747-749.
23. Zhang, Y., J. Mu, Y. Feng, Y. Kang and J. Zhang *et al.*, 2009. Broad-spectrum antimicrobial epiphytic and endophytic fungi from marine organisms: Isolation, bioassay and taxonomy. Mar. Drugs, 7: 97-112.
24. Kjer, J., A. Debbab, A.H. Aly and P. Proksch, 2010. Methods for isolation of marine-derived endophytic fungi and their bioactive secondary products. Nat. Protocols, 5: 479-490.
25. CLSI., 2019. Performance standards for antimicrobial susceptibility testing. CLSI Supplement M100, 29th Edition, Clinical and Laboratory Standards Institute, Wayne, PA., USA., January 2019.
26. Tallei, T.E. and B.J. Kolondam, 2015. DNA barcoding of Sangihe Nutmeg (*Myristica fragrans*) using *matK* gene. HAYATI J. Biosci., 22: 41-47.
27. Lamboni, Y., K.F. Nielsen, A.R. Linnemann, Y. Gezgin and K. Hell *et al.*, 2016. Diversity in secondary metabolites including mycotoxins from strains of *Aspergillus* section *Nigri* isolated from raw cashew nuts from Benin, West Africa. PLoS ONE, Vol. 11. 10.1371/journal.pone.0164310.
28. Yang, G., L. Sandjo, K. Yun, A.S. Leutou and G.D. Kim *et al.*, 2011. Flavusides A and B, antibacterial cerebrosides from the marine-derived fungus *Aspergillus flavus*. Chem. Pharmaceut. Bull., 59: 1174-1177.
29. Lin, A.Q., L. Du, Y.C. Fang, F.Z. Wang, T.J. Zhu, Q.Q. Gu and W.M. Zhu, 2009. *iso-* α -cyclopiazonic acid, a new natural product isolated from the marine-derived fungus *Aspergillus flavus* CF-3. Chem. Nat. Compd., 45: 677-680.

30. Karim, F., M.Y. Putra, T.A. Hadi and M. Abrar, 2018. Antimicrobial and cytotoxic properties of the ascidians *Lissoclinum patella*, *Oxycyrrina fascicularis*, *Didemnum molle* and *Botryllus schlosseri*. *Pharmaceut. Sci. Res.*, 5: 65-71.
31. Lopez-Legentil, S., P.M. Erwin, M. Turon and O. Yarden, 2015. Diversity of fungi isolated from three temperate ascidians. *Symbiosis*, 66: 99-106.
32. Sanchez, J.F., A.D. Somoza, N.P. Keller and C.C.C. Wang, 2012. Advances in *Aspergillus* secondary metabolite research in the post-genomic era. *Nat. Prod. Rep.*, 29: 351-371.
33. Song, Q., X.M. Li, X.Y. Hu, X. Li, L.P. Chi, H.L. Li and B.G. Wang, 2019. Antibacterial metabolites from Ascidian-derived fungus *Aspergillus clavatus* AS-107. *Phytochem. Lett.*, 34: 30-34.
34. Jiang, W., P. Ye, C.T.A. Chen, K. Wang and P. Liu *et al.*, 2013. Two novel hepatocellular carcinoma cycle inhibitory cyclodepsipeptides from a hydrothermal vent crab-associated fungus *Aspergillus clavatus* C2WU. *Mar. Drugs*, 11: 4761-4772.
35. Huttel, W. and M. Muller, 2007. Regio-and stereoselective intermolecular oxidative phenol coupling in kotanin biosynthesis by *Aspergillus niger*. *ChemBioChem*, 8: 521-529.
36. Bajpai, V.K., 2016. Antimicrobial secondary metabolites from marine fungi: A mini review. *Indian J. Geo-Mar. Sci.*, 45: 1067-1075.
37. Ruangamnart, A., S. Buranaphalin, R. Temsiririrkkul, W. Chuakul and J. Pratuangdejkul, 2015. Chemical compositions and antibacterial activity of essential oil from dill fruits (*Anethum graveolens* L.) cultivated in Thailand. *Mahidol Univ. J. Pharmaceut. Sci.*, 42: 135-143.
38. Kapoor, G., S. Saigal and A. Elongavan, 2017. Action and resistance mechanisms of antibiotics: A guide for clinicians. *J. Anaesthesiol. Clin. Pharmacol.*, 33: 300-305.