

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

Pakistan Journal of Biological Sciences

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

The Metagenomic Analysis of Potential Pathogenic Emerging Bacteria in Fleas

¹Dina Victoria Rombot and ^{2,3}Mokosuli Yermia Samuel

¹Faculty of Medicine, Sam Ratulangi University, Manado, North Sulawesi, Indonesia

²Department of Biology, Faculty of Mathematics and Natural Sciences, Manado State University, Tondano, North Sulawesi, Indonesia

³Laboratory of Biomolecular and Bioactivity, Faculty of Mathematics and Natural Sciences, Manado State University, Tondano, North Sulawesi, Indonesia

Abstract

Background and Objective: At present many pathogenic microbes that cause disease in humans are transmitted through animals. *Ctenocephalides felis* is specific ectoparasites in cats. Metagenomic research on the digestive tract and body surface of *C. felis* has been conducted. DNA genomics was extracted from the body surface and digestive tract of *C. felis*. **Materials and Methods:** Metagenomic analysis has used the 16S rRNA gene (V3-V4 region). Sequencing was carried out using New Generation Sequencing at the First BASE Laboratory, Singapore. **Results:** Wolbachia has the most significant bacterial composition in *C. felis* (94.4%), we were found bacteria with a composition >1% that have never been reported to be associated with *C. felis*. Also, there were 0.2% of bacteria whose taxonomic status cannot be determined. **Conclusion:** The results of this study become a vital reference pathogenic bacteria that can be transmitted to humans and animals through *C. felis*. It is necessary to study the resistance of bacteria isolated from *C. felis* to antibiotics in the future.

Key words: Metagenomics, 16S rRNA, *Ctenocephalides felis*, bacteria composition, symbiotic, caulobacteriaceae, staphylococcaceae

Citation: Rombot, D.V. and M.Y. Samuel, 2021. The metagenomic analysis of potential pathogenic emerging bacteria in fleas. Pak. J. Biol. Sci., 24: 1084-1090.

Corresponding Author: Mokosuli Yermia Samuel, Department of Biology, State University of Manado, Tondano, Sulawesi Utara, Indonesia

Copyright: © 2021 Dina Victoria Rombot *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Ctenocephalides felis is the most widely distributed ectoparasite worldwide^{1,2}. This is due to human migration which also brings domestic cats to various regions around the world^{3,4}. Insects have carried out symbiotic coevolution with various types of bacteria. Metagenomic analysis has made it possible to examine the diversity of symbiotic and non-symbiotic bacteria in insects, especially species of bacteria that cannot be cultured in a laboratory^{5,6}. However, the genome and behaviour of these bacteria are the object of very deep biological studies and holds a mystery. Metagenomic exploration in insects has been carried out lately. Several insects have been performed metagenomic analysis including Apids⁷, *Megaphragma amalphitanum*⁸, blowflies and house flies^{9,10}, *Dendroctonus valens* and *D. mexicanus*¹¹. All metagenomic research results on these insects are reported to be genera and bacterial species that have never been known to be associated with insects before.

However, the composition of bacteria based on genus and family needs to be mapped to determine the abundance of bacteria at the family and genus levels found in cat fleas. Recently, pathogenic microbes that come from animals and infect humans have become a topic of much research^{6,9}. Identifying bacteria and the composition of bacteria in cat fleas is important to study, among others, the background that cats are the domestic pets in the world. As a pet, cats interact directly with humans, so the possibility of transmitting pathogenic microbes to humans is substantial¹⁰⁻¹².

Previous research found several families of potential pathogens in humans. These families include Rickettsiaceae, Burkholderiaceae, Pseudomonadaceae, Planococcaceae, Corynebacteriaceae, Caulobacteriaceae, Isosphaeraceae and Staphylococcaceae¹³. Bacterial identification using a metagenomic approach will identify 99% of the bacteria present in samples^{7,13,14}. For the identification of bacteria in this study using the 16S RNA gene. Bacterial identification using the 16S RNA gene is still empathetic to identify bacteria¹⁵⁻¹⁸.

However, the composition of bacteria at the genus and family level needs to be studied in depth. This study describes the bacterial composition of *C. felis* at the family and genus levels using a metagenomic analysis approach.

MATERIALS AND METHODS

Study area: Samples of *C. felis* were obtained from wild cats in several locations in Manado City, namely Karombasan



Fig. 1: *Ctenocephalides felis* samples from Manado North Sulawesi, Indonesia

Village, Paal Dua Village and Malalayang Village in Manado City, North Sulawesi, Indonesia. The blackish-brown cat fleas, complete with their organs, were then taken to the laboratory for analysis (Fig. 1). Morphological identification was carried out in the Biology Laboratory of Manado State University, Indonesia. Samples were prepared with 70% alcohol before being used for total DNA genomics extraction. Complete DNA extraction was carried out at the Laboratory of Bioactivity and Molecular Biology, Manado State University, North Sulawesi, Indonesia, from July, 2019 until March, 2020. Metagenomic sequencing was carried out at Axil Scientific Pte Ltd Laboratory (20-0200922-D) 41 Science Park Road, #04-08, The Gemini, Singapore Science Park II, Singapore. The study was carried out at the Department of Biology, Bimolecular Lab, Indonesia from July, 2019-March, 2020.

Extraction of DNA genomic: Total DNA *C. felis* was extracted using the Bacteria Genomic DNA Kit with the manufacturer's protocol. Total DNA *C. felis* was extracted using the Bacteria Genomic DNA Kit with the manufacturer's protocol and with conventional methods (CTAB, Cetyltrimethylammonium Bromide)^{19,20}. To determine the concentration and purity of DNA, the 1% gel electrophoresis method was used where based on the concentration, the DNA was diluted to 1 ng L⁻¹ using dd H₂O.

Amplicon generation: 16S rRNA genes of distinct regions 16S V3-V4 were amplified used a specific primer with the barcode. The primer was used as follow Table 1.

PCR products

Quantification and qualification: To identify the PCR product, mix equal quantities of 1X loading buffer (including SYB




Fig. 2: Metagenomic research workflow

Table 1: Primer sequence used in the research

Hypervariable region	Primer	References
V3	28F: 5'-GAGTTTGATCCTGGCTCAG-3'	Suchodolski <i>et al.</i> ²¹
	519R: 5'-GTNTTACNGCGGCKGCTG-3'	Turner <i>et al.</i> ²²
V4	515F: 5'-GTGCCAGCMGCCGCGG-3'	Weisburg <i>et al.</i> ²³
	907R: 5'-CCGTCAATTCMTTTRAGTTT-3'	

All PCR reactions were carried out with Phusion® High-Fidelity PCR master mix (New England Biolabs)

green) and electrophoresis on a 2% agarose gel. Further studies were conducted on samples with bright main strips ranging from 400-450 bp. TapeStation 4200, picogreen and nanodrop were used to test the quality and quantity of the V3V4 amplicon. All of the samples passed the quality-control tests and were sent straight to production line²¹.

Mixing and purification: PCR products were mixed in equidensity ratios. Then, a mixture of PCR products was purified with Qiagen Gel Extraction Kit (Qiagen, Germany). The libraries generated with TruSeq® DNA PCR-Free Sample Preparation Kit and quantified via Qubit and Q-PCR would be analyzed by HiSeq2500 PE250 (Fig. 2).

The quality of the libraries was measured using TapeStation4200, Picogreen and qPCR. All libraries passed the QC measurement. The library was then pooled according to the protocol recommended by Illumina and proceed straight to sequencing using the MiSeq platform at 2x301PE format. The libraries were prepared using Illumina 16s metagenomics library prep kit and their quality and quantity were determined using Agilent TapeStation 4200 and Picogreen.

New generation sequencing of DNA: The amplicon was sequenced on the Illumina HiSeq paired-end platform, yielding 250 bp paired-end raw reads (Raw PE), assembled and processed to get Clean Tags. Raw data would be combined and filtered to provide clean data for sequencing data analysis. The effective data is utilized to create OTU clusters and species annotations for each OTU's sequence. As a result, the relativistic²⁴.

RESULTS AND DISCUSSION

Family level: The composition of the family level results of metagenomic analysis based on the 16S rRNA gene from *C. felis* found 0.2% of the population of Unassigned bacteria,

which means it has not been recorded or identified in the world bacterial taxonomic system. The bacterial database based on the 16S rRNA gene has been stored in the NCBI gene bank. As many as 0.1% belong to the family Corynebacteriaceae, 0.2% included in the Bacillales family, 1.8% included in the family Plannococcaceae, 0% included in the Staphylococcus, 0.1% belongs to the family Methylophilaceae, 0.1% belong to the family Neisseriaceae, 0.1% belong to the Moraxellaceae family and 96.4% belong to the Rickettsiaceae family (Table 2). The Rickettsiaceae family is the dominant bacterial family of bacteria on the surface of the body and the digestive tract of *C. felis* cat fleas.

Genus level: The composition of genus-level resulted from metagenomic analysis based on 16S rRNA, *C. felis* genes was found in 0.2% of the population of unassigned bacteria, which means it has not been recorded or identified in the world bacterial taxonomic system. The bacterial database based on the 16S rRNA gene has been stored in the NCBI gene bank. As much as 0.1% belongs to the genus Corynebacterium, 0.2% included in the genus Bacillales, 1.8% belong to the family genus Plannococcaceae not yet known, 0.8% belongs to the genus Staphylococcus, 0.1% belongs to the family of the genus Methylophilaceae unknown, 0.1% belongs to the family of the genus Neisseriaceae unknown, 0.1% belong to the family Moraxellaceae genus unknown, 1.5% belong to the genus Rickettsia and 94.4% to the genus Wolbachia (Table 3). The family Rickettsiaceae genus Wolbachia is the dominant bacterium on the body's surface and the digestive tract of *C. felis*.

The results of this study indicate that a new family and genus of bacteria were first reported in cat fleas. The genus has many pathogenic species in humans, namely the genus Staphylococcus, Corynebacterium and the family Plannococcaceae. It is necessary to study more deeply the potential of *C. felis* to transmit other pathogenic bacteria to humans.

Table 2: Composition of bacteria at the family level

Numbers	Genus	Percentage
1	Corynebacteriaceae	0.1
2	Bacillales	0.2
3	Planococcaceae	1.8
4	Staphylococcus	0.0
5	Isosphaeraceae	0.3
6	Rickettsiaceae	0.5
7	Rickettsiae	96.4
8	Methylophilaceae	0.1
9	Neisseriaceae	0.1
10	Moraxellaceae	0.1

Complete composition is shown in Appendix 1

Legend	Taxonomy	Total %	RN %
	Unassigned;Other;Other;Other;Other	0.1%	0.2%
	k_Archaea;p_Euryarchaeota;c_Methanobacteria;o_Methanobacteriales;f_Methanobacteriaceae	0.0%	0.0%
	k_Bacteria;p_Acidobacteria;c_Acidobacteria;o_iii1-15;f_	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Actinopolysporaceae	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Corynebacteriaceae	12.3%	0.1%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Dermatophilaceae	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Intrasporangiaceae	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Micrococcaceae	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Nocardioidaceae	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Pseudonocardiaceae	0.0%	0.0%
	k_Bacteria;p_Actinobacteria;c_Coriobacteriia;o_Coriobacteriales;f_Coriobacteriaceae	0.0%	0.0%
	k_Bacteria;p_Bacteroidetes;c_Flavobacteriia;o_Flavobacteriales;f_Weeksellaceae	0.0%	0.0%
	k_Bacteria;p_Chloroflexi;c_Thermomicrobia;o_	0.0%	0.0%
	k_Bacteria;p_Chloroflexi;c_Thermomicrobia;o_JG30-KF-CM45;f_	0.0%	0.0%
	k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;Other	0.1%	0.2%
	k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Bacillaceae	0.0%	0.0%
	k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Planococcaceae	4.7%	1.8%
	k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Staphylococcaceae	0.4%	0.6%
	k_Bacteria;p_Firmicutes;c_Clostridia;o_Clostridiales;Other	0.0%	0.0%
	k_Bacteria;p_Firmicutes;c_Clostridia;o_Clostridiales;f_Lachnospiraceae	0.0%	0.0%
	k_Bacteria;p_Firmicutes;c_Clostridia;o_Clostridiales;f_Ruminococcaceae	0.0%	0.0%
	k_Bacteria;p_Planctomycetes;c_Planctomycetia;o_Gemmatales;f_Isosphaeraceae	0.1%	0.3%
	k_Bacteria;p_Planctomycetes;c_Planctomycetia;o_Pirellulales;f_Pirellulaceae	0.0%	0.0%
	k_Bacteria;p_Planctomycetes;c_Planctomycetia;o_Planctomycetales;f_Planctomycetaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Caulobacteriales;f_Caulobacteraceae	0.9%	0.0%
	k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rhizobiales;f_Bartonellaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rhizobiales;f_Bradyrhizobiaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rhodospirillales;f_Acetobacteraceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rickettsiales;f_Rickettsiaceae	48.2%	96.4%
	k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Sphingomonadales;f_Sphingomonadaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Burkholderiales;f_Burkholderiaceae	32.5%	0.0%
	k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Burkholderiales;f_Comamonadaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Methylophilales;f_Methylophilaceae	0.0%	0.1%
	k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Neisseriales;f_Neisseriaceae	0.1%	0.1%
	k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;Other;Other	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Aeromonadales;f_Aeromonadaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Enterobacteriales;f_Enterobacteriaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pasteurellales;f_Pasteurellaceae	0.0%	0.0%
	k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pseudomonadales;f_Moraxellaceae	0.0%	0.1%
	k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pseudomonadales;f_Pseudomonadaceae	0.3%	0.0%
	k_Bacteria;p_Thermi;c_Deinococci;o_Deinococcales;f_Deinococcaceae	0.0%	0.0%

Appendix 1: Family composition of bacteria from *C. felis*

Table 3: Composition of bacteria at the genus level

Numbers	Genus	Percentage
1	Corynebacterium	0.1
2	Bacillales	0.2
3	Planococcaceae-genus unknown	1.8
4	Staphylococcus	0.8
5	Isosphaeraceae genus unknown	0.3
6	Rickettsiaceae genus unknown	0.5
7	Rickettsia	1.5
8	Wolbachia	94.4
9	Methylophilaceae genus unknown	0.1
10	Neisseriaceae genus unknown	0.1
11	Acinetobacter	0.1

Complete composition is shown in Appendix 2

Legend	Taxonomy	Total %	RN %
Unassigned;Other;Other;Other;Other;Other		0.1%	0.2%
k_Archaea;p_Euryarchaeota;c_Methanobacteria;o_Methanobacteriales;f_Methanobacteriaceae;g_Methanobacterium		0.0%	0.0%
k_Bacteria;p_Acidobacteria;c_Acidobacteria-6;o_iii1-15;f_g		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Actinopolysporaceae;Other		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Corynebacteriaceae;g_Corynebacterium		12.3%	0.1%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Dermatophilaceae;Other		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Intrasporangiaceae;g		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Micrococcaceae;g		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Micrococcaceae;g_Kocuria		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Micrococcaceae;g_Rothia		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Nocardioidaceae;g		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Actinobacteria;o_Actinomycetales;f_Pseudonocardiaceae;g_Saccharopolyspora		0.0%	0.0%
k_Bacteria;p_Actinobacteria;c_Coriobacteriia;o_Coriobacteriales;f_Coriobacteriaceae;g_Collinsella		0.0%	0.0%
k_Bacteria;p_Bacteroidetes;c_Flavobacteriia;o_Flavobacteriales;f_[Weeksellaceae];g_Cloacibacterium		0.0%	0.0%
k_Bacteria;p_Chloroflexi;c_Thermomicrobia;o_f_g		0.0%	0.0%
k_Bacteria;p_Chloroflexi;c_Thermomicrobia;o_JG30-KF-CM45;f_g		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;Other;Other		0.1%	0.2%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Bacillaceae;g_Bacillus		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Planococcaceae;Other		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Planococcaceae;g		4.7%	1.8%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Staphylococcaceae;g_Macrococcus		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Staphylococcaceae;g_Salinicoccus		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Bacilli;o_Bacillales;f_Staphylococcaceae;g_Staphylococcus		0.4%	0.6%
k_Bacteria;p_Firmicutes;c_Clostridia;o_Clostridiales;Other;Other		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Clostridia;o_Clostridiales;f_Lachnospiraceae;g		0.0%	0.0%
k_Bacteria;p_Firmicutes;c_Clostridia;o_Clostridiales;f_Ruminococcaceae;g		0.0%	0.0%
k_Bacteria;p_Plantomycetes;c_Plantomycetia;o_Gemmatales;f_Isosphaeraceae;g		0.1%	0.3%
k_Bacteria;p_Plantomycetes;c_Plantomycetia;o_Pirellulales;f_Pirellulaceae;g		0.0%	0.0%
k_Bacteria;p_Plantomycetes;c_Plantomycetia;o_Plantomycetales;f_Plantomycetaceae;g_Plantomycetes		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Caulobacteriales;f_Caulobacteraceae;g		0.9%	0.0%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rhizobiales;f_Bartonellaceae;g_Bartonella		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rhizobiales;f_Bradyrhizobiaceae;g		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rhodospirillales;f_Acetobacteraceae;g		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rickettsiales;f_Rickettsiaceae;Other		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rickettsiales;f_Rickettsiaceae;g		0.2%	0.5%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rickettsiales;f_Rickettsiaceae;g_Rickettsia		0.8%	1.5%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Rickettsiales;f_Rickettsiaceae;g_Wolbachia		47.2%	94.4%
k_Bacteria;p_Proteobacteria;c_Alphaproteobacteria;o_Sphingomonadales;f_Sphingomonadaceae;g_Kaistobacter		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Burkholderiales;f_Burkholderiaceae;g_Burkholderia		32.5%	0.0%
k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Burkholderiales;f_Comamonadaceae;g_Delftia		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Methylophilales;f_Methylophilaceae;g		0.0%	0.1%
k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Neisseriales;f_Neisseriaceae;g		0.1%	0.1%
k_Bacteria;p_Proteobacteria;c_Betaproteobacteria;o_Neisseriales;f_Neisseriaceae;g_Neisseria		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;Other;Other		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Aeromonadales;f_Aeromonadaceae;g		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Enterobacteriales;f_Enterobacteriaceae;Other		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Enterobacteriales;f_Enterobacteriaceae;g		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pasteurellales;f_Pasteurellaceae;g_Actinobacillus		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pseudomonadales;f_Moraxellaceae;g_Acinetobacter		0.0%	0.1%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pseudomonadales;f_Pseudomonadaceae;g		0.0%	0.0%
k_Bacteria;p_Proteobacteria;c_Gammaproteobacteria;o_Pseudomonadales;f_Pseudomonadaceae;g_Pseudomonas		0.3%	0.0%
k_Bacteria;p_[Thermi];c_Deinococci;o_Deinococcales;f_Deinococcaceae;g_Deinococcus		0.0%	0.0%

Appendix 2: Genus composition of bacteria from *C. felis*

Although the *Wolbachia* genus has the most total bacterial composition in *C. felis* (94.4%), there are bacteria with a composition >1% that have never been reported to be associated with *C. felis*. Furthermore, there are 0.2% of bacteria whose taxonomy has yet to be determined. The results of this study become a vital reference pathogenic bacteria that can be transmitted in humans and animals through *C. felis*. It is necessary to study the resistance of bacteria isolated from *C. felis* to antibiotics in the future.

This study confirms some bacteria associated with cat fleas such as *Bartonella*, *Rickettsia* and *Wolbachia*²⁵. *Rickettsia* and *Bartonella* infections occur worldwide and may cause serious diseases in people^{19,26,27}. Bacteria of the genera *Staphylococcus* and *Streptococcus* are found in cat fleas²⁸. However, in this study, it was found that the composition of the genus *Rickettsia* bacteria was the bacteria with the largest composition. *Rickettsia asembonensis* and *R. felis* were reported to be human pathogens²⁹. The dominance of the genus *Rickettsia* in cat fleas was also reported by Douglas *et al.*³⁰, Billeter *et al.*³¹, Roucher *et al.*³². The bacterial species found in *C. felis* in this study are medically relevant bacteria (MR) in humans, based on the definition of the International Statistical Classification of Diseases and Related Health Problems, WHO³³. This study proposes a recent study of the bacterial composition of *C. felis* with a metagenomic approach. This is the basis for the study of bacteria that have the potential to infect humans in the future. However, further research is needed to identify bacteria in *C. felis* using marker genes other than 16S rRNA.

CONCLUSION

The results of metagenomic analysis of bacteria from *C. felis* found many species of bacteria that conventional bacterial isolation methods can not isolate. Species of bacteria are found to be potentially infecting humans because they belong to medically relevant bacteria. The dominant bacterial genus in cat fleas is *Wolbachia*. Found a genus that has never been reported found in *C. felis*, where this genus has many species of pathogens in humans.

SIGNIFICANCE STATEMENT

This study found the composition of the bacterial genus associated with *C. felis* which could be useful for the study of potential bacterial pathogens not only in cats but in humans in the future. This study will help researchers to uncover critical areas of metagenomic bacteria associated with *C. felis* that many other researchers have not explored. Thus, this

study provides scientific information about the composition of *C. felis* bacteria through a metagenomic study approach.

ACKNOWLEDGMENT

We want to thank the chairperson and staff of the bioactivity and molecular biology laboratory, Manado State University, for their assistance in metagenomic analysis.

REFERENCES

1. Mendes-de-Almeida, F., A.L. Crissiuma, L.C. Gershony, L.M.V. Willi, J.P. Paiva, J. Guerrero and N. Labarthe, 2011. Characterization of ectoparasites in an urban cat (*Felis catus* Linnaeus, 1758) population of Rio de Janeiro, Brazil. *Parasitol. Res.*, 108: 1431-1435.
2. Pérez-Orsorio, C.E., J.E. Zavala-Velázquez, J.J.A. León and J.E. Zavala-Castro, 2008. *Rickettsia felis* as emergent global threat for humans. *Emerging Infect. Dis.*, 14: 1019-1023.
3. Chamany, K., D. Allen and K. Tanner, 2008. Making biology learning relevant to students: Integrating people, history and context into college biology teaching. *CBE—Life Sci. Educ.*, 7: 267-278.
4. Crkvenec, N. and J. Šlapeta, 2019. Climate change models predict southerly shift of the cat flea (*Ctenocephalides felis*) distribution in Australia. *Parasites Vectors*, Vol. 12. 10.1186/s13071-019-3399-6.
5. Oliveira, C., L. Gunderman, C. Coles, J. Lochmann and M. Parks *et al.*, 2017. 16S rRNA gene-based metagenomic analysis of ozark cave bacteria. *Diversity*, Vol. 9. 10.3390/d9030031.
6. Simandjuntak, S. and M. Samuel, 2018. Isolation and identification of thermophilic bacteria, producer of amylase enzyme, from lake Linow, North Sulawesi. *J. Pure Appl. Microbiol.*, 12: 543-554.
7. Clerck, C.D., A. Fujiwara, P. Joncour, S. Léonard and M.L. Félix *et al.*, 2015. A metagenomic approach from aphid's hemolymph sheds light on the potential roles of co-existing endosymbionts. *Microbiome*, Vol. 3. 10.1186/s40168-015-0130-5.
8. Nedoluzhko, A.V., F.S. Sharko, S.V. Tsygankova, E.S. Boulygina and A.S. Sokolov *et al.*, 2017. Metagenomic analysis of microbial community of a parasitoid wasp *Megaphragma amalphantum*. *Genomics Data*, 11: 87-88.
9. Junqueira, A.C.M., A. Ratan, E. Acerbi, D.I. Drautz-Moses and B.N.V. Premkrishnan *et al.*, 2017. The microbiomes of blowflies and houseflies as bacterial transmission reservoirs. *Sci. Rep.*, Vol. 7. 10.1038/s41598-017-16353-x.
10. Singh, K.M., V.B. Ahir, A.K. Tripathi, U.V. Ramani and M. Sajani *et al.*, 2012. Metagenomic analysis of Suri buffalo (*Bubalus bubalis*) rumen: A preliminary study. *Mol. Biol. Rep.*, 39: 4841-4848.

11. Hernández-García, J.A., R. Gonzalez-Escobedo, C.I. Briones-Roblero, C. Cano-Ramírez, F.N. Rivera-Orduña and G. Zúñiga, 2018. Gut bacterial communities of *Dendroctonus valens* and *D. Mexicanus* (Curculionidae: Scolytinae): A metagenomic analysis across different geographical locations in Mexico. *Int. J. Mol. Sci.*, Vol. 19. 10.3390/ijms19092578.
12. Wolfe, N.D., C.P. Dunavan and J. Diamond, 2007. Origins of major human infectious diseases. *Nature*, 447: 279-283.
13. Trinh, P., J.R. Zaneveld, S. Safranek and P.M. Rabinowitz, 2018. One health relationships between human, animal and environmental microbiomes: A mini-review. *Front. Public Health*, Vol. 6. 10.3389/fpubh.2018.00235.
14. Damborg, P., E.M. Broens, B.B. Chomel, S. Guenther and F. Pasmans *et al.*, 2016. Bacterial zoonoses transmitted by household pets: State-of-the-art and future perspectives for targeted research and policy actions. *J. Comp. Pathol.*, 155: S27-S40.
15. Ye, Z.W., S. Yuan, K.S. Yuen, S.Y. Fung, C.P. Chan and D.Y. Jin, 2020. Zoonotic origins of human coronaviruses. *Int. J. Biol. Sci.*, 16: 1686-1697.
16. Rombot, D.V., J. Pelealu, M. Tulung, J. Memah and M.Y. Samuel, 2019. The position of the species of cat lice (*Ctenocephalides felis*) based on molecular barcoding of the sub unit 1 (CO1) cytochrome oxidase gene. *Int. J. Entomol. Res.*, 4: 29-35.
17. Koo, H., J.A. Hakim, C.D. Morrow, M.R. Crowley, D.T. Andersen and A.K. Bej, 2018. Metagenomic analysis of microbial community compositions and cold-responsive stress genes in selected antarctic lacustrine and soil ecosystems. *Life*, Vol. 8. 10.3390/life8030029.
18. De Clerck, C., A. Fujiwara, P. Joncour, S. Léonard and M.L. Félix *et al.*, 2015. A metagenomic approach from aphid's hemolymph sheds light on the potential roles of co-existing endosymbionts. *Microbiome*, Vol. 3. 10.1186/s40168-015-0130-5.
19. Suddin, S., Y.S. Mokosuli, W. Marcelina, N. Orbanus and K. Ardi, 2019. Molecular barcoding based 16s rRNA gene of thermophilic bacteria from volcanic sites, Linow Lake, Tomohon. *Mater. Sci. Forum*, 967: 83-92.
20. Zhang, J. and J.M. Stewart, 2000. Economical and rapid method for extracting cotton genomic DNA. *J. Cotton Sci.*, 4: 193-201.
21. Datta, S., M.S. Samuel and E. Selvarajan, 2021. Exploring the bacterial community composition of soil from a tropical dry evergreen forest in Tamil Nadu, India. *Res. Square*, 10.21203/rs.3.rs-218538/v1.
22. Lee, P.Y., J. Costumbrado, C.Y. Hsu and Y.H. Kim, 2012. Agarose gel electrophoresis for the separation of DNA fragments. *J. Visual. Exp.*, Vol. 62. 10.3791/3923.
23. Weisburg, W.G., S.M. Barns, D.A. Pelletier and D.J. Lane, 1991. 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.*, 173: 697-703.
24. Chen, B., T. Yu, S. Xie, K. Du and X. Liang *et al.*, 2018. Comparative shotgun metagenomic data of the silkworm *Bombyx mori* gut microbiome. *Sci. Data*, Vol. 5. 10.1038/sdata.2018.285.
25. Rombot, D., J. Pelealu and M.Y. Samuel, 2021. The diversity and composition of new pathogenic bacteria in cat fleas. *Int. J. Pharm. Res.*, Vol. 13. 10.31838/ijpr/2021.13.02.347.
26. Rombot, D. and M.Y. Samuel, 2021. Biochemical characteristics and antibiotic resistance of bacterial isolate from *Ctenocephalides felis*. *J. Phys.: Conf. Ser.*, Vol. 1968. 10.1088/1742-6596/1968/1/012006.
27. Rolain, J.M., M. Franc, B. Davoust and D. Raoult, 2003. Molecular detection of *Bartonella quintana*, *B. koehlerae*, *B. henselae*, *B. clarridgeiae*, *Rickettsia felis* and *Wolbachia pipientis* in cat fleas, France. *Emerg. Infect. Dis.*, 9: 338-342.
28. Pornwiroon, W., M.T. Kearney, C. Husseneder, L.D. Foil and K.R. Macaluso, 2007. Comparative microbiota of *Rickettsia felis*-uninfected and -infected colonized cat fleas, *Ctenocephalides felis*. *ISME J.*, 1: 394-402.
29. Ferreira, F.C., D.M. Fonseca, G. Hamilton and D. Price, 2020. Metagenomic analysis of human-biting cat fleas in urban Northeastern United States of America reveals an emerging zoonotic pathogen. *Sci. Rep.*, Vol. 10. 10.1038/s41598-020-72956-x.
30. Douglas, G., A. Tsakris, S. Beleri, E. Patsoula, M. Linou, C. Billinis and J. Papaparaskevas, 2021. Molecular evidence of a broad range of pathogenic bacteria in *Ctenocephalides* spp.: Should we re-examine the role of fleas in the transmission of pathogens? *Trop. Med. Infect. Dis.*, Vol. 6. 10.3390/tropicalmed6010037.
31. Billeter, S.A., P.P.V. de Paiva Diniz, L.A. Jett, A.L. Wournell and A.M. Kjemtrup *et al.*, 2016. Detection of *Rickettsia* species in fleas collected from cats in regions endemic and nonendemic for flea-borne rickettsioses in California. *Vector-Borne Zoonotic Dis.*, 16: 151-156.
32. Roucher, C., O. Mediannikov, G. Diatta, J.F. Trape and D. Raoult, 2012. A new *Rickettsia* species found in fleas collected from human dwellings and from domestic cats and dogs in senegal. *Vector-Borne Zoonotic Dis.*, 12: 360-365.