

<http://www.pjbs.org>

PJBS

ISSN 1028-8880

Pakistan Journal of Biological Sciences

ANSI*net*

Asian Network for Scientific Information
308 Lasani Town, Sargodha Road, Faisalabad - Pakistan



Research Article

SWEET Gene Family in Sugar Beet (*Beta vulgaris*): Genome-Wide Survey, Phylogeny and Expression Analysis

¹Hong Viet La, ²Ha Duc Chu, ²Quyen Thi Ha, ³Thi Thanh Huyen Tran, ⁴Hai Van Tong, ⁵Tien Van Tran, ⁶Quynh Thi Ngoc Le, ⁴Huong Thi Thu Bui and ⁷Phi Bang Cao

¹Institute of Research and Application, Hanoi Pedagogical University 2, Phuc Yen City, Vinh Phuc 280000, Vietnam

²Faculty of Agricultural Technology, University of Engineering and Technology, Vietnam National University Hanoi, Xuan Thuy Road, Cau Giay District, Hanoi 122300, Vietnam

³Faculty of Biology, Hanoi National University of Education, Xuan Thuy Road, Cau Giay District, Hanoi 122300, Vietnam

⁴Faculty of Biotechnology, Vietnam National University of Agriculture, Gia Lam District, Hanoi 122300, Vietnam

⁵National Academy of Public Administration, Nguyen Chi Thanh Street, Dong Da District, Hanoi 122300, Vietnam

⁶Department of Biotechnology, Thuyloi University, Tay Son Street, Dong Da District, Hanoi 122300, Vietnam

⁷Faculty of Natural Sciences, Hung Vuong University, Nguyen Tat Thanh Street, Nong Trang Wards, Viet Tri City, Phu Tho Province 35000, Vietnam

Abstract

Background and Objective: The *SWEET* (Sugars Will Eventually be Exported Transporter) proteins play important roles in modulating the growth and development processes in plants. However, little information is available on the *SWEET* family in sugar beet (*Beta vulgaris*). The objectives of this present study were to genome-wide identify and characterize the *BvSWEET* family in sugar beet. **Materials and Methods:** Based on the available genome, proteome and transcriptome databases of sugar beet, various computational tools have been used to analyze the nucleotide and full-length protein sequences of members of the *BvSWEET* family. **Results:** A total of 16 members of the *BvSWEET* family has been identified in sugar beet at the genome-wide scale. Structural analysis indicated that the *BvSWEET* family exhibited variable characteristics. Furthermore, the *BvSWEET* family in sugar beet could be categorized into four distinct groups like in other plant species. Of our interest, we found that some *BvSWEET* genes exhibited strongly preferential expression in major organs/tissues under adverse environmental stimuli. **Conclusion:** The results provided a comprehensive foundation for further functional characterization of the *BvSWEET* gene family.

Key words: Identification, characterization, expression patterns, sugar beet, *SWEET*

Citation: La, H.V., H.D. Chu, Q.T. Ha, T.T.H. Tran and H.V. Tong *et al.*, 2022. *SWEET* gene family in sugar beet (*Beta vulgaris*): Genome-wide survey, phylogeny and expression analysis. Pak. J. Biol. Sci., 25: 387-395.

Corresponding Author: Huong Thi Thu Bui, Faculty of Biotechnology, Vietnam National University of Agriculture, Gia Lam District, Hanoi 122300, Vietnam
Phi Bang Cao, Faculty of Natural Sciences, Hung Vuong University, Nguyen Tat Thanh Street, Nong Trang Wards, Viet Tri City, Phu Tho Province 35000, Vietnam

Copyright: © 2022 Hong Viet La *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Sugar beet or red beetroot (*Beta vulgaris*) has been known as one of the most essential dicotyledonous plants, which is widely grown in the world. Supplying approximately 30% of the global sucrose, sugar beet was recognized as the second-largest source of refined table sugar¹. This economically important edible beet is also daily used as a source of a variety of natural pigments, like bioactive molecules (such as betalains, pectic-oligosaccharides and phenolics) and phytochemicals (such as flavonoid, carotenoid and vitamin)², which was reportedly related to numerous health benefits³. Thus, the efficiency of cultivation of this sugar-yielding crop has been comprehensively investigated.

To investigate and evaluate the molecular changes in sugar beets under adverse environmental conditions, many studies have been reported the roles of transporter protein families in the facilitation of sucrose accumulation⁴. Among them, one key sucrose transporter family has been known as 'SWEET-s' (Sugars Will Eventually be Exported Transporters). Participating in sucrose translocation and storage⁵, the SWEET family has been reportedly modulated many biological processes during the growth and development of plants⁶. Recently, the SWEET families have been comprehensively identified and characterized in various crop species, like rice⁷, tomato⁸, soybean⁹, oilseed rape¹⁰, cucumber¹¹, banana¹², Chinese white pear¹³, cotton¹⁴, tea plant¹⁵, apple¹⁶, litchi¹⁷, wheat¹⁸, *Medicago truncatula*¹⁹ and walnut²⁰. Unfortunately, no report has been made to a genome-wide survey of this critical gene family in sugar beet.

Thus, the purpose of this research was to comprehensively identify and characterize the SWEET family in sugar beet. Particularly, a genome-wide survey has been performed to seek all putative members of the SWEET family. We also analyzed their general characteristics, like physic-chemical properties, subcellular localization, gene structure, phylogenetic tree and transmembrane helices. Furthermore, the expression patterns of the *SWEET* gene in major tissues/organs of sugar beet were determined by re-analyzing the previous transcriptome atlas.

MATERIALS AND METHODS

Study area: This study was carried out from March, 2021 to January, 2022. All analyses were conducted at Hanoi City and Vinh Phuc and Phu Tho provinces, Vietnam.

Genome-wide survey of the SWEET family in sugar beet:

To comprehensively seek SWEET-s in sugar beet, the well-characterized domain of plant's SWEET⁵ obtained from the Pfam tool²¹ was applied to search against the latest assembly of sugar beet (Accession number: GCF_000511025.2)²² in the NCBI (National Center for Biotechnology Information) and Phytozome 12.0²³ portals. The major annotated features of the identified SWEET-s, including peptide sequence, coding DNA sequence (CDS), genomic DNA sequence (gDNA), identifier code and chromosomal distribution were consequently obtained for further analyses.

Analysis of protein properties of the SWEET family in sugar beet:

The physic-chemical characteristics of the SWEET proteins, like length, molecular weight, iso-electric point, aliphatic index, hydropathicity, were theoretically calculated by the ExPASy ProtParam tool²⁴ as previously guided^{25,26}.

Prediction of the subcellular localization of the BvSWEET-s:

The subcellular localization of the SWEET proteins were predicted by using the web-based YLOC tool²⁷. The full-length protein sequence of each member of the SWEET family in sugar beet was used as a query for a search in the YLOC system²⁷. Particularly, a total of 10 locations, including the nucleus, cytoplasm, mitochondrion, plasma membrane, extracellular space, endoplasmic reticulum, peroxisome, Golgi apparatus, vacuole and chloroplast were predicted for the plant model²⁷.

Analysis of the membrane-bound SWEET proteins in sugar

beet: To predict the transmembrane helices in SWEET proteins in sugar beet, the TMHMM 2.0 tool²⁸ was used. The full-length protein sequences of the SWEET family in sugar beet were used as seed sequences for seeking the transmembrane helices²⁸.

Construction of phylogenetic tree of the SWEET family in sugar beet:

To construct the phylogenetic tree of the SWEET family, the full-length protein sequences of all members of the SWEET family in sugar beet were collected as raw material for the MEGA (Molecular Evolutionary Genetics Analysis) 7.0 software²⁹. The Maximum Likelihood estimation with 1,000 bootstrapped replications was used as the statistical model for generating the phylogenetic tree. Additionally, also collected the full-length protein sequences of the SWEET families from *Arabidopsis thaliana*³⁰, rice⁷ and cassava in the Phytozome²³ to analyze the categorization of the SWEET families.

Analysis of gene structures of the *SWEET* family in sugar beet:

The number of exons and introns of genes encoding the *SWEET* family in sugar beet was analyzed by using the GSDS (Gene Structure Display Server) 2.0 website³¹. The order of members of the *SWEET* gene family was arranged based on the Maximum-Likelihood-based phylogenetic tree²⁹. The result was then illustrated by using Adobe Illustrator^{25,26}.

Analysis of the expression patterns of the *SWEET* family in sugar beet:

To investigate the expression patterns of the *SWEET* gene family in sugar beet, we re-analyzed the previous transcriptome databases by searching against the NCBI GEO (NCBI Gene expression omnibus)³², like GSE135555³³, GSE138039³⁴, GSE107627³⁵ and GSE114968³⁶. Particularly, the first microarray database (GSE135555) provided expression data of fifteen-day-old roots of the sugar beet plants treated with a beet cyst nematode, namely *Heterodera schachtii*³³, while the second dataset (GSE138039) contained expression data of leaves on five-week-old sugar beet plants inoculated with a fungal pathogen, namely *Sclerotinia sclerotiorum*³⁴. Additionally, we also downloaded two datasets, GSE107627³⁵ and GSE114968³⁶, which provided two expression data of leaf tissues from alkaline-treated seedlings and root tissues from salinity-treated seedlings, respectively. Differentially expressed gene was assigned by a fold-change cut-off ($|\text{fold-change}| > 2.0$). The cluster heatmap of the *SWEET* genes in sugar beet was then visualized in R software with the g plots³⁷.

RESULTS AND DISCUSSION

Identification, annotation and chromosome localization of the *SWEET* family in sugar beet: First of all, we seek the *SWEET* family in sugar beet by Blast-ing a conserved domain

of *SWEET* molecules, namely 'PF03083'¹⁵ against the available sugar beet's assembly²² from Phytozome²³ and NCBI databases. As a result, a total of 16 putative members of the *SWEET* family has been found and validated in sugar beet in Table 1.

As compared with the *SWEET* families in other plant species, we found that the amount of members from this important family is critically variable³⁸. For example, 21 putative members have been identified and characterized in rice⁷. In the cucumber and tea plant, 17 and 13 members of the *SWEET* family have been reported recently, respectively^{11,15}, while 25 *SWEET*-s were found in Medicago's assembly¹⁹. In four cotton species, 22, 31, 55 and 60 members of the *SWEET* family have been listed in *Gossypium arboreum*, *G. raimondii*, *G. hirsutum* and *G. barbadense*, respectively¹⁴. The number of members of the *SWEET* family could also reach 52 and 108 in soybean and wheat, respectively^{9,18}. Quite recently, a total of 25 members of the *SWEET* family has been studied in walnut²⁰. Here, our genome-wide survey indicated that the number of *SWEET* family in sugar beet (16) is more than in tea plant (13)¹⁵ but less than many plant species, like cucumber (17)¹¹, rice (21)⁷, *G. arboreum* (22)¹⁴, Medicago (25)¹⁹, walnut (25)²⁰, *G. raimondii* (31)¹⁴, soybean (52)⁹, two other cotton species (55 and 60 in *G. hirsutum* and *G. barbadense*, respectively)¹⁴ and wheat (108)¹⁸.

Next, the chromosomal locations of all 16 members of the *SWEET* gene family were manually retrieved from the NCBI database. According to the chromosomal localization of *SWEET* genes in Fig. 1, we named 16 annotated genes, from *BvSWEET01* to 16 by their order of occurrence on the chromosomes. It was noted that 16 *BvSWEET* genes were located on the sugar beet genome with an even ratio. Particularly, the chromosomes 2, 6 and 7 harboured only one *BvSWEET* gene each, while two chromosomes, like 3 and 5

Table 1: BvSWEET family in sugar beet

Gene name	Locus name	Length	Mass	pI	AI	GRAVY	SL	TMH
<i>BvSWEET01</i>	EL10Ac1g00932	234	26.39	9.27	115.30	0.70	PM	7
<i>BvSWEET02</i>	EL10Ac1g01460	255	27.95	9.50	116.59	0.59	PM	7
<i>BvSWEET03</i>	EL10Ac2g02891	162	18.71	8.71	121.42	0.81	PM	4
<i>BvSWEET04</i>	EL10Ac3g05343	251	27.86	8.89	113.03	0.49	PM	7
<i>BvSWEET05</i>	EL10Ac3g06628	246	27.22	8.35	121.99	0.68	PM	6
<i>BvSWEET06</i>	EL10Ac3g06631	241	26.47	7.70	109.13	0.53	PM	7
<i>BvSWEET07</i>	-	162	18.61	8.47	122.04	0.79	PM	5
<i>BvSWEET08</i>	EL10Ac4g08428	234	26.03	6.25	112.48	0.81	PM	7
<i>BvSWEET09</i>	EL10Ac4g09802	262	29.42	8.93	117.14	0.59	PM	7
<i>BvSWEET10</i>	EL10Ac4g09898	223	24.49	8.93	130.67	0.89	PM	7
<i>BvSWEET11</i>	EL10Ac5g10774	234	26.74	9.22	122.35	0.72	PM	6
<i>BvSWEET12</i>	EL10Ac5g11351	324	36.27	6.52	105.49	0.34	PM	7
<i>BvSWEET13</i>	EL10Ac5g12959	257	28.22	9.59	123.62	0.77	PM	7
<i>BvSWEET14</i>	EL10Ac5g12960	273	30.36	8.99	114.62	0.58	PM	7
<i>BvSWEET15</i>	EL10Ac6g13562	258	28.86	9.25	123.49	0.65	PM	7
<i>BvSWEET16</i>	EL10Ac7g17402	335	37.62	8.19	108.81	0.30	PM	7

pI: Iso-electric point, AI: Aliphatic index, GRAVY: Grand average of hydropathicity, SL: Sub-cellular localization, TMH: Transmembrane helices and PM: Plasma membrane




Fig. 1: Chromosomal distribution of the *SWEET* gene family in sugar beet genome

contained the highest number of *BvSWEET* genes (four each) (Fig. 1). We also found that two and three *BvSWEET* genes were mapped on chromosomes 1 and 4, respectively (Fig. 1). However, no *BvSWEET* gene was found on chromosomes 8 and 9.

Interestingly, two *BvSWEET* genes, like *BvSWEET13* and *14* were mapped on the sub-telomeric areas of chromosome 5. This trend was also reported in the *SWEET* gene families in other plant species, like rice⁷, soybean⁹, cucumber¹¹, tea plant¹⁵, four cotton species¹⁴, Medicago¹⁹ and wheat¹⁸. We hypothesized that genes on the sub-telomeric region of the chromosome, like two *BvSWEET* genes in this study, may act in the chromosome pairing and recognition during the meiosis process in sugar beet, as previously reported in other plant species³⁹.

Investigation on the physico-chemical properties of the *SWEET* family in sugar beet: To get insight into the *BvSWEET* family in sugar beet, the physico-chemical characteristics of each *BvSWEET*'s member were analyzed by the online ExPASy ProtParam tool²⁴. Particularly, five parameters, including protein length, mass, iso-electric point, aliphatic index and hydropathicity were explored by applying the full-length protein sequences to tool²⁴. All features of the *BvSWEET* family were then provided in Table 1.

The results revealed that the lengths of gDNA *SWEET*-s varied from 489 (*BvSWEET03* and *07*) to 15046 bp (*BvSWEET15*), whereas, the protein lengths ranged from 162 (*BvSWEET03* and *07*) to 335 amino acid residues (*BvSWEET16*) (Table 1). Moreover, the molecular weights of the *BvSWEET* family were changed from 18.61 (*BvSWEET07*) to 37.62 kDa (*BvSWEET16*) (Table 1). A majority of the *BvSWEET* proteins (15 out of 16) had alkaline properties (the pI values were more than 7.0), whereas, only *BvSWEET08* had acidic property (the pI value was less than 7.0) (Table 1). Interestingly, *in silico* analysis indicated that all members of the *BvSWEET* family are hydrophilic with positive GRAVY values (Table 1), while all these molecules contained a large amount of aliphatic amino

acid residues, with the aliphatic index ranging from 105.49 (*BvSWEET12*) to 130.67 (*BvSWEET10*) (Table 1). Findings suggested that the variable features of the *BvSWEET* family may be related to their potential roles during the growth and development of sugar beet plants¹².

Previously, great efforts have been recorded to investigate the physico-chemical properties of the *SWEET* families in other plant species. For example, the *SWEET* family from tomato, oilseed rape, banana, apple and litchi were varied from 233-308, 56-303, 171-333 (with relative molecular weight between 19.1-37.4 kDa), 215-340 and 229-300 (with relative molecular weight between 25.6-33.6 kDa) amino acid residues, respectively^{8,10,12,16,17}. Recently, the walnut *SWEET* proteins have been reported to vary from 154-301 amino acid residues in size, their molecular weights ranged from 16.63-33.24 kDa²⁰. Moreover, a majority of the members (22 out of 25) of the *SWEET* family from banana had basic properties (iso-electric points ranging from 7.56-9.83)¹², while all 16 members of the *SWEET* family in litchi also exhibited alkaline properties¹⁷. The pI values of 21 (out of 25) members of the *SWEET* family in walnut had also basic properties²⁰.

Sub-cellular localization and membrane-bound analysis of the *SWEET* family in sugar beet: It is thought that the subcellular localization of proteins may be strongly related to their specific function⁴⁰. Thus, the study carried out a prediction of the sub-cellular distribution of *BvSWEET* proteins in sugar beet by using the YLOC portal²⁷. As expected, we found that all 16 *BvSWEET* proteins may be localized on the plasma membrane (Table 1). Findings were also confirmed in the *SWEET* proteins in other plant species, like tomato⁸, walnut²⁰. For example, five members of the *SWEET* family in tomato were predicted to distribute on the plasma membrane⁸, while 22 (out of 25) *SWEET* proteins from walnut were also localized on the plasma membrane²⁰. Moreover, the prediction of subcellular localization in a previous study showed that all 168 *SWEET* proteins from four cotton species were located in the plasma membrane¹⁴.




Fig. 2: Phylogenetic tree diagram of SWEET families from sugar beet, *Arabidopsis thaliana*, rice and cassava

Next, the presence of transmembrane helices was then investigated on the full-length protein sequences of the BvSWEET family in sugar beet. The predicted result was provided in Table 1. We found that a majority of members (12 out of 16) of the BvSWEET family contained seven transmembrane helices (Table 1). Two members, including BvSWEET05 and 11, were noted to contain six transmembrane helices, while BvSWEET07 and 03 harboured five and four transmembrane helices, respectively (Table 1). Previously, SWEET families in litchi¹⁷ and some other plant species⁹ were recognized to contain seven transmembrane helices⁵. Taken together, suggested that BvSWEET proteins, with the presence of these transmembrane helices, may function as membrane receptors⁴¹.

Phylogenetic analysis of the SWEET family in sugar beet:

To categorize the BvSWEET family in sugar beet, a maximum-likelihood-based phylogenetic tree of full-length SWEET proteins from sugar beet, *A. thaliana*³⁰, rice⁷ and cassava has been successfully constructed in Fig. 2. As provided in Fig. 2, all members of the BvSWEET family from sugar beet could be classified into four distinct groups. Six members, including BvSWEET01, 03, 07, 11, 13 and 14 have been fall into group I, while three (BvSWEET02, 04 and 08) and three (BvSWEET05, 06 and 10) members were categorized into group II and III, respectively (Fig. 2). Finally, group IV contained four remaining members, like BvSWEET09, 12, 15, 16 (Fig. 2). Here, we realized that proteins in the same branch shared similar characteristics (Fig. 2, Table 1). For example,




Fig. 3: Gene organizations of the *BvSWEET* gene family in sugar beet

BvSWEET03 and *07* were localized in the same clade of group I (Fig. 2), they shared similar protein sizes, molecular weights, iso-electric points, aliphatic index and average grand of hydrophathy). Thus, we suggested that these same-clade-SWEET proteins may share similar functions.

In the previous study, a phylogenetic analysis of the SWEET families in tomato and two other plant species, like rice and *A. thaliana* has been performed⁸. As expected, the SWEET family in three targeted plant species could be categorized into four classes⁸. Furthermore, a comprehensive phylogenetic tree of 173 SWEET proteins from 13 plant species, including four dicots, two monocots, two bryophytes and five algae has been well-described⁹. As expected again, four distinct clades were perceived⁹. Taken together, findings revealed that the SWEET family in sugar beet, perhaps in plant species, could be categorized into four major groups.

Gene structure of the *SWEET* family in sugar beet: To investigate the structure of the gene encoding *BvSWEET* proteins, we analyzed the exon/intron organization of each member of the *BvSWEET* gene family by using the GSDS tool³¹. The gene organizations of all *BvSWEET* genes were then generated and provided in Fig. 3. We found that most members (14 out of 16) of the *BvSWEET* gene family contained introns in their genomic sequences, whereas, two genes, including *BvSWEET03* and *07* were noted to be intronless. Next, three genes, like *BvSWEET13*, *14* and *16* contained five exons, while the remaining *BvSWEET* gene family members (11 out of 16) harboured six exons (Fig. 3).

Significantly, such a similar phenomenon of six exons in the *BvSWEET* gene family was also confirmed in the *SWEET* families from many crop species. In tomatoes, some *SWEET* genes (23 out of 29) contained six exons and five introns⁸. Also, 51 (out of 68), 12 (out of 17), 12 (out of 18) members of

the *SWEET* families from oilseed rape, cucumber, Chinese white pear, respectively^{10,11,13} contained six exons. Recently, many members of the *SWEET* gene families in the tea (11 out of 13) and litchi (14 out of 16) have been reported to harbour six exons^{15,17}. Taken together, our comparisons strongly indicated that the majority of the *SWEET* gene families in plant species might have the conserved motif of six exons and five introns.

Expression patterns of the *SWEET* family during the growth and development of sugar beet: As the main part of this study, the expression patterns of the *BvSWEET* gene family members were explored by re-analyzing four available RNA-Seq datasets³³⁻³⁶ in the NCBI GEO³². A heatmap was then generated and provided in Fig. 4. Noted that two genes, including *BvSWEET03* and *07* not expressed in all datasets. The expression of all remaining members of the *BvSWEET* gene family displayed spatial variation in major organs/tissues of sugar beet plants.

Under the alkaline stress³⁵, only three genes, like *BvSWEET04*, *10* and *11* exhibited a significantly different expression in root tissues (Fig. 4). Among them, *BvSWEET10* and *11* were noted to strongly induce and reduce in alkaline-treated roots, respectively (Fig. 4). Under the salt stress³⁶, eight genes, like *BvSWEET01*, *02*, *06*, *09*, *12*, *14*, *15* and *16* were differentially expressed in leaves from alkaline-treated seedlings. Interestingly, one (*BvSWEET12*) and two (*BvSWEET14* and *15*) genes were highly up-regulated and down-regulated in leaf tissues under the alkaline treatment, respectively. Next, under the infection of a fungal pathogen³⁴, six (*BvSWEET04*, *05*, *06*, *09*, *12* and *14*) and one (*BvSWEET10*) genes were highly reduced and induced in leaf tissues, respectively (Fig. 4). Finally, under the inoculation of *H. schachtii*³³, only two genes, *BvSWEET10* and *14* were




Fig. 4: Expression patterns of the *BvSWEET* gene family during the growth and development of sugar beet plants

noted to be up-regulated and down-regulated in root tissues (Fig. 4). Previously, various efforts have been recorded to investigate the role of *SWEET* genes in response to abiotic stress(es) in higher plants, such as banana¹², cotton¹⁴, *P. equestris* and *D. officinale*⁴² and *M. truncatula*¹⁹. Taken together, our re-analysis suggested that the *BvSWEET* gene family is differentially expressed in various organs/tissues under numerous abiotic/biotic stress conditions, which may suggest their potential functions during growth and development processes and perhaps in responses to adverse environmental conditions.

CONCLUSION

In this present study, a comprehensive analysis of 16 members of the *BvSWEET* family has been investigated in sugar beet. Various information of the *BvSWEET* family, like chromosomal locations, physic-chemical properties, subcellular localizations, exon/intron organizations, phylogenetic trees, gene structures has been analyzed by using computational tools. We re-analyzed four transcriptome datasets to explore the expression profiles of the *BvSWEET* gene family in major organs/tissues during the growth and development processes. To sum up, the data

provide valuable information for further functional characterization of sugar beet *BvSWEET* genes.

SIGNIFICANCE STATEMENT

The features and expression analyses of sugar beet *SWEET* genes were originally presented in this paper. The findings laid the groundwork for future functional studies, especially those involving the *BvSWEET* gene in these important crops.

REFERENCES

1. Zhang, Y., J. Nan and B. Yu, 2016. OMICS technologies and applications in sugar beet. *Front. Plant Sci.*, Vol. 7. 10.3389/fpls.2016.00900.
2. Chhikara, N., K. Kushwaha, P. Sharma, Y. Gat and A. Panghal, 2018. Bioactive compounds of beetroot and utilization in food processing industry: A critical review. *Food Chem.*, 272: 192-200.
3. de Oliveira, S.P.A., H.M.A. do Nascimento, K.B. Sampaio and E.L. de Souza, 2021. A review on bioactive compounds of beet (*Beta vulgaris* L. subsp. *vulgaris*) with special emphasis on their beneficial effects on gut microbiota and gastrointestinal health. *Crit. Rev. Food Sci. Nutr.*, 61: 2022-2033.

4. Jung, B., F. Ludewig, A. Schulz, G. Meißner and N. Wöstefeld *et al.*, 2015. Identification of the transporter responsible for sucrose accumulation in sugar beet taproots. *Nat. Plants*, Vol. 1. 10.1038/nplants.2014.1.
5. Julius, B.T., K.A. Leach, T.M. Tran, R.A. Mertz and D.M. Braun, 2017. Sugar transporters in plants: New insights and discoveries. *Plant Cell Physiol.*, 58: 1442-1460.
6. Chen, L.Q., 2014. SWEET sugar transporters for phloem transport and pathogen nutrition. *New Phytol.*, 201: 1150-1155.
7. Yuan, M. and S. Wang, 2013. Rice MtN3/Saliva/SWEET family genes and their homologs in cellular organisms. *Mol. Plant*, 6: 665-674.
8. Feng, C.Y., J.X. Han, X.X. Han and J. Jiang, 2015. Genome-wide identification, phylogeny, and expression analysis of the SWEET gene family in tomato. *Gene*, 573: 261-272.
9. Patil, G., B. Valliyodan, R. Deshmukh, S. Prince and B. Nicander *et al.*, 2015. Soybean (*Glycine max*) SWEET gene family: Insights through comparative genomics, transcriptome profiling and whole genome re-sequence analysis. *BMC Genomics*, Vol. 16. 10.1186/s12864-015-1730-y.
10. Jian, H., K. Lu, B. Yang, T. Wang and L. Zhang *et al.*, 2016. Genome-wide analysis and expression profiling of the *SUC* and *SWEET* gene families of sucrose transporters in oilseed rape (*Brassica napus* L.) *Front. Plant Sci.*, Vol. 7. 10.3389/fpls.2016.01464.
11. Hu, L.P., F. Zhang, S.H. Song, X.W. Tang and H. Xu *et al.*, 2017. Genome-wide identification, characterization, and expression analysis of the SWEET gene family in cucumber. *J. Integr. Agric.*, 16: 1486-1501.
12. Miao, H., P. Sun, Q. Liu, Y. Miao and J. Liu *et al.*, 2017. Genome-wide analyses of SWEET family proteins reveal involvement in fruit development and abiotic/biotic stress responses in banana. *Sci Rep.*, Vol. 7. 10.1038/s41598-017-03872-w.
13. Li, J., M. Qin, X. Qiao, Y. Cheng, X. Li, H. Zhang and J. Wu 2017. A new insight into the evolution and functional divergence of SWEET transporters in Chinese white pear (*Pyrus bretschneideri*). *Plant Cell Physiol.*, 58: 839-850.
14. Zhao, L., J. Yao, W. Chen, Y. Li and Y. Lu *et al.*, 2018. A genome-wide analysis of *SWEET* gene family in cotton and their expressions under different stresses. *J. Cotton Res.*, Vol. 1. 10.1186/s42397-018-0007-9.
15. Wang, L., L. Yao, X. Hao, N. Li and W. Qian *et al.*, 2018. Tea plant SWEET transporters: Expression profiling, sugar transport, and the involvement of CsSWEET16 in modifying cold tolerance in *Arabidopsis*. *Plant Mol. Biol.*, 96: 577-592.
16. Zhen, Q., T. Fang, Q. Peng, L. Liao, L. Zhao, A. Owiti and Y. Han, 2018. Developing gene-tagged molecular markers for evaluation of genetic association of apple SWEET genes with fruit sugar accumulation. *Hortic. Res.*, Vol. 5. 10.1038/s41438-018-0024-3.
17. Xie, H., D. Wang, Y. Qin, A. Ma and J. Fu *et al.*, 2019. Genome-wide identification and expression analysis of SWEET gene family in *Litchi chinensis* reveal the involvement of *LcSWEET2a/3b* in early seed development. *BMC Plant Biol.*, Vol. 19. 10.1186/s12870-019-2120-4.
18. Gautam, T., G. Saripalli, V. Gahlaut, A. Kumar, P.K. Sharma, H.S. Balyan and P.K. Gupta, 2019. Further studies on sugar transporter (SWEET) genes in wheat (*Triticum aestivum* L.). *Mol. Biol. Rep.*, 46: 2327-2353.
19. Hu, B., H. Wu, W. Huang, J. Song, Y. Zhou and Y. Lin, 2019. SWEET gene family in *Medicago truncatula*: Genome-wide identification, expression and substrate specificity analysis. *Plants*, Vol. 8. 10.3390/plants8090338.
20. Jiang, S., B. Balan, R. de A.B. Assis, C.H.D. Sagawa and X. Wan *et al.*, 2020. Genome-wide profiling and phylogenetic analysis of the SWEET sugar transporter gene family in walnut and their lack of responsiveness to *Xanthomonas arboricola* pv. *juglandis* infection. *Int. J. Mol. Sci.*, Vol. 21. 10.3390/ijms21041251.
21. El-Gebali, S., J. Mistry, A. Bateman, S.R. Eddy and A. Luciani *et al.*, 2019. The Pfam protein families database in 2019. *Nucleic Acids Res.*, 47: D427-D432.
22. Dohm, J.C., A.E. Minoche, D. Holtgräwe, S. Capella-Gutiérrez and F. Zakrzewski *et al.*, 2014. The genome of the recently domesticated crop plant sugar beet (*Beta vulgaris*). *Nature*, 505: 546-549.
23. Goodstein, D.M., S. Shu, R. Howson, R. Neupane and R.D. Hayes *et al.*, 2012. Phytozome: A comparative platform for green plant genomics. *Nucleic Acids Res.*, 40: D1178-D1186.
24. Gasteiger, E., A. Gattiker, C. Hoogland, I. Ivanyi, R.D. Appel and A. Bairoch, 2003. ExPASy: The proteomics server for in-depth protein knowledge and analysis. *Nucleic Acids Res.*, 31: 3784-3788.
25. Chu, H., K. Nguyen, Y. Watanabe, D. Le, T. Pham, K. Mochida and L.S. Tran, 2018. Identification, structural characterization and gene expression analysis of members of the nuclear factor-Y family in chickpea (*Cicer arietinum* L.) under dehydration and abscisic acid treatments. *Int. J. Mol. Sci.*, Vol. 19. 10.3390/ijms19113290.
26. Niu, L. H.D. Chu, C.D. Tran, K.H. Nguyen and H.X. Pham *et al.*, 2020. The *GATA* gene family in chickpea: Structure analysis and transcriptional responses to abscisic acid and dehydration treatments revealed potential genes involved in drought adaptation. *J. Plant Growth Regul.*, 39: 1647-1660.
27. Briesemeister, S., J. Rahnenfuhrer and O. Kohlbacher, 2010. YLoc-an interpretable web server for predicting subcellular localization. *Nucleic Acids Res.*, 38: W497-W502.
28. Chaturvedi, N., S. Shanker, V.K. Singh, D. Sinha and P.N. Pandey, 2011. Hidden markov model for the prediction of transmembrane proteins using MATLAB. *Bioinformation*, 7: 418-421.

29. Kumar, S., G. Stecher and K. Tamura, 2016. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. *Mol. Biol. Evol.*, 33: 1870-1874.
30. Durand, M., D. Mainson, B. Porcheron, L. Maurousset, R. Lemoine and N. Pourtau, 2018. Carbon source-sink relationship in *Arabidopsis thaliana*: The role of sucrose transporters. *Planta*, 247: 587-611.
31. Hu, B., J. Jin, A.Y. Guo, H. Zhang, J. Luo and G. Gao, 2015. GSDS 2.0: An upgraded gene feature visualization server. *Bioinformatics*, 31: 1296-1297.
32. Barrett, T., S.E. Wilhite, P. Ledoux, C. Evangelista and I.F. Kim *et al.*, 2013. NCBI GEO: Archive for functional genomics data sets-update. *Nucleic Acids Res.*, 41: D991-D995.
33. Ghaemi, R., E. Pourjam, N. Safaie, B. Verstraeten and S.B. Mahmoudi *et al.*, 2020. Molecular insights into the compatible and incompatible interactions between sugar beet and the beet cyst nematode. *BMC Plant Biol.*, Vol. 20. 10.1186/s12870-020-02706-8.
34. Sucher, J., M. Mbengue, A. Dresen, M. Barascud, M. Didelon, A. Barbacci and S. Raffaele, 2020. Phylotranscriptomics of the pentapetalae reveals frequent regulatory variation in plant local responses to the fungal pathogen *Sclerotinia sclerotiorum*. *Plant Cell*, 32: 1820-1844.
35. Zou, C., Y. Wang, B. Wang, D. Liu, L. Liu, Z. Gai and C. Li, 2020. Long non-coding RNAs in the alkaline stress response in sugar beet (*Beta vulgaris* L.). *BMC Plant Biol.*, Vol. 20. 10.1186/s12870-020-02437-w.
36. Liu, L., B. Wang, D. Liu, C. Zou and P. Wu *et al.*, 2020. Transcriptomic and metabolomic analyses reveal mechanisms of adaptation to salinity in which carbon and nitrogen metabolism is altered in sugar beet roots. *BMC Plant Biol.*, Vol. 20. 10.1186/s12870-020-02349-9.
37. Liao, Y., G.K. Smyth and W. Shi, 2019. The R package *Rsubread* is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.*, 47: e47-e47.
38. Jia, B., X.F. Zhu, Z.J. Pu, Y.X. Duan and L.J. Hao *et al.*, 2017. Integrative view of the diversity and evolution of SWEET and SemiSWEET sugar transporters. *Front. Plant Sci.*, Vol. 8. 10.3389/fpls.2017.02178.
39. del Carmen Calderón, M., M.D. Rey, A. Cabrera and P. Prieto, 2014. The subtelomeric region is important for chromosome recognition and pairing during meiosis. *Sci. Rep.*, Vol. 4. 10.1038/srep06488.
40. Goodin, M.M., 2018. Chapter Six-Protein Localization and Interaction Studies in Plants: Toward Defining Complete Proteomes by Visualization. In: *Advances in Virus Research*, Kielian, M., T.C. Mettenleiter and M.J. Roossinck (Eds.), Elsevier, Netherlands, pp: 117-114.
41. Xuan, Y.H., Y.B. Hu, L.Q. Chen, D. Sosso, D.C. Ducat, B.H. Hou and W.B. Frommer, 2013. Functional role of oligomerization for bacterial and plant SWEET sugar transporter family. *PNAS*, 110: E3685-E3694.
42. Wang, T., Z. Song, W.L. Meng and L.B. Li, 2018. Identification, characterization, and expression of the SWEET gene family in *Phalaenopsis equestris* and *Dendrobium officinale*. *Biol. Plant.*, 62: 24-32.