

PJN

ISSN 1680-5194

PAKISTAN JOURNAL OF
NUTRITION

ANSI*net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan
Mob: +92 300 3008585, Fax: +92 41 8815544
E-mail: editorpjn@gmail.com

Association of Serum C-Reactive Protein (CRP) with Some Nutritional Parameters of Maintenance Hemodialysis Patients

Azar Baradaran¹ and Hamid Nasri^{2*}

¹Department of Biochemistry, Center of Research and Reference Laboratory, Iran

²Hemodialysis Department, Hajar Medical, Educational and Therapeutic Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

Abstract: Malnutrition and inflammation are common in hemodialysis patients, and are usually closely associated. Serum C-reactive protein (CRP) concentrations have been found to be significantly elevated in hemodialysis patients and reflects chronic inflammation, and as an acute-phase reactant, is a sensitive and independent marker of malnutrition. To investigate the association of serum CRP level with some nutritional variables in diabetic and non diabetic end-stage renal failure patients undergoing regular hemodialysis, we designed a study of 36 maintenance hemodialysis patients (f = 15 m = 21), consisting of 25 non-diabetic HD patients and 11 diabetic HD patients. In this study we found a near significant difference of CRP between diabetic and non-diabetics of total patients with more values of CRP in diabetics, a significant difference of CRP between diabetic and non-diabetic of female HD patients with more values in diabetics and a significant difference of CRP between males and female of non-diabetic population with more values of CRP in males. An inverse correlation of serum CRP with serum cholesterol and triglyceride levels and a near significant positive correlations of CRP with serum ALP and with serum iPTH were found. An inverse correlation of serum CRP with dialysis efficacy was seen too. No significant association between serum CRP and serum albumin was seen. Compatible with some studies and in contrast to some other studies, the association of serum albumin with serum CRP levels in this study was insignificant. The positive correlation of high serum PTH with inflammation implies further need to control of hyperphosphatemia and secondary hyperparathyroidism in HD patients, also inverse correlation of serum CRP with cholesterol and triglyceride further support the malnutrition-inflammation complex syndrome (MICS) which frequently seen in HD patients.

Key words: End-stage renal failure, hemodialysis, C-reactive protein (CRP), malnutrition-inflammation

Introduction

C-reactive protein (CRP), is an acute phase protein whose synthesis in the liver is regulated by different cytokines, particularly interleukin 6 (IL-6). Plasma levels of CRP in the absence of active disease are low, but can rise up to 1000-fold in patients with an inflammatory reaction. Besides being a marker of inflammation, CRP itself may have proinflammatory properties since it can activate the complement system (Wolbink *et al.*, 1996; Morrow and Ridker, 2000). Thus elevated plasma concentrations of C-reactive protein (CRP), a sensitive marker of underlying systemic inflammation. (Clyne and Olshaker, 1999; McIntyre *et al.*, 1997; Haubitz *et al.*, 1996). Serum CRP concentrations have also been found to be significantly elevated in hemodialysis patients (Stenvinkel *et al.*, 2000; Panichi *et al.*, 2000) and reflects chronic inflammation, and as an acute-phase reactant, is a sensitive and independent marker of malnutrition (Ortega *et al.*, 2002). An elevated serum C-reactive protein has been shown to be strongly predictive of morbidity and mortality in dialysis patients, specially is a strong predictor of cardiovascular mortality in hemodialysis patients and (Kim *et al.*, 2005; Caglar *et*

al., 2002). Uremic malnutrition is characterized by insidious loss of somatic protein stores (reflected by lean body mass or serum creatinine) and visceral protein concentrations (reflected by serum albumin or serum prealbumin) (Fernandez-Reyes *et al.*, 2000). The etiology of malnutrition in renal failure is complex and may include many factors, e.g. poor food intake because of anorexia, nausea and vomiting due to uraemic toxicity, hormonal derangements, acidosis and increased resting energy expenditure (Fernandez-Reyes *et al.*, 2000; Fujino *et al.*, 2005). Malnutrition and inflammation are common in hemodialysis patients, and are usually closely associated (Marckmann, 1988). Indeed the prevalence of protein-energy malnutrition in hemodialysis (HD) patients is high (23 to 73%) in various studies (Enia *et al.*, 1993; Cianciaruso *et al.*, 1995; Leavey *et al.*, 1998; Lowrie and Lew, 1990). Several recent reports have focused on the association between nutritional status and clinical outcome, providing support for the hypothesis that malnutrition may cause or contribute to mortality and predict poor clinical outcome in end-stage renal disease patients. (Leavey *et al.*, 1998; Lowrie and Lew, 1990;

Baradaran and Nasri: Association of Serum C-Reactive Protein with Some Nutritional Parameters

Table 1: Mean±SD, Minimum and Maximum of age, duration and dosage hemodialysis and also laboratory results of total hemodialysis patients

Total patients n=36		Min.	Max.	Mean±SD
Age	years	16	80	46.6±17
DH*	months	2	156	32±36
Dialysis dose	sessions	36	1584	294±3393
URR	%	39	76	59±8.9
BMI	kg/m ²	16	34	22±4.4
iPTH	Pg/ml	16	1980	434±455
Ca	mg/dl	5	10	7.7±0.94
P	mg/dl	3	10	6.4±2
Alp	IU/L	175	5487	628±891
HCO3	mE/l	14	25	20±2.6
CRP	mg/l	3	40	8.7±6.7
BUN	mg/dl	30	180	83±33
Creat	mg/dl	3	18	9.5±3.6
Alb	g/dl	2.4	4.8	3.8±0.5
25-OH Vit D	nMol/l	1.3	105	10.5±18
Chol	mg/dl	59	211	117±39
TG	mg/dl	29	461	130±95
LDL	mg/dl	12	122	61.6±22
HDL	mg/dl	2	70	37±11

*Duration of hemodialysis treatment

Table 2: Mean±SD, Minimum and Maximum of age, duration and dosage hemodialysis and also laboratory results of non-diabetic hemodialysis patients

Total patients n=25		Min.	Max.	Mean±SD
Age	years	16	80	44±17
DH*	months	2	156	40±40.8
Dialysis dose	sessions	36	1584	370±452
URR	%	60	76	61±7.5
BMI	kg/m ²	16	36	21±4.6
iPTH	Pg/ml	22	1980	537±483
Ca	mg/dl	6	9	7.8±0.70
P	mg/dl	4	10	6.6±1.8
ALP	IU/L	190	5487	760±1044
HCO3	mE/l	14	25	20±2.8
CRP	mg/l	2	20	7.4±3.8
BUN	mg/dl	30	180	81±32
Creat	mg/dl	4	15	9.8±2.9
Alb	g/dl	2.4	4.7	3.8±0.50
25-OH Vit D	nMol/l	1.3	105	12.6±21
Chol	mg/dl	59	171	110±33
TG	mg/dl	61	461	130±85
LDL	mg/dl	12	99	61±2
HDL	mg/dl	25	70	38±11

*Duration of hemodialysis treatment

Goldwasser *et al.*, 1993). Low serum albumin as a parameter of malnutrition in HD patients is principally associated with infection/inflammation but not with malnutrition alone (Stenvinkel *et al.*, 2000). Preliminary studies showed the association of CRP level and the serum albumin level (Teruel *et al.*, 2005). Recently in a study on a group of hemodialysis patients no correlation between baseline CRP and serum albumin was found (Nascimento *et al.*, 2004), in this regard we sought to investigate the association of serum CRP level with

Table 3: Mean±SD, Minimum and Maximum of age, duration and dosage hemodialysis and also laboratory results of diabetic hemodialysis patients

Total patients n=11		Min.	Max.	Mean±SD
Age	years	27	75	53±15.8
DH*	months	6	24	14.5±6
Dialysis dose	sessions	54	216	123±54
URR	%	39	75	53.5±9.8
BMI	kg/m ²	20	34	23±3.9
iPTH	Pg/ml	16	860	202±277
Ca	mg/dl	5	10	7.5±1.3
P	mg/dl	3	10	6±2
ALP	IU/L	175	584	327±148
HCO3	mE/l	18	25	20±1.8
CRP	mg/l	4	40	12±10
BUN	mg/dl	30	140	87±36
Creat	mg/dl	3	18	9±4.8
Alb	g/dl	3	4.8	3.8±0.50
25-OH Vit D	nMol/l	1.5	3.6	5.8±10
Chol	mg/dl	60	211	133±49
TG	mg/dl	29	456	130±120
LDL	mg/dl	25	122	62±26
HDL	mg/dl	20	46	35±9.5

*Duration of hemodialysis treatment

some nutritional variables consisting serum albumin in our diabetic and non diabetic end-stage renal failure patients undergoing regular hemodialysis patients.

Materials and Methods

This cross-sectional study was conducted on patients with end-stage renal disease (ESRD), who were undergoing maintenance hemodialysis treatment with acetate basis dialysate and polysulfone membranes. According to the severity of secondary hyperparathyroidism, each patient being treated for secondary hyperparathyroidism was given oral active vitamin D3 (Rocaltrol), calcium carbonate, and Rena-Gel capsules at various doses. According to the severity of anemia, patients were under IV iron therapy with Iron Sucrose (venofer) at various doses after each dialysis session, all patients were under treatments of 6mg folic acid daily, 500mg L-Carnitine daily, oral Vitamin B-complex tablet daily and also 2000U IV Eprex (recombinant human erythropoietin (rHuEPO) unique for each patient after each dialysis session routinely. Exclusion criteria were active or chronic infection. For patients levels of serum predialysis creatinine (Creat), post and predialysis blood urea nitrogen(BUN), Serum calcium (Ca), phosphorus (P), albumin (Alb) and also lipid profile containing serum Triglyceride (TG), Cholesterol (Chol) and high density lipoprotein (HDL-C) as well as serum C-reactive protein (CRP) were measured using standard kits. Also Intact serum PTH (iPTH) was measured by the radioimmunoassay (RIA) method using DSL-8000 of USA (normal range of values is 10-65 pg/ml). Serum 25-hydrox VitD (25-OH Vit D) level (normal range of values is 25 to 125 nmol/l) was




Fig. 1: Near significant difference of CRP between diabetic and non-diabetics.




Fig. 2: Significant difference of CRP between diabetic and non-diabetic of female HD patients.




Fig. 3: Significant difference of CRP between males and female of non-diabetic HD patients.

measured by enzyme-linked immunosorbent assay (ELISA) method using DRG kit of Germany. Plasma HCO_3^- and blood P^{H} was measured by arterial blood gas. For the efficacy of hemodialysis the urea reduction rate (URR) was calculated from pre- and post-blood urea nitrogen (BUN) data (Boag, 1994). The body mass index (BMI) was calculated using weight and height

(kg/m^2) (<http://www.halls.md/body-mass-index/av.htm>). Serum LDL-C was calculated using Friedewald's formula (Friedewald *et al.*, 1972). Duration and dosages of hemodialysis treatment were calculated from the patients' records. The duration of each hemodialysis session was 4 hours. Statistical analysis were performed on total hemodialysis (HD), females, males, diabetics and non diabetics populations separately. For statistical analysis, the data are expressed as the mean \pm SD. Comparison between the groups was done using Student's t-test. Statistical correlations were assessed using partial correlation test. All statistical analyses were performed using SPSS (version 11.5.00). Statistical significance was determined at a p-value < 0.05 .

Results

Total patients were 36 (f = 15 m = 21), consisting of 25 (f = 11 m = 14) non-diabetic HD patients and 11 (f = 4 m = 7) diabetic HD patients. Table 1, 2 and 3 show the patients' mean \pm SD age, the length of time they were on hemodialysis, the dialysis dosage, and the results of laboratory tests. The mean patient's age was 46.6 (± 17) years. In total patients, no significant difference of CRP between males and females of total patients was seen, also no significant difference of CRP between diabetic and non-diabetic male HD patients was seen (p N.S.), however a near significant difference of CRP between diabetic and non-diabetics of total patients was seen (p = 0.056; Fig. 1), also a significant difference of CRP between diabetic and non-diabetic of female HD patients was seen (p < 0.001; Fig. 2), moreover a significant difference of CRP between males and female of non-diabetic population was seen (p = 0.032; Fig. 3). In this study the associations between serum CRP levels and variables consisting of age, duration of hemodialysis, hemodialysis doses, serum lipids, parameters of bone activity, Serum 25-hydroxy Vit D levels, dialysis adequacy and also with plasma HCO_3^- in all groups containing total, non-diabetic, diabetic, female and male HD patients were as follows; In total HD patients, a near significant inverse correlation of serum CRP with serum cholesterol (r = -0.29, p = 0.090; Fig. 4) and a significant positive correlation of serum CRP with serum phosphorus (r = 0.33, p = 0.050; Fig. 5) were found (adjusted for duration of dialysis for two correlations). In this group also a near significant positive correlation of serum CRP with serum BUN (r = 0.33, p = 0.050) (adjusted for dialysis dosage) was found too. In non-diabetic HD patients a significant positive correlation of serum CRP with serum ALP (r = 0.77, p < 0.001; Fig. 6) (adjusted for age) and a near significant inverse correlations of serum CRP with serum Creat (r = -0.38, p = 0.065; Fig. 7) and with dialysis efficacy as determined by URR (r = -0.37, p = 0.074; Fig. 8) were seen too (adjusted for dialysis doses for both two correlations). In diabetic HD patients a significant positive correlation of serum CRP with serum iPTH (r =




Fig. 4: Near significant inverse correlation of serum CRP with serum cholesterol in total HD patients




Fig. 5: Significant positive correlation of serum CRP with serum phosphorus

0.68, $p=0.030$; Fig. 9) (adjusted for dialysis doses), and a near significant inverse correlation of serum CRP with plasma HCO_3^- ($r = -0.52$, $p=0.054$; Fig. 10) (adjusted for duration of dialysis) were found too. More over in this group we found a near significant positive correlation of serum CRP with age ($r = 0.50$, $p=0.065$) (adjusted for duration of dialysis). In male HD patients there was a near significant inverse correlations of serum CRP with serum Tg ($r = -0.40$, $p=0.079$) (adjusted for age), more over we found that the previous near significant inverse correlation of serum CRP with cholesterol in total patients was significantly negative in this population ($r = -0.45$, $p=0.046$) (adjusted for dialysis doses), in this group we also found a near significant positive correlation of serum albumin and plasma HCO_3^- ($r = 0.42$, $p=0.065$, Fig. 11) (adjusted for dialysis duration). In this study no significant association between serum

CRP and serum albumin was seen (p N.S.).

Discussion

In this study we found a near significant difference of CRP between diabetic and non-diabetics of total patients with more values of CRP in diabetics, a significant difference of CRP between diabetic and non-diabetic of female HD patients with more values in diabetics and a significant difference of CRP between males and female of non-diabetic population with more values of CRP in males. we also found inverse correlation of serum CRP with serum cholesterol and triglyceride levels. A significant positive correlation of serum CRP with serum phosphorus and also near significant positive correlations of serum CRP with serum BUN, with serum ALP and serum iPTH, and also with age of patients were found too. Inverse correlations of serum CRP with serum Creat, and with dialysis efficacy and also with plasma HCO_3^- were found too. No significant association between serum CRP and serum albumin was seen. It has recently been recognized evidence of inflammation that predicts serum albumin concentration in dialysis patients (Fernandez-Reyes et al., 2000). Uremic malnutrition and chronic inflammation often coexist in ESRD patients. A recent study by Stevinkel *et al.*, 1999. demonstrated this relationship in advanced chronic kidney disease patients not yet on maintenance dialysis. Of 109 patients with advanced chronic renal failure, 44% had moderate to severe protein-calorie malnutrition, and 32% had signs of inflammation, as determined by elevated CRP concentrations (Stenvinkel *et al.*, 1999). Most importantly, 53% of the patients with malnutrition had signs of inflammation and 72% of the patients with inflammation had signs of malnutrition. A similar association is observed in chronic hemodialysis patients. Qureshi *et al.*, (1998) have demonstrated that chronic hemodialysis patients with severe malnutrition display signs of chronic inflammation (CRP >20 mg/L) approximately four times that of well-nourished chronic hemodialysis patients (Qureshi *et al.*, 1998). Recently, markers of chronic inflammation have also been associated with adverse clinical outcome in chronic renal failure patients. CRP is a significant predictor of mortality as well as morbidity in both chronic hemo- and peritoneal dialysis patients (Ikizler *et al.*, 1999; Zimmermann *et al.*, 1999; Yeun *et al.*, 2000). Similarly, increased levels of proinflammatory cytokines are associated with increased risk of mortality in ESRD patients (Stenvinkel *et al.*, 2002; Bologa *et al.*, 1998). Whereas these two unfavorable conditions independently predispose ESRD patients to increased risk of morbidity and mortality (Warner *et al.*, 2002), when uremic malnutrition and chronic inflammation coexist, they have an exponential relationship to morbidity and mortality, i.e., a small increase in the severity of either condition leads to significant worsening of morbidity and mortality (Caglar *et al.*, 2002). In a study conducted by




Fig. 6: Significant positive correlation of serum CRP with serum ALP




Fig. 7: Near significant inverse correlation of serum CRP with serum Creatinine




Fig. 8: Near significant inverse correlation of serum CRP with dialysis efficacy as determined by URR.

significant correlation between serum CRP and serum albumin neither in total nor in subgroups were found. In accordance with our finding, Nascimento *et al.*, 2004 found that baseline C-reactive (CRP) was not correlated with serum albumin level in the group of haemodialysis (HD) patients that they examined (Nascimento *et al.*, 2004) also Tsirpanlis *et al.*, 2005 in a longitudinal study on a similar patient population, reached to similar finding (Warner *et al.*, 2002). In contrast to previous studies Iseki *et al.* (1999) on 163 dialysis patients that divided into two groups according to their baseline CRP levels, with group 1 consisting of CRP < 10 mg/l (n=128) and group 2 of CRP ≥ 10 mg/l (n=35), found significant lower serum albumin in group two. Furthermore, the risk of death was significantly higher in group 2, Iseki concluded that CRP is a significant predictor of death in chronic dialysis patients, independent of serum albumin and other possible confounders (Tsirpanlis *et al.*, 2005). In contrast to the finding of this work, Panichi *et al.* (2001) conducted a study on 102 chronic pre-dialysis patients whose mean age was 53±5.8 years with a mean creatinine clearance of 52±37 ml/min. While serum CRP was greater in patients with lower creatinine clearance, the serum albumin levels were not significantly different between the groups of patients with a creatinine clearance lower than 20 mL/min (n=32) and patients with a creatinine clearance higher than 20 mL/min (n = 70) (Iseki *et al.*, 1999). In hemodialysis patients serum triglyceride and cholesterol are considered as nutritional parameters (Fernandez-Reyes *et al.*, 2000), their inverse correlation with CRP showed the effect of inflammatory state of uremia and its negative effects on lipids (Panichi *et al.*, 2001). To support of our results, Fernandez-Reyes *et al.* (2000) in a cross sectional study which was performed in 64 patients (35 males, 13% diabetics; mean age 64±12), who had been on HD for 64±58 months showed the correlation of low serum cholesterol and high level of CRP (Fernandez-Reyes *et al.*, 2000). We also showed negative correlation of dialysis adequacy with CRP, which show that in an adequate dialysis, will be accompanied with lower stimulation of proinflammatory mediators. Parathyroid hormone (PTH) promotes IL-6 secretion by osteoblasts, and may also up-regulate IL-6 production in the liver and adipose tissue; this may explain why serum IL-6 is markedly elevated in primary hyperparathyroidism, and low in hypoparathyroidism. IL-6 is the chief stimulus to hepatic production of many acute phase reactants, notably fibrinogen and C-reactive protein as mentioned (Liu *et al.*, 2004). In hemodialysis, hyperphosphatemia is accompanied by uncontrolled secondary hyperparathyroidism (SHPTH) and high serum PTH level, hence association of hyperphosphatemia and high iPTH with serum CRP in this study further support the previous confirmed data (Liu *et al.*, 2004), which explain the need to better control of SHPTH in these patients. In this regard, *in vitro*

Fujino *et al.* on 389 HD patients, significant negative correlations were seen between fat mass changes and CRP was mentioned (Fujino *et al.*, 2005). In our study no




Fig. 9: Significant positive correlation of serum CRP with serum iPTH.




Fig. 10: Near significant inverse correlation of serum CRP with plasma HCO₃⁻




Fig. 11: Near significant positive correlation of serum albumin and plasma HCO₃⁻

studies have shown that 1, 25 dihydroxyvitamin D3 [1,25(OH)₂D₃] decreases cytokine production by monocytes and lymphocytes. In addition, intravenous or

oral pulse calcitriol treatment suppresses interleukin 6 (IL6) and interleukin1beta (IL1 beta) in hemodialysis patients (McCarty, 2005). In this study we also found an inverse association of plasma HCO₃⁻ with serum CRP. Metabolic acidosis has been suggested to be an important factor responsible for protein and caloric malnutrition (Gannage-Yared *et al.*, 2003; Mitch, 1996; Gennari, 1985). In contrast to the metabolic studies, many epidemiologic studies in maintenance dialysis patients have indicated a paradoxically inverse association between mildly decreased serum bicarbonate and improved markers of protein-energy nutritional state (Movilli *et al.*, 1998; Kalantar-Zadeh *et al.*, 2004). Hence metabolic acidosis may be considered as yet another element of the reverse epidemiology in ESRD patients, to support this finding we could show the positive association of plasma HCO₃⁻ with serum albumin in our male population. Taken together, we could not show the association of serum albumin with serum CRP level, we showed the positive correlation of SHPTH with inflammation implies further need to control of hyperphosphatemia in HD patients, also inverse correlation of serum CRP with cholesterol and triglyceride further support the malnutrition-inflammation complex syndrome (MICS) which frequently seen in HD patients.

Acknowledgements

We would like to thank Dr. K. Kalantar-Zadeh, Assistant Professor of Medicine of UCLA School of Medicine for sending his articles to our center and his comments .

References

Boag, J.T., 1994. Basic Truths in Optimal Hemodialysis, Dialysis & Transplantation, Volume 23, Number 11, Page 636.

Bologa, R.M., D.M. Levine and T.S. Parker, 1998. Interleukin-6 predicts hypoalbuminemia, hypocholesterolemia, and mortality in hemodialysis patients. *Am. J. Kidney Dis.*, 32: 107-114.

Caglar, K., R.M. Hakim and T.A. Ikizler, 2002. Approaches to the reversal of malnutrition, inflammation, and atherosclerosis in end-stage renal disease. *Nutr. Rev.*, 60: 378-87.

Cianciaruso, B., G. Brunori, J.D. Kopple, G. Traverso, G. Panarello, G. Enia, P. Strippoli, A. De Vecchi, M. Querques, G. Viglino, E. Vonesh and R. Maiorca, 1995. Cross-sectional comparison of malnutrition in continuous ambulatory peritoneal dialysis and hemodialysis patients. *Am. J. Kidney Dis.*, 26: 475-486.

Clyne, B. and J.S. Olshaker, 1999. The C-reactive protein. *J. Emerg. Med.*, 17: 1019 -1025.

Enia, G., C. Sicuso, G. Alati and C. Zocalli, 1993. Subjective global assessment of nutrition in dialysis patients. *Nephrol. Dial. Transplant.*, 8: 1094-1098.

- Fernandez-Reyes, M.J., F. Alvarez-Ude, R. Sanchez, C. Mon, P. Iglesias and A. Vazquez, 2000. Nutritional status, comorbidity, and inflammation in hemodialysis. *Nefrologia*, 20: 540-9.
- Friedewald, W.T., R. Levy and D.S. Fredrickson, 1972. Estimation of the concentration of Low-density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge. *Clin. Chem.*, 18: 799-502.
- Fujino, Y., E. Ishimura, S. Okuno, N. Tsuboniwa, K. Maekawa and T. Izumotani, 2005. C-reactive protein is a significant predictor of decrease in fat mass in hemodialysis patients. *Biomed. Pharmacother.*, 13.
- Gannage-Yared, M.H., M. Azoury, I. Mansour, R. Baddoura, G. Halaby and R. Naaman, 2003. Effects of a short-term calcium and vitamin D treatment on serum cytokines, bone markers, insulin and lipid concentrations in healthy post-menopausal women. *J Endocrinol. Invest.*, 26: 748-53.
- Gennari, F., 1985. Acid-base balance in dialysis patients. *Kidney Int.*, 28: 678-688.
- Goldwasser, P., N. Mittman, A. Antignani, D. Burrell, M.A. Michel, J. Collier and M.M. Avram, 1993. Predictors of mortality in hemodialysis patients. *J. Am. Soc. Nephrol.*, 3: 1613-1622.
- Haubitz, M., R. Brunkhorst, E. Wrenger, P. Froese, M. Schulze and K.M. Koch, 1996. Chronic induction of C-reactive protein by hemodialysis, but not by peritoneal dialysis therapy. *Perit. Dial. Int.*, 16: 158-162.
- Ikizler, T.A., R.L. Wingard, J. Harvell, Y. Shyr and R.M. Hakim, 1999. Association of morbidity with markers of nutrition and inflammation in chronic hemodialysis patients: a prospective study. *Kidney Int.*, 55: 1945-1951.
- Iseki, K., M. Tozawa, S. Yoshi and K. Fukiyama, 1999. Serum C-reactive protein (CRP) and risk of death in chronic dialysis patients. *Nephrol. Dial. Transplant.*, 14: 1956-1960.
- Kalantar-Zadeh, K., R. Mehrotra, D. Fouque and J.D. Kopple, 2004. Metabolic acidosis and malnutrition-inflammation complex syndrome in chronic renal failure. *Semin Dial.*, 17: 455-65.
- Kim, B.S., D.S. Jeon, M.J. Shin, Y.O. Kim, H.C. Song and S.H. Lee, 2005. Persistent Elevation of C-Reactive Protein May Predict Cardiac Hypertrophy and Dysfunction in Patients Maintained on Hemodialysis. *Am. J. Nephrol.*, 29: 25: 189-195.
- Leavey, S.F., R.L. Strawderman, C.A. Jones, F.K. Port and P.J. Held, 1998. Simple nutritional indicators as independent predictors of mortality in hemodialysis patients. *Am. J. Kidney Dis.*, 31: 997-1006.
- Lowrie, E.G. and N.L. Lew, 1990. Death risk in hemodialysis patients: The predictive value of commonly measured variables and an evaluation of death rate differences between facilities. *Am. J. Kidney Dis.*, 15: 458-482.
- Liu, Y., J. Coresh, J.A. Eustace, J.C. Longenecker, B. Jaar and N.E. Fink, 2004. Association between cholesterol level and mortality in dialysis patients: role of inflammation and malnutrition. *JAMA.*, 28: 291: 451-9.
- Marckmann, P., 1988. Nutritional status of patients on hemodialysis and peritoneal dialysis. *Clin. Nephrol.*, 29: 75-78.
- McCarty, M.F., 2005. Secondary hyperparathyroidism promotes the acute phase response -- a rationale for supplemental vitamin D in prevention of vascular events in the elderly. *Med Hypotheses*, 64: 1022-6.
- McIntyre, C., I. Harper, I.C. Macdougall, A.E. Raine, A. Williams and L.R. Baker, 1997. Serum C-reactive protein as a marker for infection and inflammation in regular dialysis patients. *Clin. Nephrol.*, 48: 371-374.
- Mitch, W.E., 1996. Metabolic acidosis stimulates protein metabolism in uremia. *Miner Electrolyte Metab.*, 22: 62-65.
- Morrow, D.A. and P.M. Ridker, 2000. C-reactive protein, inflammation, and coronary risk. *Med. Clin. North Am.*, 84: 149-161.
- Movilli, E., R. Zani and O. Carli, 1998. Correction of metabolic acidosis increases serum albumin concentrations and decreases kinetically evaluated protein intake in haemodialysis patients: a prospective study. *Nephrol. Dial. Transplant.*, 13: 1719-1724.
- Nascimento, M.M., R. Pecoits-Filho, A.R. Qureshi, S.Y. Hayashi, R.C. Manfro and M.A. Pachaly, 2004. Prognostic impact of fluctuating levels of C-reactive protein in Brazilian haemodialysis patients: a prospective study. *Nephrol. Dial. Transplant.*, 19: 2803-9.
- Ortega, O., I. Rodriguez, P. Gallar, A. Carreno, M. Ortiz and B. Espejo, 2002. Significance of high C-reactive protein levels in pre-dialysis patients. *Nephrol. Dial. Transplant.*, 17: 1105-1109.
- Panichi, V., M. Migliori, S. De Pietro, M.R. Metelli, D. Taccola and R. Perez, 2000. Plasma C-Reactive Protein in Hemodialysis Patients: A Cross-Sectional, Longitudinal Clinical Survey. *Blood Purification*, 18: 30-36.
- Panichi, V., M. Migliori, S. De Pietro, D. Taccola, A.M. Bianchi and M. Norpoth, 2001. C reactive protein in patients with chronic renal diseases. *Ren. Fail.*, 23: 551-62.
- Qureshi, A.R., A. Alvestrand and A. Danielsson, 1998. Factors predicting malnutrition in hemodialysis patients: a cross-sectional study. *Kidney Int.*, 53: 773-782.
- Stenvinkel, P., P. Barany, O. Heimbürger, R. Pecoits-Filho and B. Lindholm, 2002. Mortality, malnutrition, and atherosclerosis in ESRD: what is the role of interleukin-6? *Kidney Int.*, 61: 103-108.

Baradaran and Nasri: Association of Serum C-Reactive Protein with Some Nutritional Parameters

- Stenvinkel, P., O. Heimbürger and F. Paultre, 1999. Strong association between malnutrition, inflammation, and atherosclerosis in chronic renal failure. *Kidney Int.* 55: 1899-1911.
- Stenvinkel, P., O. Heimbürger, B. Lindholm, G.A. Kaysen and J. Bergström, 2000. Are there two types of malnutrition in chronic renal failure? Evidence for relationships between malnutrition, inflammation and atherosclerosis (MIA syndrome). *Nephrol. Dial. Transplant.*, 15: 953-60.
- Stenvinkel, P., O. Heimbürger, B. Lindholm, G.A. Kaysen and J. Bergström, 2000. Are there two types of malnutrition in chronic renal failure? Evidence for relationships between malnutrition, inflammation and atherosclerosis (MIA syndrome) *Nephrol. Dial. Transplant.*, 15: 953-960.
- Teruel, J.L., R. Marcen, J. Ocana, M. Fernandez-Lucas, M. Rivera, G. Taberero and J. Ortuno, 2005. Clinical Significance of C-Reactive Protein in Patients on Hemodialysis: A Longitudinal Study. *Nephron. Clin. Pract.*, 25; 100: c140-c145.
- Tsirpanlis, G., P. Bagos, D. Ioannou, A. Bleta, I. Marinou and A. Lagouranis, 2005. Serum albumin: a late-reacting negative acute-phase protein in clinically evident inflammation in dialysis patients. *Nephrol Dial. Transplant.*, 20: 658-9.
- Warner, C., J. Zimmermann, S. Schwedler and T. Metzger, 2002 Inflammation and cardiovascular risk in dialysis patients. *Kidney Int.*, 61: 99-102.
- Wolbink, G.J., M.C. Brouwer, S. Buysmann, I.J. ten Berge and C.E. Hack, 1996. CRP-mediated activation of complement *in vivo*: assessment by measuring circulating complement-C-reactive protein complexes. *J. Immunol.*, 157: 473-9.
- Yeun, J.Y., R.A. Levine, V. Mantadilok and G.A. Kaysen, 2000. C-Reactive protein predicts all-cause and cardiovascular mortality in hemodialysis patients. *Am. J. Kidney Dis.*, 35: 469-476.
- Zimmermann, J., S. Herrlinger, A. Pruy, T. Metzger and C. Wanner, 1999. Inflammation enhances cardiovascular risk and mortality in hemodialysis patients. *Kidney Int.*, 55: 648-658.