

**PJN**

ISSN 1680-5194

PAKISTAN JOURNAL OF  
**NUTRITION**

**ANSI***net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan  
Mob: +92 300 3008585, Fax: +92 41 8815544  
E-mail: [editorpjn@gmail.com](mailto:editorpjn@gmail.com)

## Antimicrobial Activity of Lactic Acid Bacteria Isolated from *Budu* of West Sumatera to Food Biopreservatives

Yusra, Fauzan Azima, Novelina<sup>1</sup> and Periadnadi<sup>2</sup>

<sup>1</sup>Department of Agricultural Processing Technology,  
Faculty of Agricultural Technology, Andalas University, Padang, 25163, Indonesia

<sup>2</sup>Department of Biology, Faculty of Mathematics and Natural Sciences,  
Andalas University, Padang, 25163, Indonesia

**Abstract:** *Budu* is a traditional fermented fish product which are produced and distributed within a specific area in the province of West Sumatera prepared from Spanish mackerel (*Scomberomorus guttatus*). The samples were obtained from the small scale domestic factory. The aim of this research are to isolate and identify Lactic Acid Bacteria (LAB) activity from *budu* with pathogen bacteria (*Escherichia coli*, *Staphylococcus aureus*, *Salmonella thypi*, *Bacillus subtilis* and *Listeria monocytogenes*) followed identified by PCR. The bacteria isolated were growth on medium Glucose Trypton Agar (GTA) supplement with CaCO<sub>3</sub> and then performed purification by plate out on *deMan Rogosa and Sharpe* (MRS) Agar. It was found 138 isolates of LAB showed with clear zone around the culture and 8 isolates (B2, B4, B5, B9, B11, B13, B28 and B34) has been antimicrobial activity against the growth of pathogenic bacteria. The results showed that isolates B28 had the highest antimicrobial activity against all bacteria test, with a range of inhibition zone 14-35 mm, gram positive, spore former bacil, motility and catalase positive. Based on morphological examination and PCR analysis, the isolate B28 was primarily identified as *Bacillus cereus* strain HVR22 bacteria.

**Key words:** Antimicrobial activity, *Budu*, isolated, food biopreservatives, West Sumatera

### INTRODUCTION

Antimicrobial agent is a general term used to refer to any compound which include antibiotics, food antimicrobial agents, sanitizer, disinfectants and other substances that acts against microorganisms (Katzung, 2004). Antimicrobial agents can be produced by fermentation products such as milk, vegetables and fish, such: *plasom* of Thailand (Kopermsub and Yunchalard, 2010) (Hwanhlem *et al.*, 2011), *bakasang* of Indonesia (Ijong, and Ohta, 1996) and *Budu* of Malaysia (Liasi *et al.*, 2009) and *budu* from West Sumatera.

*Budu* from West Sumatera differs from the Malaysian *budu*, not only in the type of fish but also in the processing. In Malaysia *budu* is produced using anchovies (*Stolephorus* sp.), the small sized marine fish locally known as *ikan bilis*, as the raw material. Anchovies are mixed with coarse salt at a ratio of 3:2 and fermented a large covered earthen container, concrete tank or black polyethylene tank for more than 8 months in an unshaded open area. After fermentation is complete, anchovy fries is naturally degraded into a dark brown liquid (Huda and Rosma, 2006). This fries mixed with tamarind and coconut sugar in salt solution (15%) and then fermented at room temperature in large earthen ware post (Liasi *et al.*, 2009).

The *budu* from West Sumatera is a fermented fish product mainly in the coastal areas such a Pariaman,

Tiku and Pasaman. *Budu* normally made from bigger sized marine fish such as Spanish mackerel (*Scomberomorus* sp.) and leatherskin (*Chorinemus* sp.), locally, known as *ikan tenggiri* and *ikan talang*. Reported that traditional processing of West Sumatera *budu* starts with hanging the fresh fish by its caudal fin at room temperature for 30 h. Once the fish is marked by a rift bulging fish meat, the fish is cut into a butterfly style and all of the intestinal organs and gills are removed. The fish is then washed to ensure that it is free of blood and other intestinal residues. The fish layered with coarse salt at a ratio of 1:5-1:3 for 3-4 h. After which the fishes rinsed to remove the excess salt. The fish is then sun-dried for 4-5 days.

Antimicrobials have been used increasingly as a primary intervention for inhibition or activation of pathogenic microorganisms in foods (Davidson and Zivnovic, 2003). Generally, in fermented foods, Lactic Acid Bacteria (LAB) display numerous antimicrobial activities. This is mainly due to the production of antimicrobial metabolites including lactic acid, hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), diasetil, carbon dioxide (CO<sub>2</sub>) and bacteriocins and antifungal peptides.

Bacteriocins are the peptides and protein antibiotics which are produced by several species and have antimicrobial properties usually against other closely related species (Olivera *et al.*, 2004). These

antimicrobial peptides are gaining more and more attention not only as an alternative therapeutic agent for the prevention and treatment of infections but also as preservatives in food industries to avoid deterioration and spoilage of food (Anthony *et al.*, 2009). Bacteriocins are generally recognized as naturally occurring food preservatives able to influence the quality and safety of foods (Settanni and Corsetti, 2008). The aims of this research are to isolate and identify lactic acid bacteria from traditional fermented fish product "budu" and to screen and determine the activity antimicrobial with pathogen bacteria (*Escherichia coli*, *Staphylococcus aureus*, *Salmonella thypi*, *Bacillus subtilis* and *Listeria monocytogenes*) in order to apply for food biopreservatives.

## MATERIALS AND METHODS

Three samples of budu were collected in sterile plastics from manufactures small scale domestic factory in West Sumatera (Sirah river and Gasan district Padang Pariaman and Sasak district Pasaman). The samples were transferred immediately to the laboratory for microbiological analysis.

**Pathogenic bacteria strains and culture medium:** *Escherichia coli*, *Staphylococcus aureus*, *Salmonella thypi*, *Bacillus subtilis* and *Listeria monocytogenes*. The culture medium in this research: GTA+CaCO<sub>3</sub> plate a consisted of 20 gr glucose, 2, 5 gr tripton, 20 gr bacto agar (Oxoid) and 15 gr CaCO<sub>3</sub>, MRS broth (Merck) and Nutrient Agar (NA) (Merck).

**Isolation of lactic acid bacteria:** At first, the samples were homogenized in mortar and pestle, taking 10 g of well mixed sample blended in 90 ml of 0, 85% (w/v NaCl) physiological saline. Further serial dilutions to 10<sup>-8</sup>. For isolation of LAB, of each dilute 0, 1 ml inocula were plated out GTA+CaCO<sub>3</sub> and the plates were incubated at room temperature for 48-72 h. The existence of clear zone around the colonies suspected to be the lactic acid bacteria. Colonies were selected at random from plates and the isolates purified by repeated streaking onto MRS agar. Purified strains of LAB were inoculated into MRS broth (pH 6.5) and incubated 24 h at 30°C. All purified strains were kept in MRS agar containing 20% glycerol at -20°C (Modified Rahayu and Margino, 1997).

**Characterization of lactic acid bacteria:** The identified of the isolates were determined by the standard procedure of gram staining, catalase test, motility and spore former test (Hadjoetomo, 1985; Fardiaz, 1989; Lay, 1994). Characterization bacterial strains of LAB determination the using standard methods "Manual for the identification of medical bacteria" (Cowan and Steel's, 1975).

**Antimicrobial activity test:** The modified methods of Savodogo *et al.* (2004) and Girum *et al.* (2005) were used to determine the antibacterial activities of the isolates. A single isolated colonies were selected from MRS agar plates and transferred to grow in sterile MRS broth. The broth culture was incubated erobically at 37°C for 48 h. After incubation, the culture was centrifuged at 10.000 rpm for 20 minute at 4°C to obtain the culture supernatant. The indicator microorganisms (*Escherichia coli*, *Staphylococcus aureus*, *Salmonella thypi*, *Bacillus subtilis* and *Listeria monocytogenes*) were grown in NA for 24 h at 37°C. A sterile cotton swab was dipped into the culture of the indicator microorganisms and rotated several times and the swab was then pressed firmly on the inside wall of the tube above the fluid level to remove excess inoculums. The dried surface of NA was inoculated by streaking the swab over the entire agar surface. This procedure was repeated by streaking two or more times while rotating the plate each time to ensure an even distribution of inoculums. For the bioassay, the sterile filter disc was dipped into the culture supernatant and touched to the side of container to remove excess liquid and it was then placed on a NA plate. After 24 hours at 37°C of aerobic incubation, each plate was evaluated and the diameters of the inhibition zones, including the diameter of the disc, were measured using a transparent ruler (Assefa *et al.*, 2008).

**DNA extract, PCR and sequencing:** The Genomic DNA was extracted from pure culture of the *Bacillus* isolate using genomic DNA extraction kit following instructions of the Pitcher *et al.* (1989, modified) (White *et al.*, 1990). The 1.5 kb 16S rDNA gene were amplified by PCR using a pair of universal bacterial 16S rDNA gene primers 9F: 5-AAG GAG GTG ATC CAG CC-3 and primers1541 R: 5-GAG TTT GAT CCT GGC TCA G-3. The PCR was carried out according to (Hiraishi *et al.*, 1995). To sequence part of the 16S-rRNA genes, the 16S-rRNA genes were amplified by PCR using specific primers of 63f and 1387r from genomic DNA (200 ng) on Ready-To-GO PCR Beads (Pharmacia-Biotech, Uppsala, Sweden). Phenol-chloroform-isoamyl alcohol (25:24:1) treatment, ethanol precipitation and agarose gel electrophoresis were used to purify the genomic DNA. Total volume of the PCR reaction (25 ul) consisted of 1.5 U *Tag* DNA Polymerase, 10mM Tris-HCl (pH 9 at room temperature), 50 mM KC1, 1.5 mM MgCl<sub>2</sub>, 200 uM of each dNTPs and stabilizer including BSA. The reaction was incubated in a Gene Amp PCR System 2400 Thermocycler (Perkin-Elmer Cetus, Norwalk, Conn.). Part of 16S-rRNA gene was sequenced to infer the closest related organism from the Ribosomal Database Project (RDP) maintained at the University of Illinois, Urbana-Champaign. The sequencing reactions were done by using the Big Dye Ready Reaction Dye Deoxy Terminator kit, purified by

ethanol-sodium acetate precipitation. The reactions were run on an ABI PRISM 3130 Genetic Analyzer (PE Applied Biosystems, Foster City, CA).

## RESULTS AND DISCUSSION

**Isolation of LAB:** The isolation of LAB from budu was performed GTA supplemented with CaCO<sub>3</sub> which was obtained at the room temperature, 72 h of incubation and MRS agar was used as a preliminary screening medium for LAB. It was found that 138 isolates exhibited a clear zone and growth on MRS agar. Calcium carbonate was used as an indicator for acid-producing strains since it dissolved when interact with acid then a clear zone is observed. All of isolates were identified as LAB using the criteria of being gram positive and catalase positive. Of these, 135 isolates (98%) from budu were identified as rod shape and 3 (2%) isolates were identified as cocci shape.

In the production of fermented foods, the initial food matrix presents abiotic and biotic conditions that select for the growth of specific microbial communities (Giraffa, 2004). In the case of *budu*, the initial concentration of salt select for the growth of lactic acid bacteria. Hwanhlem *et al.* (2011) reported of lactic acid bacteria from Thailand fermented fish (*plasom*) were identified 133 isolates, 25 isolates were cocci, 75 isolates were short rods and 33 isolates were rods. Tanasupawat *et al.* (2009) observed from fish sauce (*nam-pla*) in Thailand have fifteen isolates were moderately rod-shaped halophilic extremely, rod-shaped halophilic bacteria and sphere shaped bacterium. Kim *et al.* (2002) isolated *Bacillus* sp. SFF34, from Korean traditional fermented flatfish. Anihouvi *et al.* (2006) from *lanhouin*, a fermented fish product in Republic Benin get *Micrococci* and *Bacilli* were the predominant organisms enumerated in product.

**Detection of antagonistic activity:** A total 138 isolated randomly picked from *budu* samples for the morphological identification was selected isolated gave 8 (B2, B4, B5, B9, B11, B13, B28 and B34) isolates has the most extensive zone of inhibition when compared to other isolates. Based on the data it appears that the fluid extracellular of eight isolates of the inhibition of the bacterial pathogen/indicator and it appears that the inhibitory activity extracellular B4 and B28 had a higher power resistor. While isolates had the highest inhibitory power is B28 isolates with inhibition of the *Escherichia coli* 14 mm, against *Staphylococcus aureus* 18 mm, against *Salmonella thypi* 15 mm, against *Bacillus subtilis* 35 mm and the *Listeria monocytogenes* 19 mm (Table 1).

Isolates that have the widest zone of inhibition against *Escherichia coli* isolates B4 which is 20 mm with inhibitory activity mm<sup>2</sup>/ml 14287, against *Staphylococcus aureus* isolates B4 is the inhibition zone diameter of 24

Table 1: Antibacterial activity of 8 isolates of LAB obtained from *budu* samples against different indicator microorganisms

No. of strains	Diameter of inhibition zone (mm)				
	<i>E. coli</i>	<i>S. aureus</i>	<i>S. thypi</i>	<i>B. subtilis</i>	<i>L. monocytogenes</i>
B2	13	20	15	20	20
B4	20	24	13	20	18
B5	12	21	15	19	20
B9	12	16	11	20	20
B11	12	16	15	23	20
B13	12	13	12	35	15
B28	14	18	15	35	19
B34	10	13	10	33	20

mm with inhibition activity 21217 mm<sup>2</sup>/ml, the bacteria *Salmonella thypi* is isolate B2, B5, B11 and B28 that is measuring 15 mm by 7418.25 mm<sup>2</sup>/ml inhibition activity, the bacterium *Bacillus subtilis* is isolat B13 and B28 that is inhibition zone diameter 35 mm with inhibition activity 46668.250 mm<sup>2</sup>/ml and the bacteria *Listeria monocytogenes* is isolate B2, B5, B9, B11 and B34 that is 20 mm wide with 14287 mm<sup>2</sup>/ml inhibition activity. The results showed an average inhibition zone of *Escherichia coli* ranging from 10 mm (isolate B34) to 20 mm (isolate B4), *Staphylococcus aureus* ranging from 13 mm (isolate B13 and B34) to 24 mm (isolate B4), *Salmonella thypi* ranging from 10 mm (isolate B34) to 15 mm (isolate B2, B5, B11 and B28), *Bacillus subtilis* ranging from 19 mm (isolate B5) to 35 mm (isolate B13 and B28) and *Listeria monocytogenes* ranging from 15 mm for isolate B13 to 20 mm for isolate B2, B5, B9, B11 and B34.

According to Ammor *et al.* (2006) LAB produce a wide range of products from low molecular mass compounds, such as hydrogen peroxide, carbon dioxide and diacetyl, to high molecular mass compounds, such as bacteriocins. Organic acid produced by LAB leads to a reduction in pH levels and increases the production of hydrogen peroxide (Ponce *et al.*, 2008), enzymes (lactoperoxidase system with hydrogen peroxide and lysozyme), low-molecular metabolites (reuter in, diacetyl and fatty acids) and bacteriocins (nisin and others) (Holzapfel *et al.*, 1995). These products exhibit antibacterial activity against various pathogenic microorganisms, including gram-positive and gram-negative bacteria (Maragkoudakis *et al.*, 2009).

Savodogo *et al.* (2004) isolated eighty strains of lactic acid bacteria producing bacteriocin were isolated from Burkina Faso fermented milk samples. These strains were identified to species *Lactobacillus fermentum*, *Pediococcus* sp., *Leuconostoc mesenteroides* and *Lactococcus*. Isolated bacteriocin exhibited antibacterial activity against *Enterococcus faecalis*, *Bacillus cereus*, *Staphylococcus aureus* and *Escherichia coli* using the agar drop diffusion test. The inhibition diameters obtained with bacteriocin are between 8 and 12 mm. Gram positive indicator bacteria were most inhibited. Ogunshe *et al.* (2007) isolated 50 bacteriocin producing

*Lactobacillus* strains from some Nigerian indigenous fermented foods and beverages and characterized as *Lactobacillus acidophilus*, *Lactobacillus casei*, *Lactobacillus fermentum*, *Lactobacillus lactis* and *Lactobacillus plantarum* were screened for these inhibitory potentials against food borne pathogenic from the same or similar to fermented food sources and against clinical indicator bacterial isolates. De Vuyst and

Leroy (2007) described the production, purification and food applications of bacteriocins from lactic acid bacteria. In fermented foods, lactic acid bacteria display numerous antimicrobial activities due to the production of organic acids and other compounds such as bacteriocins.

**Morphological, physiological and biochemical examination of isolate B28 bacteria:** Bacteria isolated from the *budu* (sample B28) was identified as *Bacillus* sp. on physiological and biochemical characteristics (Table 2). The isolate B28 as gram positive, rod shape and positive for catalase test having smooth round colonies on the MRS media. The strain was capable of fermenting sugars, namely glucose and sucrose. These tests based on Manual for the Identification of Medical Bacteria (Cowan and Steel's, 1975) and the bacteria belong to *Bacillus* genus Phylogenetically, bacteria

Test	Isolate B28
Colony morphology	Creamy, little sticks and Smooth round colonies
Gram staining	Gram positive, rod
Spore-former	Bacil
Coloni Surface	Convex
Coloni size	2 mm
Motility	Positive
Growth in MRS broth	Uniform turbidity
Catalase	Positive
Oxidase	Negative




Fig. 1(a-e): The antimicrobial activity of *Bacillus cereus* strain HVR22 against pathogens bacteria:(a) *Escherichia coli*, (b) *Staphylococcus aureus*, (c) *Salmonella thypi*, (d) *Bacillus subtilis* and (e) *Listeria monocytogenes*, by agar well diffusion methods

```

GGCGGCGTGCCTAATACATGCAAGTCGAGCGAATGGATTAAGAGCTTGCTCTTATGAAGT
TAGCGGCGGACGGGTGAGTAACACGTGGGTAACTGCCATAAGACTGGGATAACTCCGG
GAAACCGGGGCTAATACCGGATAACATTTTGAACCGCATGGTTCGAAATTGAAAGCGGC
TTCGGCTGTCACTTATGGATGGACCCGCGTCGCATTAGCTAGTTGGTGAGGTAACGGCTC
ACCAAGGCAACGATGCGTAGCCGACCTGAGAGGGTGATCGGCCACACTGGGACTGAGACA
CGGCCAGACTCCTACGGGAGGCGAGCAGTAGGGAATCTTCGCAATGGACGAAAAGTCTGA
CGGAGCAACGCCGCGTGAGTGATGAAGGCTTTCGGGTCGTAATACTCTGTTGTTAGGGAA
GAACAAGTGCTAGTTGAATAAGCTGGCACCTTGACGGTACCTAACCAGAAAGCCACGGCT
AACTACGTGCCAGCAGCCGCGTAATACGTAGGTGGCAAGCGTTATCCGGAATTATTGGG
CGTAAAGCGCGCGCAGGTGGTTTCTTAAGTCTGATGTGAAAGCCCACGGCTCAACCGTGG
AGGGTCATTGGAACTGGGAGACTTGAGTGACAGAAGAGGAAAGTGGAAATCCATGTGTAG
CGGTGAAATGCGTAGAGATATGGAGGAACACCACTGGCGAAGGCGACTTTCTGGTCTGTA
ACTGACACTGAGGCGCGAAAAGCGTGGGGAGCAACAGGATTAGATACCCTGGTAGTCCAC
GCCGTAACGATGAGTGCTAAGTGTTAGAGGGTTTCCGCCCTTATGTGCTGAAGTTAACG
CATTAAAGCACTCCGCCTGGGAGTACGGCCGCAAGGCTGAAACTCAAAGGAATTGACGGG
GGCCCGCACAAAGCGGTGGAGCATGTGGTTAATTCGAAGCAACGCCGAAGAACCCTTACCAG
GTCTTGACATCCTCTGAAAACCCTAGAGATAGGGCTTCTCCTTCGGGAGCAGAGTGACAG
GTGGTGATGGTTGTCGTACGCTCGTGTGAGATGTTGGGTTAAGTCCCACAACGAGC
GCAACCCCTGATCTTAGTTGCCATCATTAAAGTTGGGCACTCTAAGGTGACTGCCGGTGAC
AAACCGGAGGAAGGTGGGGATGACGTCAAATCATCATGCCCTTATGACCTGGGCTACAC
ACGTGCTACAATGGACGGTACAAAGAGCTGCAAGACCGCGAGGTGGAGCTAATCTCATAA
AACCGTTCTCAGTTCGGATTGTAGGCTGCAACTCGCCTACATGAAGCTGGAATCGCTAGT
AATCGCGGATCAGCATGCCGCGGTGAATACGTTCCCGGGCCTGTACACACCGCCCGTCA
CACCACGAGAGTTTGTAAACCCGAAGTCGGTGGGGTAACC
    
```

Fig. 2: Nucleotide sequence of *Bacillus* sp B28 by 16S rDNA sequence (full sequence; 1,462 base pairs)

belonging to the genus *Bacillus* belong to class I of the phylum Firmicutes i.e., the bacilli. Members of the genus *Bacillus* are gram-positive, aerobic and endospore-forming bacteria that are characterized by their rod-shaped cell morphology, catalase production and their ubiquitous distribution. They are found in diverse environments such as soil and clays, rocks, dust, aquatic environments, vegetation, food and the gastrointestinal tracts of various insects and animals (Nicholson, 2002).

Members of the *Bacillus* group *sensu-lato* are considered good producers of antimicrobial substances, including peptide and lipopeptide antibiotics and bacteriocins (Stein, 2005). The production of antimicrobial substances and sporulation capacity confer *Bacillus* strains with a double advantage in terms of their survival in different habitats. The presence of *Bacillus* species in food does not always imply spoilage or food poisoning and some species or strains are even used in human and animal food production such as, for example, *Bacillus subtilis* strains that are used in Natto, an East Asian fermented food production (Hosoi and Kiuchi, 2003). Furthermore, specific *B. subtilis* strains are also used as a starter culture for fermenting soybeans into the traditional West African condiment dawadawa (Terlabie *et al.*, 2006) or for fermenting African mesquite seeds in the production of the Nigerian food condiment okpehe (Oguntoyinbo *et al.*, 2007). A nontoxigenic *Bacillus cereus* ssp. *toyoi* with probiotic properties is also used as an animal feed additive (Lodemann *et al.*, 2008).

*Bacillus* is an interesting genus to be investigated for antimicrobial activity because *Bacillus* sp. produces a large number of peptides with biological activities. e.g., cerecin 7 (Oscariz *et al.*, 1999) produced by *B. cereus* BC7, tochicin (Paik *et al.*, 1997) from *B. thuringensis*, thuricin 7 (Cherif *et al.*, 2003), subpeptin JM4-A and subpeptin JM4-B produced by *B. subtilis* JM4 (Wu *et al.*, 2005). The chemical and physical diversity of peptide antibiotics makes them ideal candidates not only for therapeutic applications but also in other areas, especially the agri-food industry.

#### Identification of lactic acid bacteria by 16SrRNA:

Methods for the detection and identification of *B. cereus* are e.g., serotyping, pyrolytic gas chromatography, pyrolytic mass spectrometry, ribotyping, phage typing, plasmid profiles, electrophoresis in pulse electric field and Polymerase Chain Reaction (PCR) using genera-specific and species-specific primers. Isolate B28 was DNA extraction for GES methods (Pitcher *et al.*, 1989; Modified), amplification PCR of its 16S rDNA with specific primers 9F: 5'-AAG GAG GTG ATC CAG CC-3' and primers 1541 R: 5'-GAG TTT GAT CGT GGC TCA G-3' (White *et al.*, 1990), purification PCR product by PEG precipitation methods (Hiraishi *et al.*, 1995) and ethanol purification methods. Result of purification to analysis against by automated DNA sequencer (ABI PRISM 3130 Genetic Analyzer, Applied Biosystems), thus for trimming and assembling by BioEdit program at BLAST program at the NCBI database, showed that the new isolate was taxonomically very close to *Bacillus cereus* strain HVR22 (Fig. 1).

*Bacillus cereus* produces several Bacteriocin like Inhibitory Substances (BLIS) or cereins, such as cerein 8A produced by *B. cereus* 8A (Bizani and Brandelli, 2002; Bizani *et al.*, 2005), cerein GN105 produced by *B. cereus* GN105 (Naclerio *et al.*, 1993) and cereins produced by *B. cereus* strains 30/11, 15/5, 8/10 and 8/2 (Torkar and Matijas, 2003). Cerein 8A from *B. cereus* 8A isolated from soil of native woodlands of Southern Brazil (Bizani and Brandelli, 2002) is protease sensitive, stable in the pH range of 2-11 and relatively thermostable, only losing activity at temperatures 47°C for 30 min (Bizani *et al.*, 2005). This cerein has an apparent molecular weight of 26 kDa by SDS-PAGE and contains acyl group (s) and aromatic amino acids such as phenylalanine and tyrosine residues which may play a critical role in antimicrobial activity (Sheehan, 2002; Bizani *et al.*, 2005). Cerein 8A inhibits several pathogenic and food-spoilage microorganisms such as *L. monocytogenes* and *B. cereus*, apparently by disturbing their membrane function (Bizani and Brandelli, 2002; Bizani *et al.*, 2005). This BLIS is also active against intact spores of *B. cereus* (Bizani *et al.*, 2005).

**Conclusions:** The isolation of LAB from *budu* was found that 138 isolates exhibited a clear zone, 135 isolates (98%) identified as rod shape and 3 (2%) isolates as cocci shape. For the morphological identification was selected isolated gave 8 (B2, B4, B5, B9, B11, B13, B28 and B34) isolates has been antimicrobial activity against the growth of pathogenic bacteria (*E. coli*, *B. cereus*, *S. thypi*, *B. subtilis* and *L. monocytogenes*). Isolates B28 had the highest antimicrobial activity against all bacteria test, with a range of inhibition zone 14-35 mm, gram positive, rod shape, spore former bacil, motility and catalase positive, Based on morphological examination and PCR analysis, the isolate B28 was primarily identified as *Bacillus cereus* HVR22 strain bacteria.

#### ACKNOWLEDGEMENTS

The author are very grateful to Directorate general of Higher Education that was funded this experiment by Hibah Fundamental Project 2013 Directorate General of Higer Education, Department of National Education Republic of Indonesia.

#### REFERENCES

Ammor, S., G. Tauveron, E. Dufour and I. Chevallier, 2006. Antibacterial activity of lactic acid bacteria against spoilage and pathogenic bacteria isolated from the same meat small-scale facility: 1- Screening and characterization of the antibacterial compounds. *Food Control*, 17: 454-461.

Anihouvi, V.B., G.S. Ayemor, J.D. Hounhouigan and E.S. Dawson, 2006. Quality and characteristic of lanhouin: a traditionally processed fermented fish product in the Republic of Benin. *Ajvan Online. Afr. J. Food Agric. Nutr. Dev.*, pp: 1-15.

Anthony, T., T. Rajesh, N. Kayalvizhi and P. Gunasekaran, 2009. Influence of medium components and fermentation conditions on the production of bacteriocins by *Bacillus licheniformis* AnBa9. *Bioresour. Technol.*, 100: 872-877.

Assefa, E., F. Beyene and A. Santhanam, 2008. Effect of temperature and pH on the antimicrobial activity of inhibitory substances produced by lactic acid bacteria isolated from Ergo, an Ethiopian traditional fermented milk. *Afr. J. Microbiol. Res.*, 2: 229-234.

Bizani, D. and A. Brandelli, 2002. Characterization of a bacteriocin produced by a newly isolated *Bacillus* sp. strain 8A. *J. Applied Microbiol.*, 93: 512-519.

Bizani, D., A.S. Motto, J.A. C. Morissy, R.M.S. Terra, A.A. Souto and A. Brandelli, 2005. Antibacterial activity of cerein 8A, a bacteriocin-like peptida produced by *Bacillus cereus*. *Int. Microbiol. J.*, 8: 125-131.

Cherif, A., S. Chehimi, F. Limem, B.M. Hansen, N.B. Hendriksen, D. Daffonchio and A. Boudabous, 2003. Detection and characterization of the novel bacteriocin entomocin 9 and safety evaluation of its producer, *Bacillus thuringiensis* subsp. entomocidus HD9. *J. Applied Microbiol.*, 95: 990-1000.

Cowan, S.T. and Steel's, 1975. *Manual for the Identification Medical Bacteria*. Cambridge University Press, Cambridge, London.

Davidson, P.M. and S. Zivanovic, 2003. Food antimicrobials. In: Davidson, P.M., J.N. Sofos and A.L. Branen (Eds.), *Antimicrobials in foods.*: CRC press, USA.

De Vuyst, L. and F. Leroy, 2007. Lactic acid bacteria with health claim-interactions and interference with gastro intestinal flora. *Int. Dairy J.*, 9: 25-35.

Fardiaz, 1989. *Penuntun Praktikum Mikrobiologi Pangan*. Fakultas Teknologi Pertanian. Institut Pertanian Bogor, Bogor.

Giraffa, G., 2004. Studying dynamics of microbial populations during food fermentation. *FEMS Microbiol.*, 3: 127-129.

Girum, T., E. Eden and A. Mogessie, 2005. Assessment of the antimicrobial activity of lactic acid bacteria isolated from Borde and Shameta, traditional Ethiopian fermented beverages, on some foodborne pathogens and effect of growth medium on the inhibitory activity. *Int. J. Food Saf.*, 5: 13-20.

Hadioetomo, 1985. *Mikrobiologi Dasar dalam Praktek*. PT. Gramedia. Jakarta.

Hiraishi, A., Y. Kamagata and N. Nakamura, 1995. Polymerase chain reaction amplification and restriction fragment lenght polymorphism analysis of 16S rRNA genes from methanogens. *J. Ferment. Bioengineering*, 79: 523-529.

Holzappel, W.H., R. Geisen and U. Schillinger, 1995. Biological preservation of foods with reference to protective cultures, bacteriocins and food-grade enzymes. *Int. J. Food Microbiol.*, 24: 343-362.

- Hosoi, T. and K. Kiuchi, 2003. Natto-a food made by fermenting cooked soybeans with *Bacillus subtilis* (natto). Handbook of Fermented Functional Foods, Farnworth, E.R. (Ed.), CRC Press, Boca Raton, FL.
- Huda, N. and A. Rosma, 2006. Budu and tukai endemic fermented fish products from West Sumatera. Infofish Int., 3: 49-51.
- Hwanhlem, N., S. Buradaleng, S. Wattanachant and S. Benjakul, 2011. Isolation and screening of lactic acid bacteria from Thai traditional fermented fish (*plasom*) and production of *plasom* from selected strains. J. Food Control, 22: 401-407.
- Ijong, F.G. and Y. Ohta, 1996. Physicochemical and microbiological changes associated with bakasang processing - a traditional Indonesian fermented fish sauce. J. Sci. Food Agri., 71: 69-74.
- Katzung, B.G., 2004. Basic and clinical pharmacology, Lange Medical Books. McGraw-Hill, 9th edition, New York, USA.
- Kim, K.P., C.H. Rhee and H.D. Park, 2002. Degradation of cholesterol by *Bacillus subtilis* SFF34 isolated from Korean traditional fermented flatfish, Letters in Applied Microbiol., 35: 468-472.
- Kopermsub, P. and S. Yunchalard, 2010. Identification of lactic acid bacteria associated with the production of plaa-som, a traditional fermented fish product of Thailand. Int. J. Food Microbiol., 138: 200-204.
- Lay, W.B., 1994. Microbes analysis in laboratory. Raja Grafindo Persada, Jakarta. Indonesia.
- Liasi, S.A., T.I. Azmi, M. Hassan, M. Shuhaimi, M. Rosfarizan and A.B. Ariff, 2009. Antimicrobial activity and antibiotic sensitivity of three isolate of lactic acid bacteria from fermented fish product, *budu*. Malaysian J. Microbiol., 5: 33-37.
- Lodemann, U., B.M. Lorenz, K.D. Weyrauch and H. Martens, 2008. Effects of *Bacillus cereus* var. *toyoi* as probiotic feed supplement on intestinal transport and barrier function in piglets. Arch. Anim. Nutr., 62: 87-106.
- Maragkoudakis, P.A., K.C. Mountzouris, D. Psyrras, S. Cremonese, J. Fischer, M.D. Canter and E. Tsakalidou, 2009. Functional properties of novel protective lactic acid bacteria and application in raw chicken meat against *Listeria monocytogenes* and *Salmonella enteridis*. Int. J. Food Microbiol., 130: 219-226.
- Naclerio, G., E. Risco, M. Sacco and M. De Felice, 1993. Antimicrobial activity of a newly identified bacteriocin of *Bacillus cereus*. Applied Environ. Microbiol., 59: 4313-4316.
- Nicholson, W.L., 2002. Roles of *Bacillus* endospores in the environment. Cell Molecular Life Sci., 59: 410-416.
- Ogunshe, K., M.J. Waites and J.R.M. Hammond, 2007. Nisin and brewing. J. Instrumental Brewing, 94: 233-238.
- Oguntoyinbo, F.A., A.I. Sanni, C.M.A.P. Franz and W.H. Holzapfel, 2007. Phenotypic diversity and technological properties of *Bacillus subtilis* species isolated from *Okpehe*, a traditional fermented condiment. World J. Microbiol. Biotechnol., 23: 401-410.
- Olivera, C.F., G.R. Caron and A. Brandelli, 2004. Bacteriocin-like substance production by *Bacillus licheniformis* strain P40. Letters in Applied Microbiology, 38: 251-256.
- Oscariz, J.C., I. Lasa and A.G. Pisabarro, 1999. Detection and characterization of cerein 7, a new bacteriocin produced by *Bacillus cereus* with a broad spectrum of activity. FEMS Microbiol. Lett., 178: 337-341.
- Paik, H.D., S.S. Bae and J.G. Pan, 1997. Identification and partial characterization of tochicin, a bacteriocin produced by *Bacillus thuringiensis* subsp. *tochigiensis*. J. Indonesia Microbiol. Biotechnol., 19: 294-298.
- Pitcher, D.G., N.A. Saunders and R.J. Owen, 1989. Rapid extraction of bacterial genomic DNA with guanidium thiocyanate. Letters in Applied Microbiol., 8: 151-156.
- Ponce, A.G., M.R. Moreira, C.E. Valle and S.I. Roura, 2008. Preliminary characterization of bacteriocin-like substances from lactic acid bacteria isolated from organic leafy vegetables. LWT-Food Sci. Technol., 41: 432-441.
- Rahayu, E.S. and S. Margino, 1997. Lactic acid bacteria: Isolation and Identification. Workshop Materials. Organized PAU Food and Nutrition, University of Gadjah Mada, Yogyakarta, 13 and June 14.
- Savodogo, A., C.A.T. Quattara, I.H.N. Bassole and S.A. Traore, 2004. Antimicrobial activities of lactic acid bacterial strains isolated from Burkina faso fermented milk. Pak. J. Nutr., 3: 174-179.
- Settanni, L. and A. Corsetti, 2008. Application of bacteriocins in vegetable food biopreservation. Int. J. Food Microbiol., 121: 123-138.
- Sheehan, D., 2002. Physical biochemistry: principles and applications. John Wiley and Sons, Chichester.
- Stein, T., 2005. *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. Moleculer Microbiol., 56: 845-857.
- Tanasupawat, S., N. Sirilak., K. Takuji and I. Takashi, 2009. Identification of halophilic bacteria from fish sauce (*nam-pla*) in Thailand. J. Culture Collections, 6: 69-75.
- Terlabie, N.N., E.S. Dawson and W.K.A. Awua, 2006. The comparative ability of four isolates of *Bacillus subtilis* to ferment soybeans into dawadawa. Int. J. Food Microbiol., 106: 145-152.
- Torkar, K.G. and B.B. Matijasic, 2003. Partial characterisation of bacteriocins produced by *Bacillus cereus* isolates from milk and milk products. Food Technol. Biotechnol., 41: 121-129.



- White, T.J., T.D. Bruns, S.B. Lee and J.W. Taylor, 1990. Amplified and direct sequencing of fungal RNA genes for phylogenetic. In: Innis, M.A., D.H. Gelfand, J.J. Sninsky and T.J. White (Eds.), PCR protocols. Academic, San Diego, pp: 315-322.
- Wu, S., S. Jia, D. Sun, M. Chen, X. Chen, J. Zhong and L. Huan, 2005. Purification and characterization of two novel antimicrobial peptides supeptin JM4-A and subpeptin JM4-B produced by *Bacillus subtilis* JM4. *Curr. Microbiol. Int. J.*, 51: 292-296.