

PJN

ISSN 1680-5194

PAKISTAN JOURNAL OF
NUTRITION

ANSI*net*

308 Lasani Town, Sargodha Road, Faisalabad - Pakistan
Mob: +92 300 3008585, Fax: +92 41 8815544
E-mail: editorpjn@gmail.com



Research Article

Optimization of Glutamate Production from *Lactobacillus plantarum* Originating from Minangkabau Fermented Food as a Feed Supplement for Broiler

¹Vabera Maslami, ¹Yetti Marlida, ¹Mirnawati, ²Jamsari and ¹Yuliaty Shafan Nur

¹Department of Animal Science, Andalas University, West Sumatra, Indonesia

²Department of Crop Science, Faculty of Agriculture, Universitas Andalas, West Sumatra, Indonesia

Abstract

Background and Objective: Glutamate is a non-essential amino acid and it improves the perception of the taste umami and serves as a building block of protein and physiological functions of the body. Increased use of glutamate in animal feed causes glutamate to rise globally. The aim of this study was to obtain the optimal conditions for glutamate production by *Lactobacillus plantarum* VM. **Materials and Methods:** *Lactobacillus plantarum* VM (*L. plantarum* VM) is a lactic acid bacteria originating from Minangkabau fermented foods and produces glutamate. The increased production of glutamate from *Lactobacillus plantarum* VM can be achieved by improving the nutrition and the growth environment of the bacteria. This study was designed in the form of a laboratory experiment protocol and was repeated 3 times. The variables measured in this study were the medium pH, temperature, incubation time, carbon source and nitrogen source. **Results:** The results of this study showed an optimum 5.5 pH (161.519 mg L⁻¹), incubation time (36 h), temperature (36°C) (350.001 mg L⁻¹), 11% glucose (566,535 mg L⁻¹) and 0.5% peptone (680.525 mg L⁻¹). **Conclusion:** Optimization of the initial pH of the media, incubation time, temperature, source C and source N can increase glutamate production.

Key words: Glutamate, umami, building block, *Lactobacillus plantarum*, Minangkabau, fermentation food

Received: March 06, 2018

Accepted: May 14, 2018

Published: June 15, 2018

Citation: Vabera Maslami, Yetti Marlida, Mirnawati, Jamsari and Yuliaty Shafan Nur, 2018. Optimization of glutamate production from *Lactobacillus plantarum* originating from minangkabau fermented food as a feed supplement for broiler. Pak. J. Nutr., 17: 336-343.

Corresponding Author: Yetti Marlida, Department of Animal Science, Andalas University, West Sumatra, Indonesia

Copyright: © 2018 Vabera Maslami *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

The use of amino acids as a supplement in animal feed is practiced extensively. One of the amino acids that is added to animal feed is glutamate. Glutamate is a non-essential amino acid, it enhances the perception of the taste umami and it is a building block for proteins^{1,2}. Apart from its role as a flavor enhancer and a building block for proteins, glutamate acts as a neurotransmitter in the brain and has a number of physiological functions³. Glutamate as a dietary supplement has been widely used in broiler diet. The addition of glutamate in broiler chicken rations can increase body weight and decrease mortality, crude protein ration and fossil ammonia^{4,6}. In addition to improve performance, glutamate can also improve carcass quality. According to Berres *et al.*⁷ and Fujimura *et al.*⁸, adding glutamate can lower abdominal fat, reduce bruising in carcasses and increase the perception of the taste umami in meat.

Increased use of glutamate as a feed supplement for livestock causes an increased glutamate demand globally. According to Sano⁹, the total world glutamate production with fermentation is estimated at 2 million tons/year. Glutamate is an amino acid that has dominated demand and was valued at over 8 billion USD in 2014¹⁰. Therefore, it is necessary to innovate to increase glutamate production. Innovations to increase glutamate production can be done with new organisms, increasing the nutrient content of media and altering the growth environment^{11,12}. According to Niaz *et al.*¹³ and Zareian *et al.*¹⁴, temperature, pH and incubation time can all affect glutamate production. Glutamate is excreted by many bacteria in response to different nutritional conditions and production of glutamate can then be influenced by the concentration of C and N in the culture media. Nadeem *et al.*¹⁵ suggests that the optimization of C and N sources can increase glutamate fermentation using various sources of carbon and nitrogen in the production media. One of the bacteria that can produce glutamate is *Lactobacillus plantarum* VM.

Lactobacillus plantarum VM is a lactic-acid bacteria (LAB) isolated from Minangkabau fermented food. To increase the production of glutamate, it is necessary to optimize the production of *Lactobacillus plantarum* VM. The purpose of this study was to obtain optimal glutamate production by improving the nutrient content of the medium and growth environment of *Lactobacillus plantarum* VM.

MATERIALS AND METHODS

Time and place of study: The study was conducted from 1st August to 1st November, 2017. The study was implemented

at the Industry Feed Technology Laboratory, Faculty of Animal Husbandry, University Andalas.

Experimental design: The research was designed in the form of a laboratory experiment descriptive protocol and repeated 3 times.

Microbes and media content: This study was conducted using *Lactobacillus plantarum* VM bacteria isolated from Minangkabau fermented foods. The *Lactobacillus plantarum* VM sample was a collection of bacteria from previous studies. *Lactobacillus plantarum* VM was cultured on MRS broth media.

Optimizing glutamate production: The glutamate production with minerals media contained (g L^{-1}): 1 g KH_2PO_4 , 0.4 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.01 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ and 0.01 g $\text{MnSO}_4 \cdot 4\text{-}5\text{H}_2\text{O}$ ¹⁶. The first optimization of glutamate production was testing the pH of the media. To decrease the pH, 2 M NaOH was used and it was raised using 2 M HCl to pH values of 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5 and 7. Second, the optimum temperature was determined (30, 33, 36, 39, 42 and 45 °C). Third, the total incubation time was optimized (12, 24, 36, 48, 60, 72, 96 and 108 h). Fourth, the optimal carbon source and dose were determined. Carbon sources used were glucose, sucrose, maltose and lactose. After the best carbon source was determined, the optimum concentration of that carbon source was determined (1, 3, 5, 7, 9, 11, 13 and 15%). Fifth, the optimal sources and concentration of nitrogen were determined (peptone, yeast extract, skim milk, NH_4NO_3 and KNO_3). After obtaining the best nitrogen source, a determination of the optimum dosage of nitrogen source (0.1, 0.3, 0.5, 0.7, 0.9, 1.1, 1.3 and 1.5%) was performed.

Determination of glutamate content: Quantitative measurements of glutamate were performed using HPLC analysis using the method described by Yang *et al.*¹⁷. The mobile phase comprised a mixture of 60% solution A (aqueous solution of 10.254 g of sodium acetate, 0.5 mL tri-ethylamine and 0.7 mL of acetic acid in 1000 mL, with a final pH of 5.8), 12% solution B (acetonitrile) and 28% solution C (aquabides). A gradient HPLC separation was performed on a Shimadzu (Kyoto, Japan) LC 20AT apparatus. A Prevail C18 column (250 × 4.6 mm I.D., particle size 5 $\mu\text{m L}^{-1}$, Alltech, IL, USA) was used during the analysis. The mobile phase from the gradient elution was pumped at a 0.6 ML min^{-1} flow rate at 27 °C and glutamic acid detection was performed at 254 nm.




Fig. 1: Effect of pH medium on glutamate production




Fig. 2: Effect of incubation time on glutamate production

RESULTS AND DISCUSSION

Optimum pH medium: Optimization of the production of glutamate from growing *Lactobacillus plantarum* VM isolates at different pH values (pH 2-7) can be seen in Fig. 1.

The results show that at pH values from 2-7, the isolates could still grow, as is indicated by the turbidity formed in the media and glutamate production. The optimum pH for growth of *Lactobacillus plantarum* VM in this study was acidic (pH 5). According to Zareian *et al.*¹⁴, an acidic pH can trigger the *gdh* gene of LAB to produce higher glutamate. The pH of the growth media of *Lactobacillus plantarum* VM, however is different compared to other studies. As Zareian *et al.*¹⁸ reported the optimum pH for the growth of LAB for glutamate production is pH 4.5. Furthermore, pH 6 has been reported to be the optimum condition of *Lactobacillus plantarum* to produce glutamate¹⁹.

The medium pH is an important factor for *Lactobacillus* growth in biological processes and glutamate production^{20,21}. Decreases in the initial pH of the media may inhibit the growth of *Lactobacillus*, thereby causing a redirection of 2-Oxoglutarate efflux to glutamate production that also increases glutamate excretion^{19,22}. In addition to growth factors, glutamate production increases due to *Lactobacillus plantarum* production of ammonia in an acidic environment, as this contributes to the pH of homeostasis to allow the survival of microorganisms through neutralizing pH²³. Thus, the ammonia produced can be used to form glutamate.

Incubation time: The incubation time for *Lactobacillus plantarum* VM can be seen in Fig. 2. The best incubation time was 36 h, with a production 260,551 mg L⁻¹. The optimal incubation time in this study differs from several other studies. Lawal *et al.*²⁴ reported the optimum incubation time using




Fig. 3: Effect of temperature on glutamate production

Bacillus spp. as 96 h. According to Zariaen *et al.*¹⁸, the best incubation time of *Lactobacillus plantarum* was 96 h for glutamate production at 1.032 mmol. Furthermore, Ahmad *et al.*²⁵ reported 100 h as the best incubation time for producing glutamate.

The optimum time difference of bacterial growth in producing glutamate is influenced by the growth rate of *Lactobacillus* cells. According to Nampoothiri *et al.*²⁶, the growth of *Lactobacillus* cells increases exponentially between 18 and 72 h of fermentation in MRS broth media. This stage is recognized as the log phase or exponential phase. The exponential growth phase has a certain time limit. This is because the nutrients in the media will decrease and the nutrients used for bacterial growth will affect the production of glutamate produced¹⁸. In addition, differences in bacterial strains and media nutrition are another factor responsible for differences in outcomes¹⁴.

Optimum temperature: To determine the optimum temperature of *Lactobacillus plantarum* VM in producing glutamate, incubation at a range of temperatures (30, 33, 36, 39, 42 and 43°C) was performed. The effect of temperature on glutamate production can be seen in Fig. 3.

The results showed that the optimum temperature of *Lactobacillus plantarum* VM in producing glutamate was 36°C (366.76 mg L⁻¹). There was a difference in the production of glutamate with temperature shifts that can be seen in Fig. 2. The optimum temperature obtained in this study is different from that reported by other studies. According to Ahmet *et al.*²⁶, 31°C is the optimum temperature of *Corynebacterium glutamicum* for producing glutamate. Furthermore, Zareian *et al.*¹⁴ and Lawal *et al.*²⁴ stated that the optimum temperature of *Lactobacillus plantarum* in producing glutamate is 37 and 32°C is the optimum temperature for *Bacillus* spp.

The occurrence of differences in glutamate production at each temperature change caused every microbe to have an optimum, maximum and minimum temperature for growth. If the environmental temperature is less than the minimum temperature or greater than the maximum temperature for growth, then enzyme activity can stop and at too high temperature, enzyme denaturation will occur²⁷. According to Lehniger *et al.*²⁸, lower temperatures decrease the fluidity of cell membranes by increasing the fraction of fatty acids containing shorter carbon chains and *cis*-double bonds. This change in lipid composition increases the fluidity of the lipid double layer in the cell membrane and will increase the transport of metabolites and nutrients to maintain the same level as normal growth temperatures. Therefore, as an increase in membrane fluidity, is achieved by increasing the temperature of cultivation, the production of glutamate will increase¹¹. This is the same as is conveyed by Uy *et al.*²⁹ increasing the temperature of cultivation can inhibit the dehydrogenase α -ketoglutarate complex. Thus, it can cause the transfer of 2-Oxoglutarate flux to glutamate production and thus increase the excretion of glutamate²².

Carbon source optimization: Determination of the best carbon source was then performed, comparing different carbon sources to produce glutamate. The carbon sources used in this study were glucose, lactose, sucrose and maltose. Once the highest carbon source to produce glutamate was selected, then a determination of the optimum concentration was performed (1, 3, 5, 7, 9, 11, 13 and 15%). The effect of carbon source and optimum concentration can be seen in Fig. 4-5.

The results showed that the best carbon source for glutamate production was glucose at 351.14 mg L⁻¹. However, this is not much different when compared with sucrose, which produced 150 mg L⁻¹ of glutamate. The difference in




Fig. 4: Effect of carbon sources on glutamate production




Fig. 5: Effect of carbon concentration on glutamate production

glutamate production results from the ability of different microbes to deregulate different carbon sources. According to Nadeem *et al.*¹⁶, each microbe has a different metabolism in the degradation of carbon sources that affects the formation of biomass and the production of primary or secondary metabolites. Research on the production of glutamate from carbon sources has been reported, suggesting that glucose is the best source of carbon for producing glutamate^{14,24}. High glutamate production is obtained because the bacterium has changed glucose through the use of Krebs cycle intermediates³⁰. Glucose can be changed to pyruvic acid through glycolysis, the TCA cycle and the electron transport chain³¹. However, different results were reported by Kiefer *et al.*³², where fructose and sucrose were the best carbon sources for producing glutamate. Thus, the production of glutamate from microbes can be influenced by available carbon sources.

Glucose is the best source of carbon in producing glutamate and author then sought to determine the optimal glucose concentration. The effect of glucose concentration on

glutamate production can be seen in Fig. 7. The results showed that the optimum glucose concentration was 11%, which was higher than in other studies. The best glucose concentrations for glutamate production obtained in this study were lower than that of Zareian *et al.*¹⁸, who reported a concentration of 12% of a carbon source. The high production of glutamate at 11% concentration is due to microbial growth itself. This was also stated by Nampoothiri and Pandey²⁶ that the utilization of sugars in the media by bacteria will continue in accordance with the growth pattern of the bacteria.

Optimization of nitrogen source: Determination of the best nitrogen source was made by comparing the sources of nitrogen (peptone, yeast extract, skim milk, NH₄NO₃ and KNO₃) for glutamate production. After obtaining the best nitrogen source, it was then determined the optimum concentration of nitrogen. The nitrogen source and the optimum concentration of nitrogen can be seen in Fig. 6-7.

The results showed that the best source of nitrogen for producing glutamate was peptone, with a production of




Fig. 6: Effect of nitrogen source on glutamate production




Fig. 7: Effect of nitrogen concentration on glutamate production

415.23 mg L⁻¹. In this study, the best results obtained differed from those reported by Li *et al.*³³ and Nadeem *et al.*¹⁵, where ammonium sulfate was the best source of nitrogen to produce glutamate. According to Savijoki *et al.*³⁴, the determination of suitable nitrogen sources is essential for the needs of amino acids and microbial peptides in producing glutamate. The ability of microbes in synthesizing nitrogen sources has an effect on the production of glutamate expression³⁴. Furthermore, each LAB strain has different needs on each nitrogen source and nitrogen is an essential growth factor or stimulating factor^{35,36}.

After obtaining results indicating peptone as the best source of nitrogen, then the optimum concentration of peptone was determined, which can be seen in Fig. 7. The results showed that the best peptone concentration was 0.9%. Nitrogen plays an important role in the fermentation of glutamate-producing bacteria. Nitrogen is taken up by bacterial cells and then assimilated to achieve metabolism³⁷.

The uptake of nitrogen sources into bacterial cells occurs through passive diffusion (ammonium) or active transport³⁸. The ammonia concentration is low so that as diffusion into the cells becomes limited, the ammonium transporter (AmtB) activates to overcome this nitrogen deficiency and the nitrogen is assimilated with glutamine synthetase³⁹. Conversely, in high nitrogen concentrations, nitrogen diffusion (NH₃) occurs across cytoplasmic membranes. This encourages the growth of bacterial cells and the nitrogen is assimilated by glutamate dehydrogenase to form glutamate. De Angelis *et al.*⁴⁰ also showed that most of the nitrogen is assimilated by glutamate dehydrogenase to form glutamate, which has been shown to exhibit high activity in *Lactobacillus plantarum*.

The implications of this study are that glutamate production from *Lactobacillus plantarum* originating from Minangkabau fermented food can be increased. Optimizing glutamate production by determining the optimum pH,

temperature, incubation time and carbon and nitrogen sources can all increase glutamate production. It is recommended to use more efficient carbon and nitrogen sources to further increase glutamate production and reduce production costs.

CONCLUSION

An increase in glutamate production was obtained using the optimized *Lactobacillus plantarum* VM. Optimization as well as the adjustment of nutrients and environmental conditions affects glutamate fermentation, resulting in the increased production of glutamate. The media pH, incubation time, temperature, C source and N source all impacted glutamate production.

SIGNIFICANCE STATEMENT

This study determined optimal conditions for glutamate production from *Lactobacillus plantarum* originating from Minangkabau fermented food. This study helps researchers to increase the production of glutamate by defining the optimal media, temperature, incubation time, carbon source and nitrogen source, which has not been explored by many other researchers. Thus, a new theory and new lactic-acid bacteria isolated from Minangkabau fermented foods can be used as a feed supplement for poultry, such as broiler chickens, layer or any monogastric animal, leading to increases in the quality immune systems of carcasses.

ACKNOWLEDGMENT

This study was funded by the Ministry of Research, Technology and Higher Education of the Republic of Indonesia through PMDSU No: 1387/E4./2015. We are very grateful to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia and the Rector of Universitas Andalas for their support.

REFERENCES

1. Yamaguchi, S. and K. Ninomiya, 2000. Umami and food palatability. *J. Nutr.*, 130: 921S-926S.
2. Pierre-Andre, G. and M. Yves, 2004. Amino acids: Beyond the building blocks. The Poultry Federation.
3. Meffert, M.K., J.M. Chang, B.J. Wiltgen, M.S. Fanselow and D. Baltimore, 2003. NF- κ B functions in synaptic signaling and behavior. *Nat. Neurosci.*, 6: 1072-1078.
4. Zulkifli, I., M. Shakeri and A.F. Soleimani, 2016. Dietary supplementation of L-glutamine and L-glutamate in broiler chicks subjected to delayed placement. *Poult. Sci.*, 95: 2757-2763.
5. Ebadiasl, G., 2011. Effects of supplemental glutamine and glutamate on growth performance, gastrointestinal development, jejunum morphology and *Clostridium perfringens* count in caecum of broilers. Ph.D. Thesis, Department of Animal Nutrition and Management, Swedish University of Agricultural Science, Sweden.
6. Bezerra, R.M., F.G.P. Costa, P.E.N. Givisiez, E.R. Freitas and C.C. Goulart *et al.*, 2016. Effect of L-glutamic acid supplementation on performance and nitrogen balance of broilers fed low protein diets. *J. Anim. Physiol. Anim. Nutr.*, 100: 590-600.
7. Berres, J., S.L. Vieira, W.A. Dozier III, M.E.M. Cortes, R. de Barros, E.T. Nogueira and M. Kutschenko, 2010. Broiler responses to reduced-protein diets supplemented with valine, isoleucine, glycine and glutamic acid. *J. Applied Poul. Res.*, 19: 68-79.
8. Fujimura, S., F. Sakai and M. Kadowaki, 2001. Effect of restricted feeding before marketing on taste active components of broiler chickens. *J. Anim. Sci.*, 72: 223-229.
9. Sano, C., 2009. History of glutamate production. *Am. J. Clin. Nutr.*, 90: 728S-732S.
10. Radian Insight, 2004. Global amino acid market size is estimated to reach 10.1 million tons by 2022. http://www.abnewswire.com/pressreleases/global-amino-acids-market-to-reach-101-million-tons-by-2022_146363.html
11. Choi, S.U., T. Nihira and T. Yoshida, 2004. Enhanced glutamic acid production of *Brevibacterium* sp. with temperature shift-up cultivation. *J. Biosci. Bioeng.*, 98: 211-213.
12. Shirai, T., A. Nakato, N. Izutani, K. Nagahisa and S. Shioya *et al.*, 2005. Comparative study of flux redistribution of metabolic pathway in glutamate production by two coryneform bacteria. *Metab. Eng.*, 7: 59-69.
13. Niaz, B., S. Nadeem, H.M. Muzammil, J.A. Khan and T. Zahoor, 2009. Optimization of fermentation conditions for enhanced glutamic acid production by a strain of *Corynebacterium glutamicum* NIAB BNS-14. *Pak. J. Zool.*, 41: 261-267.
14. Zareian, M., A. Ebrahimpour, A.K.S. Mohamed and N. Saari, 2013. Modeling of glutamic acid production by *Lactobacillus plantarum* MNZ. *Electron. J. Biotechnol.*, 16: 1-16.
15. Nadeem, S., B. Niaz, H.M. Muzammil, S.M. Rana, M.I. Rajoka and A.R. Shakoori, 2011. Optimising carbon and nitrogen sources for L-glutamic acid production by *Brevibacterium* strain NIAB SS-67. *Pak. J. Zool.*, 43: 285-290.
16. Nakamura, J., S. Hirano, H. Ito and M. Wachi, 2007. Mutations of the *Corynebacterium glutamicum* NCgl1221 gene, encoding a mechanosensitive channel homolog, induce L-glutamic acid production. *Applied Environ. Microbiol.*, 73: 4491-4498.

17. Yang, S.Y., F.X. Lu, Z.X. Lu, X.M. Bie, Y. Jiao, L.J. Sun and B. Yu, 2008. Production of γ -aminobutyric acid by *Streptococcus salivarius* subsp. *thermophilus* Y2 under submerged fermentation. *Amino Acids*, 34: 473-478.
18. Zareian, M., A. Ebrahimpour, F.A. Bakar, A.K.S. Mohamed, B. Forghani, M.S.B. Ab-Kadir and N. Saari, 2012. A glutamic acid-producing lactic acid bacteria isolated from Malaysian fermented foods. *Int. J. Mol. Sci.*, 13: 5482-5497.
19. Zacharof, M.P. and R.W. Lovitt, 2010. Development of an optimised growth strategy for intensive propagation, lactic acid and bacteriocin production of selected strains of *Lactobacilli* genus. *Int. J. Chem. Eng. Applic.*, 1: 55-63.
20. Yang, E., L. Fan, J. Yan, Y. Jiang, C. Doucette, S. Fillmore and B. Walker, 2018. Influence of culture media, pH and temperature on growth and bacteriocin production of bacteriocinogenic lactic acid bacteria. *AMB Express*, Vol. 8, No. 1. 10.1186/s13568-018-0536-0.
21. Eggeling, L. and M. Bott, 2005. *Handbook of Corynebacterium glutamicum*. CRC Press, Boca Raton, FL., USA.
22. Asakura, Y., E. Kimura, Y. Usuda, Y. Kawahara, K. Matsui, T. Osumi and T. Nakamatsu, 2007. Altered metabolic flux due to deletion of *odhA* causes L-glutamate overproduction in *Corynebacterium glutamicum*. *Applied Environ. Microbiol.*, 73: 1308-1319.
23. Jaichumjai, P., R. Valyasevi, A. Assavanig and P. Kurdi, 2010. Isolation and characterization of acid-sensitive *Lactobacillus plantarum* with application as starter culture for Nham production. *Food Microbiol.*, 27: 741-748.
24. Lawal, A.K., B.A. Oso, A.I. Sanni and O.O. Olatunji, 2011. L-Glutamic acid production by *Bacillus* spp. isolated from vegetable proteins. *Afr. J. Biotechnol.*, 10: 5337-5345.
25. Ahmed, Y.M., J.A. Khan, K.A. Abulnaja and A.L. Al-Malki, 2013. Production of glutamic acid by *Corynebacterium glutamicum* using dates syrup as carbon source. *Afr. J. Microbiol.*, 7: 2071-2077.
26. Nampoothiri, K.M. and A. Pandey, 1996. Urease activity in a glutamate producing *Brevibacterium* sp. *Process Biochem.*, 31: 471-475.
27. Sanchez-Peinado, M.D.M., J. Gonzalez-Lopez, B. Rodelas, V. Galera, C. Pozo and M.V. Martinez-Toledo, 2008. Effect of linear alkylbenzene sulfonates on the growth of aerobic heterotrophic cultivable bacteria isolated from an agricultural soil. *Ecotoxicology*, 17: 549-557.
28. Lehniger, A.L., D.L. Nelson and M.M. Cox, 1993. *Principles of Biochemistry*. 2nd Edn., Worth, New York, pp: 268-274.
29. Uy, D., S. Delaunay, P. Germain, J.M. Engasser and J.L. Goergen, 2003. Instability of glutamate production by *Corynebacterium glutamicum* 2262 in continuous culture using the temperature-triggered process. *J. Biotechnol.*, 104: 173-184.
30. Rizal, Y. and G. Wu, 2012. *Metabolisme Protein dan Asam-Asam Amino*. Andalas University Press, Padang.
31. Williams, A.G., S.E. Withers, E.Y. Brechany and J.M. Banks, 2006. Glutamate dehydrogenase activity in *Lactobacilli* and the use of glutamate dehydrogenase-producing adjunct *Lactobacillus* spp. cultures in the manufacture of cheddar cheese. *J. Applied Microbiol.*, 101: 1062-1075.
32. Kiefer, P., E. Heinzle and C. Wittmann, 2002. Influence of glucose, fructose and sucrose as carbon sources on kinetics and stoichiometry of lysine production by *Corynebacterium glutamicum*. *J. Ind. Microbiol. Biotechnol.*, 28: 338-343.
33. Li, P., Y.L. Yin, D. Li, S.W. Kim and G. Wu, 2007. Amino acids and immune function. *Br. J. Nutr.*, 98: 237-252.
34. Savijoki, K., H. Ingmer and P. Varmanen, 2006. Proteolytic systems of lactic acid bacteria. *Applied Microbiol. Biotechnol.*, 71: 394-406.
35. Letort, C. and V. Juillard, 2001. Development of a minimal chemically-defined medium for the exponential growth of *Streptococcus thermophilus*. *J. Applied Microbiol.*, 91: 1023-1029.
36. Barrangou, R., S.J. Lahtinen, F. Ibrahim and A.C. Ouwehand, 2011. Genus *Lactobacilli*. In: *Lactic Acid Bacteria: Microbiological and Functional Aspects*, Lahtinne, S., S. Salminen, A. von Wright and A. Ouwehand (Eds.), CRC Press, London, pp: 77-92.
37. Burkovski, A., 2003. Ammonium assimilation and nitrogen control in *Corynebacterium glutamicum* and its relatives: An example for new regulatory mechanisms in actinomycetes. *FEMS Microbiol. Rev.*, 27: 617-628.
38. Meier-Wagner, J., L. Nolden, M. Jakoby, R. Siewe, R. Kramer and A. Burkovski, 2001. Multiplicity of ammonium uptake systems in *Corynebacterium glutamicum*: role of Amt and AmtB. *Microbiology*, 147: 135-143.
39. Jakoby, M., L. Nolden, J. Meier Wagner, R. Kramer and A. Burkovski, 2000. AmtR, a global repressor in the nitrogen regulation system of *Corynebacterium glutamicum*. *Mol. Microbiol.*, 37: 964-977.
40. De Angelis, M., M. Calasso, R. di Cagno, S. Siragusa, F. Minervini and M. Gobbetti, 2010. NADP glutamate dehydrogenase activity in nonstarter lactic acid bacteria: Effects of temperature, pH and NaCl on enzyme activity and expression. *J. Applied Microbiol.*, 109: 1763-1774.