

Asian Journal of
Applied
Sciences

Review Article

Phytochemical and Pharmacological Aspects of Anthocyanins

Md. Sohail Akhtar and Wegdan Ali Shehata

College of Pharmacy and Nursing, University of Nizwa, P.O. Box 33, Post Code 616, Birkat Al Mouz, Nizwa, Sultanate of Oman, Oman

Abstract

Anthocyanin is a natural compound in plants which comes under the phenolic compounds class. It is responsible for imparting attractive colors in the plants like red, purple and blue. Since, ancient times, plants containing anthocyanins have been used in herbal remedies to treat different medical conditions. The aim of this review article was to provide comprehensive information about the chemistry, stability, pH influences and different pharmacological activities exhibited by anthocyanins and emphasis its importance. Anthocyanins are known to possess antioxidant, anti-inflammatory, anti-cancer, anti-diabetic, cardiovascular and neuroprotective activity. It was also reported that anthocyanins contribute in enhancing learning and memory. In industrial field, anthocyanins are used as natural food colorants and this is for their relative safety when compared to the synthetic food colorants.

Key words: Anthocyanin, antioxidant, pigment, pharmacological activity, natural colorants, phenolic compounds, phytochemical

Citation: Md. Sohail Akhtar and Wegdan Ali Shehata, 2020. Phytochemical and pharmacological aspects of anthocyanins. Asian J. Applied Sci., 13: 94-99.

Corresponding Author: Md. Sohail Akhtar, College of Pharmacy and Nursing, University of Nizwa, P.O. Box 33, Post Code 616, Birkat Al Mouz, Nizwa, Sultanate of Oman, Oman Tel: +968-25-446-423

Copyright: © 2020 Md. Sohail Akhtar and Wegdan Ali Shehata. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Phenolic compounds are vital constituents that are abundant in a wide variety of fruits and vegetables and they have a role in their sensory properties like bitterness and astringency¹. As described in Holy Quran, "The fruits like olive, grape, date, fig and pomegranate are gifts and heavenly fruits of God", these fruits contain large amounts of phenolic compounds like anthocyanins, flavonoids, coumarins and tannins². Anthocyanins are classified as phenolic compounds and they are water-soluble pigments³.

The origin of the word anthocyanin comes from a Greek word that was used to describe a pigment which is blue, that cornflower (*Centaurea cyanus*) contained⁴. They are responsible for most of the bright and shiny colors as blue, pink, purple, red and orange that can be seen in most of the fruits, leaves, flowers and cereal grains⁵.

The type and quantity of anthocyanins range extensively among diverse plants species. This result in a variation of anthocyanin intake levels widely by season, region and individuals factors like culture and education⁶.

ANTHOCYANINS CHEMISTRY

Anthocyanins are considered as flavonoids, but due to their ability to form flavylium cation (Fig. 1) they can be easily identified or distinguished from the rest of flavonoids compounds⁷. They can be classified into two types, glycosides and acyl glycosides based on their aglycone anthocyanidins (Fig. 2)⁸. In a fresh plant, it is rare to find aglycones. Seventeen anthocyanidins can be found in nature, but only cyanidin, petunidin, pelargonidin, delphinidin, malvidin and peonidin are widely distributed and available (Fig. 3)⁹. Anthocyanins vary in the position and number of methoxyl and hydroxyl groups on the primary anthocyanidin structure¹⁰. This means that the anthocyanin component is identified




Fig. 1: Flavylium cations structure

Source: Miguel⁷

Anthocyanidin	R1	R2
Pelargonidin	H	H
Cyanidin	OH	H
Peonidin	OCH ₃	H
Delphinidin	OH	OH
Malvidin	OCH ₃	OCH ₃
Petunidin	OCH ₃	OH
R ₃	OH, glycosyl	
R ₄	OH, glycosyl	

Fig. 2: Structure of anthocyanidin

Source: Ghosh⁸




Fig. 3: Structure of naturally occurring anthocyanidins

Source: Pervaiz *et al.*⁹

based on the number and position of the sugar linked. In addition, it is determined by the acylating agent identity and sugar acylation identity.

Anthocyanin occurs mainly as glycosylated anthocyanidin which means that one or more sugar or acylated sugar is attached to aglycone. There are four common substituted mono-sides: Arabinose, rhamnose, glucose and galactose¹¹. The glycosylation process provides stability to anthocyanins and anthocyanidins are found to be less stable. Basically, anthocyanins structure has 2 parts: Chromophore (a light-absorbent) which will be called anthocyanidin with a glycoside attachment. Cyanidin-3-glucoside is one of the common anthocyanin structure found in plants¹². The chromophore which is basic in anthocyanins is a 7-hydroxy flavylum ion. Anthocyanins that occur naturally mainly have hydroxyl substituents at positions 3 (Permanently glycosylated, adding thermal stability) and 5 (occasionally glycosylated) and one or additional methoxy or hydroxy substituents is contained in the 2-phenyl or B-ring¹³.

STABILITY OF ANTHOCYANINS

Anthocyanins stability is influenced by many factors like changes in pH, oxygen existence, the chemical composition of the whole structure, applying UV-radiation chemical interactions with consumed food causing metal ions chelation and changes in temperature and light¹⁴. The influence of these factors results in changes and degradation of the colors. However, some studies showed that if other synergistic antioxidant constituents present, it may contribute to the inhibition or decreasing of anthocyanin degradation¹⁵.

In a study conducted to evaluate the temperature influence on the degradation of anthocyanins, it was found that temperature has a significant effect on the anthocyanin pigment stability whether light is present or absent¹⁶. In a study performed on anthocyanins present in sorghum plant, it was found that the methanol which was acidified first and used for the extraction had a higher stability than the methanol extract without the acidification process and the discoloration by 2 ways of extraction was directly proportional with the increase in temperature, time and light¹⁷.

The change of color of plants containing anthocyanins sometimes occurs due to enzymatic degradation or browning. This occurs due to the oxidation of phenolic compounds as they convert to o-quinones enzymatically. There are 2 enzymes responsible for this process, polyphenol oxidase (PPO) and peroxidase (POD), because of the structure of

anthocyanin which contains a sugar moiety, the PPO enzyme has low affinity against anthocyanin as a result of steric hindrance. If phenols are present the activity of PPO against anthocyanins increases due to increased oxygen consumptions. The PPO can be found in plum, strawberry, grape and blueberry^{18,19}.

pH INFLUENCE

Basically, anthocyanins color intensity is greater in acidic condition. The change in anthocyanins color is more in alkaline conditions due to instability and degradation. The explanation for this behavior is possibly due to the interaction of the red flavylum cation with co-pigments present which will result in changing the adsorption properties of the solution containing anthocyanins and this at the end will create a more stable anthocyanin extract and protect the color features of the particular solution²⁰.

At different pH conditions, the structure of anthocyanin differs. In a solution of pH = 1, the red color predominates as flavylum cation. When the pH increases, a competition which is thermodynamic and kinetic starts between the proton transfer reaction associated with the acidic hydroxyl groups of the aglycone and the hydration reaction related to the flavylum cation. In a range of pH = 2-4, blue color predominate as a quinoidal species. At pH values range from 5-6, carbinol pseudo base and chalcone species which are colorless present. When the pH values are greater than 7 degradation of anthocyanins occurs which depends on the substituent groups. Depending on the pH in a range of 4-6 anthocyanins can be identified in different 4 forms which are flavylum cation, carbinol without any color, quinoidal base which is anhydrous and a chalcone with a light yellow color²¹.

PHARMACOLOGICAL ACTIVITIES REPORTED OF ANTHOCYANINS

Anti-oxidant activity: It was observed in studies that anthocyanins contribute to protecting cells by preventing cells and tissue damage caused by oxidant stress. Consumption of foods rich in anthocyanin could contribute to modulating specific secondary physiological changes of the oxidant stress and thus preventing obesity²². Phenolic compounds like anthocyanins extensively distributed in plants which have been reported to exert different pharmacological effect, including antioxidant activity²³. The 1,1-diphenyl-2-picrylhydrazyl (DPPH) mostly used for the estimation of free radical-scavenging activity of plant antioxidants²⁴.

Anti-inflammatory activity: In a study that was conducted to analyze the anti-inflammatory properties identified in anthocyanins, it was proved that cyanidin-3-glucoside metabolites were effective under physiological conditions as anti-inflammatory agents for the endothelial cells in the vascular system of humans²⁵.

Anti-cancer activity: Secondary metabolites are major sources of new chemical compounds for development of new medicines including anticancer agents²⁶.

The fast growth in the cancer problem denotes a crisis for healthcare systems around world and there is need for search of better drug control cancer²⁷.

Due to anti-oxidant, anti-mutagenesis and anti-inflammation properties it was found that anthocyanins have the ability to inhibit propagation of cells by modifying the pathways of signal transduction, which will result in stimulating cell cycle arrest and inducing apoptosis or in other term autophagy of tumor cells. It was observed that anthocyanins have the ability to increase the tumor cells sensitivity to chemotherapy drugs and thereby reversing and reducing drug resistance²⁸.

Cardiovascular effect: It was found that the consumption of anthocyanins contributes to the treatment of chronic diseases such as; Cardiovascular Diseases (CVD). This is due to the chemical structure of anthocyanin, preventing endothelial dysfunction and inflammatory processes²⁹.

Antidiabetic activity: Many studies supported that anthocyanins prevent the occurrence of diabetes mellitus³⁰. It was found that anthocyanins by the up regulation of insulin-regulated glucose transporter (GLUT4), it controlled the carbohydrate metabolism and that peroxisome proliferator-activated receptor- γ (PPAR γ) found in skeletal muscles and adipose tissue was activated by anthocyanins³¹. Anthocyanins works on α -glucosidase along with pancreatic α -amylase to metabolize, digest carbohydrates and thus, decrease the glucose level in the blood and this mechanism mimics to some extent the way acarbose works on our body. Along with that, it was also found that anthocyanins improve insulin resistance, raise insulin sensitivity and uptake of glucose³².

Neuroprotective activity: Anthocyanins properties such as; anti-neuroinflammatory, anti-apoptotic and antioxidant have a therapeutic effect in the treatment of neurodegenerative diseases such as; Parkinson's disease³³.

Learning and memory enhancing effects: In a study conducted to evaluate the effects of anthocyanins in learning and memory, it was found that it decreased the level of lipid peroxidation damage in the brain which will slow down memory damage that result from oxidative stress. Overall, it was found that anthocyanins provide an improvement in learning and memory³⁴.

MEDICINAL USES

From many years ago, anthocyanin was contained in the human diet and was used traditionally in form of herbals owing to their physiological ability to treat many conditions such as; liver disorders, hypertension, diarrhea and dysentery, pyrexia, common cold and urinary problems³⁵.

For instance, *Hibiscus* sp. which contains anthocyanins have been used in folk medicines in the past to treat hypertension and liver dysfunction. Anthocyanins found in bilberry (*Vaccinium*) were historically used as a traditional remedy for diarrhea, microbial infections and vision disorders³⁶.

INDUSTRIAL USES

To improve the appearance of foods, industries use synthetic colorants. However, synthetic colorants caused considerable side effects and natural colorants extracted from anthocyanins proved they have more benefits with almost absence of adverse effects^{37,38}. Anthocyanins provide a wide range of intensive pigments such as; orange, purple, blue and red colors. Examples of plants reach with anthocyanins pigments are red grapes, blueberries, purple corn, strawberry, elderberries, black chokeberries, black currants, cranberry and radish potato³⁹. Extraction of the natural pigments is usually done by homogenization and then solvent extraction. In industries ethanol is preferred for the extraction because methanol is associated with toxicity⁴⁰.

CONCLUSION

Anthocyanins in many food and drinks appear to have an important role in the prevention of diverse diseases. In the last few years, synthetic food dyes have been banned in many countries because of their poisoning and carcinogenicity. Anthocyanin are coloured natural compounds that are easily obtained from fruits and vegetables can be considered as a possible alternative to banned food dyes. In fact, have shining attractive colors, while its high solubility allows for easy

integration into aquatic diets. The nutrient-rich extracts are increasingly attractive to the food industry as natural substitutes for the synthetic dye.

SIGNIFICANCE STATEMENT

This review discovers the phytochemicals and pharmacological activities of anthocyanins that can be beneficial for healthcare management. This study will help the researcher to uncover the critical areas of pharmacological activities of anthocyanins that many researchers were not able to explore.

ACKNOWLEDGMENT

The authors are grateful to University of Nizwa for providing necessary facility for writing this review.

REFERENCES

1. Mozetic, B., P. Trebse and J. Hribar, 2002. Determination and quantitation of anthocyanins and hydroxycinnamic acids in different cultivars of Sweet Cherries (*Prunus avium* L.) from Nova Gorica region (Slovenia). Food Technol. Biotechnol., 40: 207-212.
2. Akhtar, M.S., S.D. Al-Balushi and M.M.S. Awladthani, 2018. Phytochemical investigation, cytotoxicity, antibacterial and anti-oxidant activity of *Olea europea* fruits grown in Oman. Eur. J. Biomed. Pharmaceut. Sci., 5: 98-102.
3. Jakobek, L., M. Seruga, M. Medvidovic-Kosanovic and I. Novak, 2007. Anthocyanin content and antioxidant activity of various red fruit juices. Deutsche Lebensmittel-Rundschau, 103: 58-64.
4. Ponmozhi, P., M. Geetha, S.M. Kumar and S.P. Devi, 2011. Extraction of anthocyanin and analysing its antioxidant properties from *Pithecellobium dulce* fruit pericarp. Asian J. Pharmaceut. Clin. Res., 4: 41-45.
5. Sun, L.L., W. Gao, M.M. Zhang, C. Li, A.G. Wang, Y.L. Su and T.F. Ji, 2014. Composition and antioxidant activity of the anthocyanins of the fruit of *Berberis heteropoda* Schrenk. Molecules, 19: 19078-19096.
6. Fang, J., 2015. Classification of fruits based on anthocyanin types and relevance to their health effects. Nutrition, 31: 1301-1306.
7. Miguel, M.G., 2011. Anthocyanins: Antioxidant and/or anti-inflammatory activities. J. Applied Pharm. Sci., 1: 7-15.
8. Ghosh, D., 2005. Anthocyanins and anthocyanin-rich extracts in biology and medicine: Biochemical, cellular and medicinal properties. Curr. Top. Nutraceut. Res., 3: 113-124.
9. Pervaiz, T., J. Songtao, F. Faghihi, M.S. Haider and J. Fang, 2017. Naturally occurring anthocyanin, structure, functions and biosynthetic pathway in fruit plants. J. Plant Biochem. Physiol., Vol. 5, No. 2. 10.4172/2329-9029.1000187.
10. Wu, X., G.R. Beecher, J.M. Holden, D.B. Haytowitz, S.E. Gebhardt and R.L. Prior, 2006. Concentrations of anthocyanins in common foods in the United States and estimation of normal consumption. J. Agric. Food Chem., 54: 4069-4075.
11. Chaiyasut, C., B.S. Sivamaruthi, N. Pengkumsri, S. Sirilun, S. Peerajan, K. Chaiyasut and P. Kesika, 2016. Anthocyanin profile and its antioxidant activity of widely used fruits, vegetables and flowers in thailand. Asian J. Pharmaceut. Clin. Res., 9: 218-224.
12. Arnnok, P., C. Ruangviriyachai, R. Mahachai, S. Techawongstien and S. Chanthai, 2012. Determination of total phenolics and anthocyanin contents in the pericarp of hot chilli pepper (*Capsicum annum* L.). Int. Food Res. J., 19: 235-243.
13. Quina, F.H., P.F. Moreira Jr., C. Vautier-Giongo, D. Rettori and R.F. Rodrigues *et al.*, 2009. Photochemistry of anthocyanins and their biological role in plant tissues. Pure Applied Chem., 81: 1687-1694.
14. Lima, A.D.J.B., A.D. Correa, A.A. Saczk, M.P. Martins and R.O. Castilho, 2011. Anthocyanins, pigment stability and antioxidant activity in jabuticaba [*Myrciaria cauliflora* (Mart.) O. Berg]. Rev. Bras. Fruticult., 33: 877-887.
15. Oancea, S. and O. Draghici, 2013. PH and thermal stability of anthocyanin-based optimised extracts of romanian red onion cultivars. Czech J. Food Sci., 31: 283-291.
16. Rakkimuthu, R., S. Palmurugan and A. Shanmugapriya, 2016. Effect of temperature, light, pH on the stability of anthocyanin pigments in *Cocculus hirsutus* fruits. Int. J. Multidisciplin. Res. Mod. Educ., 2: 91-96.
17. Devi, P.S., M. Saravanakumar and S. Moh, 2012. The effects of temperature and pH on stability of anthocyanins from red sorghum (*Sorghum bicolor*) bran. Afr. J. Food Sci., 6: 567-573.
18. Ducamp-Collin, M.N., M. Lebrun, H. Ramarson and G. Self, 2007. Anthocyanins and anthocyanin-degrading enzymes in Kwai May and Wai Chee cultivars of litchis grown in Reunion Island and Spain. Fruits, 62: 353-359.
19. Jiang, Y., 2000. Role of anthocyanins, polyphenol oxidase and phenols in lychee pericarp browning. J. Sci. Food Agric., 80: 305-310.
20. Sipahli, S., V. Mohanlall and J.J. Mellem, 2017. Stability and degradation kinetics of crude anthocyanin extracts from *H. sabdariffa*. Food Sci. Technol., 37: 209-215.
21. Castaneda-Ovando, A., M. de Lourdes Pacheco-Hernandez, M.E. Paez-Hernandez, J.A. Rodriguez and C.A. Galan-Vidal, 2009. Chemical studies of anthocyanins: A review. Food Chem., 113: 859-871.
22. Wu, H.Y., K.M. Yang and P.Y. Chiang, 2018. Roselle anthocyanins: Antioxidant properties and stability to heat and pH. Molecules, Vol. 23, No. 6. 10.3390/molecules 23061357.

23. Akhtar, M., S. Padmini, S. Prabodh, S. Alok and M. Alok, 2010. Comparative *in-vitro* evaluation of *Vitex leucoxylo* Linn. bark for antioxidant activity. Int. J. Pharm. Sci. Res., 1: 27-33.
24. Akhtar, M.S., M.A. Hossain and S.A. Said, 2017. Isolation and characterization of antimicrobial compound from the stem-bark of the traditionally used medicinal plant *Adenium obesum*. J. Trad. Complement. Med., 7: 296-300.
25. Blando, F., N. Calabriso, H. Berland, G. Maiorano, C. Gerardi, M.A. Carluccio and O.M. Andersen, 2018. Radical scavenging and anti-inflammatory activities of representative anthocyanin groupings from pigment-rich fruits and vegetables. Int. J. Mol. Sci., Vol. 19, No. 1. 10.3390/ijms19010169.
26. Wel, A.M., R.S. Al-Burtamani, M.S. Al-Azzawi, M.S. Akhtar and S.A. Said, 2019. Antimicrobial, antioxidant and cytotoxic activity of *Maytenus dhofarensis* growing in Oman. J. Applied Sci., 19: 229-234.
27. Said, S.A., Y. Tamimi, M.S. Akhtar, A.M. Wel, S.S. Al-Khanjari and Q.A. Al-Riyami, 2017. *In vitro* anticancer activity of selected medicinal plants from Oman. J. Pharmaceut. Res. Int., 15: 1-8.
28. Lin, B.W., C.C. Gong, H.F. Song and Y.Y. Cui, 2017. Effects of anthocyanins on the prevention and treatment of cancer. Br. J. Pharmacol., 174: 1226-1243.
29. Reis, J.F., V.V.S. Monteiro, R. de Souza Gomes, M.M. do Carmo, G.V. da Costa, P.C. Ribera and M.C. Monteiro, 2016. Action mechanism and cardiovascular effect of anthocyanins: A systematic review of animal and human studies. J. Transl. Med., Vol. 14, No. 1. 10.1186/s12967-016-1076-5.
30. Sivamaruthi, B.S., P. Kesika, K. Subasankari and C. Chaiyasut, 2018. Beneficial effects of anthocyanins against diabetes mellitus associated consequences-a mini review. Asian Pac. J. Trop. Biomed., 8: 471-477.
31. Rozanska, D. and B. Regulska-Ilow, 2018. The significance of anthocyanins in the prevention and treatment of type 2 diabetes. Adv. Clin. Exp. Med., 27: 135-142.
32. Belwal, T., S.F. Nabavi, S.M. Nabavi and S. Habtemariam, 2017. Dietary anthocyanins and insulin resistance: When food becomes a medicine. Nutrients, Vol. 9, No. 10. 10.3390/nu9101111.
33. Winter, A.N., M.C. Brenner, N. Punessen, M. Snodgrass, C. Byars, Y. Arora and D.A. Linseman, 2017. Comparison of the neuroprotective and anti-inflammatory effects of the anthocyanin metabolites, protocatechuic acid and 4-hydroxybenzoic acid. Oxid. Med. Cell. Longevity, Vol. 2017. 10.1155/2017/6297080.
34. Kangwan, N., K. Pintha, W. Preedapirom, P. Tantipaiboonwong, O. Chumphukam and M. Suttajit, 2015. Learning and memory enhancing effects of anthocyanin in black rice extract on cerebral ischaemia in mice. ScienceAsia, 41: 315-321.
35. Shipp, J. and E.S.M. Abdel-Aal, 2010. Food applications and physiological effects of anthocyanins as functional food ingredients. Open Food Sci. J., 4: 7-22.
36. Lila, M.A., 2004. Anthocyanins and human health: An *in vitro* investigative approach. J. Biomed. Biotechnol., 5: 306-313.
37. Jackman, R.L., R.Y. Yada, M.A. Tung and R.A. Speers, 1987. Anthocyanins as food colorants—a review. J. Food Biochem., 11: 201-247.
38. He, X.L., X.L. Li, Y.P. Lv and Q. He, 2015. Composition and color stability of anthocyanin-based extract from purple sweet potato. Food Sci. Technol., 35: 468-473.
39. Shaik, A., K.N. Killari and J. Panda, 2018. A review on anthocyanins: A promising role on phytochemistry and pharmacology. Int. Res. J. Pharm., 9: 1-9.
40. Hua, Z., D. Yuesheng, X. Ge, L. Menglu, D. Liya, A. LiJia and X. Zhilong, 2013. Extraction and purification of anthocyanins from the fruit residues of *Vaccinium uliginosum* Linn. J. Chromatogr. Sep. Tech., Vol. 4, No. 2. 10.4172/2157-7064.1000167.