

ISSN 1819-1894

Asian Journal of
Agricultural
Research

Egg Quality Defects in Poultry Management and Food Safety

O.K. Chukwuka, I.C. Okoli, N.J. Okeudo, A.B.I. Udedibie, I.P. Ogbuwu, N.O. Aladi,
O.O.M. Iheshiolor and A.A. Omede

Department of Animal Science and Technology, Federal University of Technology, P.M.B 1526, Imo State,
Nigeria

*Corresponding Author: O.K. Chukwuka, Department of Animal Science and Technology, Federal University of
Technology, P.M.B 1526, Imo State, Nigeria Tel: +2347038494388*

ABSTRACT

Poultry egg is a vehicle for reproduction; it also serves as a source of food for human consumption. The size and shape of avian eggs differs among the various species of birds, but all eggs have three main parts-yolk, albumen and shell. The quality of eggs depends on physical make up and chemical composition of its constituent parts. Due to the diversity in the potential uses of poultry eggs and the subsequent consumer demands that egg quality become extremely difficult to define. Egg quality is the more important price contributing factor in table and hatching eggs. It is obvious that quality of egg is important from producer's point of view. One of the biggest challenges for the poultry industry is to provide consistent quality egg products to the consumer. Thus breeding companies are shifting selection emphasis to improved egg quality. Problems associated with egg quality include: egg shell defect and internal defects which can be broadly categorized into three groups namely: defects affecting yolk quality, defects affecting albumin quality and defects affecting overall quality. Egg quality defects are usually easily resolved, but can be costly if they are not dealt with quickly.

Key words: Yolk quality, albumen quality, shell quality, egg size, colour, management

INTRODUCTION

Kramer (1951) defined food quality as the sum of characteristics of a given food item which influence the acceptability or preference for that food by the consumer. Based on this definition, it is clear that egg quality will mean different things to different people and the consumer's perception of quality is likely to vary depending on their intended use of the egg and their own preferences. This is clearly illustrated by a brief review of the regulatory requirements for eggs sold around the world.

Under the European Union Egg Marketing Regulations, enforced in all European Union Countries, eggs are classed as either class A or B eggs (Council of the European Union, 2006; European Commission, 2003) and only eggs graded class A can be sold for direct human consumption or retailed (Council of the European Union, 2006). The characteristics of grade A and B eggs are detailed in Table 1. Similarly, the United States Department of Agriculture (USDA) has developed three grades of eggs based on the interior quality of the egg, appearance and condition of the egg shell (USDA Food Safety and Inspection Service, undated). US grade AA and US grade A eggs are usually retailed while U.S. grade B eggs are usually sent for further processing. In contrast, no defined grades have been developed for shell eggs sold in Nigeria and little attention has been focused on internal egg quality factors, such as yolk or albumin quality.

Table 1: Summary of standards for exterior quality of chicken eggs

Factor	Grade AA or A	Grade B	Dirty
Stain	- Must be clean - May show small specks, stains or cage marks that do not detract from general appearance of the egg - May show traces of processing oil	- Slight stains - Moderate stains: localized (single) <1/32 of shell - Scattered (2 or more) <1/16 of shell	- Prominent stains - Moderate stains: - Localized (single) >1/32 of shell - Scattered >1/16 of shell
Adhering dirt or foreign material	None	None	Adhering dirt or foreign material
Egg shape	Approximately the usual elliptical shape	Unusual or decidedly misshapen (very long or distorted)	
Shell texture	May have rough areas and small calcium deposits that do not materially affect shape or strength	- Extremely rough areas that may be faulty in soundness or strength - May have large calcium deposits	
Ridges	May have slight ridges that do not materially affect shape or strength	May have pronounced ridges	
Shell thickness	Must be free from thin spots	May show pronounced thin spots	

Source: Jacob *et al.* (2000)

However, the Animal Products Specification for Products Intended for Human Consumption Notice 2000 requires that shell eggs intended to be traded in the shell must:

- Be visibly clean
- Have no cracks that are visible on candling (or equivalent) unless they have been treated by a process that destroys pathogenic organisms
- Have no evidence of embryo development or putrefaction, and no significant blood clots
- Not have been incubated
- Be handled and stored under conditions that minimise condensation on the surface of the eggs

Eggs which do not meet these minimum requirements can only be sold for human consumption if they have been pasteurized (or undergone an equivalent process) and meet the microbiological criteria. Thus, although, grading systems for shell eggs may vary from country to country or region to region, however, regardless of the grading or classification system used egg shell quality and interior quality are important factors in determining egg quality.

Egg shell quality: The vast majority of eggs sold in Nigeria are sold in their shell and a consumer's first impression of any egg purchased is based on their perception of shell quality (Okoli and Udedibie, 2000). Shell quality is one of the most important factors that influence hatchability (Roque and Soares, 1994). The productivity and quality of breeding eggs have an overall significant for the continuity of the flocks and for an economic breeding (Ojedapo *et al.*, 2009). Peebles and Brake (1987) reported that reduction in eggs shell quality depressed hatchability and result in weakening of embryos. Egg shell quality has significant impact on the reproductive fitness of the parent (Bennett, 1992; Abanikannda *et al.*, 2007). The shell thickness and porosity help to regulate the exchange of carbon dioxide and oxygen between the developing embryo and the air during incubation (Roque and Soares, 1994). Shell thickness also has a very significant effect on moisture loss during incubation (Bennett, 1992). Thin-shelled eggs lose more

Table 2: Causes of shell quality problems

Condition of shell	Possible causes
A. Odd shaped	<ol style="list-style-type: none"> 1. Inherited 2. Disease: Newcastle disease, infectious bronchitis, laryngotracheitis, egg drop syndrome 76 3. Age of hens: incidence is higher in older hens
B. Thin, porous or shell-less	<ol style="list-style-type: none"> 1. Inheritance influences porosity and ability to produce strong shells 2. Lack of sufficient calcium, phosphorus, manganese or vitamin D 3 3. Vitamin D 2 mistakenly substituted for D 3 4. Excess phosphorus consumption, especially by older hens 5. Ingestion of sulfanilamide (sulfa drugs) 6. Disease: Newcastle disease, infectious bronchitis, avian influenza, egg drop syndrome 76 7. Hens exposed to temperature over 85-90°F 8. Age of hens: incidence higher with older hens 9. Premature laying of the egg
C. Rough or abnormal shell texture	<ol style="list-style-type: none"> 1. Inherited 2. Newcastle disease or infectious bronchitis 3. Excessive use of antibiotics 4. Excess calcium consumption by the hens 5. Copper deficiency
D. Mottled shells	<ol style="list-style-type: none"> 1. Primarily caused by high or low extremes in humidity 2. Inherited 3. Manganese deficiency 4. Artificially induced 5. White strain layers producing tinted
E. White strain layers producing tinted eggs	<ol style="list-style-type: none"> 1. Primarily inherited
F. Yellow shells	<ol style="list-style-type: none"> 1. Extended use of high levels of certain antibiotics
G. Tremulous or loose air cells	<ol style="list-style-type: none"> 1. Newcastle disease 2. Infectious bronchitis 3. Rough handling of eggs 4. Eggs stored large end down
H. Depigmented brown shell	<ol style="list-style-type: none"> 1. Infectious bronchitis 2. High stress in the flock 3. Egg Drop Syndrome 76

Source: Jacob *et al.* (2000)

moisture than do thick-shelled eggs, causing the chick to have difficulty hatching (Roque and Soares, 1994). The Egg Producers Federation of New Zealand (Inc) (EPF) Code of Practice (EPF and NZFSA, 2002) lists 14 possible eggs shell defects, although, these can be grouped into five main categories, defects associated with egg shell integrity, texture, shape, colour and cleanliness. Some of the causes of shell quality problems are detailed in Table 2.

Egg shell integrity: Defects considered under the category of egg shell integrity include gross cracks, hairline cracks, star cracks and thin shelled or shell-less eggs. As cracked eggs cannot be made available for retail sale (Food Standards Australia New Zealand, 2006), high number of cracked eggs will have a negative impact on the profitability of any egg producer. One of the most obvious reasons for egg shell cracks (including gross cracks, hairline cracks and star cracks) is

mechanical damage (Awosanya *et al.*, 1998) caused either by the birds themselves or as a result of poor management practices, such as infrequent collection of eggs, rough handling and poor design and/or maintenance of the cage floor. Egg shell strength ultimately affects the soundness of the shell, with weaker shelled eggs more prone to cracks and breakages and subsequently microbial contamination (Yoruk *et al.*, 2004). Shell strength can be affected by a wide range of factors including:

Egg size: Smaller eggs have stronger shells than larger ones, as hens have a finite capacity to deposit calcium in the shell and as a result, the same amount of calcium is spread over a larger area (Butcher and Miles, 2003; Rajkumar *et al.*, 2009).

Age of bird: Older birds tend to lay bigger eggs and have a higher egg output, which impacts on shell strength as described above (Butcher and Miles, 2003). Very young birds with immature shell glands may produce shell-less eggs or eggs with very thin shells (Rajkumar *et al.*, 2009). Delaying the onset of sexual maturity by one to two weeks will prevent this (Coutts and Wilson, 1990).

Stress: A single stress or disturbance to a flock of laying hens can be enough to de-synchronise the process of egg formation for several days, during which time, a number of different egg quality faults may be seen (Clunies *et al.*, 1992). For example:

- Any factor which results in oviposition prior to completion of shell deposition will result in soft or thin-shelled eggs. Activities which create disturbances in and around the layer shed should be minimised (Coutts and Wilson, 1990)
- If an egg is retained in the shell gland, any subsequent egg laid may spend less time than normal in the shell gland, resulting in insufficient shell deposition and a soft shelled or shell-less egg (Clunies *et al.*, 1992). Solomon (1991) noted that once an imperfect mammillary layer has formed (as a result of stress experienced before the egg reaches the shell gland), subsequent layers are disorganized and thin or soft shelled eggs are a common phenomenon after stress

Elevated environmental temperature: High (above 25°C) environmental or shed temperatures may affect the feed (and therefore calcium) intake of the bird, thus, resulting in a decreased availability of calcium for shell deposition (Okoli *et al.*, 2006). As well as decreasing feed intake, laying hens will try to overcome heat stress by panting (Koelkebeck, 1999). However, this causes a decrease in the amount of carbon dioxide (CO₂) in the hens' blood, a condition known as respiratory alkalosis (Noddegaard, 1992; Koelkebeck, 1999). As egg shells are made up of 95% calcium carbonate (CaCO₃), this decrease in blood CO₂ levels, combined with an increase in blood pH and a subsequent decrease in Ca²⁺ ions for shell formation leads to an increase in the number of thin or soft shelled eggs produced. Arima *et al.* (1976) found that the egg quality of older hens was more severely affected by increased temperature than younger hens.

Nutrition and water quality: The provision of adequate dietary minerals and vitamins is essential for good eggshell quality (Yoruk *et al.*, 2004). Similarly, as water quality varies from country to country and region, the role of drinking water in mineral and trace element supply should not be overlooked (Anyanwu *et al.*, 2008). Calcium and phosphorous are essential macro

minerals with calcium forming a significant component of the shell and phosphorous playing an important role in skeletal calcium deposition (Frost and Roland, 1991) and subsequent availability of calcium for egg shell formation during the dark period (Boorman *et al.*, 1989). Coetzee (2002) investigated the effect of calcium levels in drinking water on shell integrity in South African laying hens and demonstrated that birds supplied an additional 200 mg of calcium per litre of drinking water laid eggs with mean shell strength of 42.6 ± 9.0 . This was in comparison to those receiving un-supplemented water, whose eggs had mean shell strength of 38.9 ± 7.0 .

However, the feeding of calcium levels above the requirement of the bird for production has not been shown to improve shell quality (Keshavarz and Nakajima, 1993). Indeed, feeding hens high levels of calcium may interfere with the availability of other minerals (NRC, 1994) and can have a negative impact on the ability of the bird to utilise calcium, particularly if calcium levels in the diet are subsequently decreased. This is particularly true in young pullets. Wideman and Lent (1991) also reported that high levels of calcium can cause latent kidney damage, which has a long term impact on bird survivability.

Similarly, feeding high levels of dietary phosphorous have also been shown to have a negative effect on eggshell quality (Keshavarz and Austic, 1990; Harms, 1982a, b; Boorman *et al.*, 1989; Taylor, 1965; Arscott *et al.*, 1962). Although, Miles and Harms (1982) showed a clear linear negative correlation between specific gravity and plasma phosphate, it remains unclear whether excess plasma phosphorous interferes with the mobilization of skeletal phosphorous reserves or the shelling process itself (Boorman *et al.*, 1989; Ahmad and Balander, 2004) or if elevated levels of phosphorous increases calcium excretion (Keshavarz and Austic, 1990).

Keshavarz and Austic (1990) also examined the interaction of phosphorus and chloride and the role of chloride in egg shell integrity. As with phosphorous, elevated dietary levels of chloride resulted in decreased eggshell quality and lower levels of blood acid-base indicators. This is supported by the work of Balnave *et al.* (1989), who observed an increase in shell defects with no changes in egg production, egg weight, feed or water intake, blood-acid base levels and electrolyte levels for birds provided with 2000 mg of sodium chloride (NaCl) per litre drinking water. This was in comparison to those hens provided water with 600 mg NaCl per litre. Birds receiving 2000 mg of NaCl per litre drinking water also had an increased incidence of shell-less eggs. Egg shell defects persisted after the sodium chloride was removed from the drinking water. In contrast, Hess and Britton (1989) fed hens diets low in chloride and found virtually no effect on shell quality.

Trace mineral nutrition is a complex area of animal nutrition and a wide range of interactions and antagonisms can result in poor absorption or utilization of minerals essential for shell formation (Frost and Roland, 1991). It should also be noted that not all trace mineral sources are equally available and consideration should be given to this aspect of premix formulation. Burley and Vadehra (1989), Miles (2001) and Mas and Arola (1985) reported varying amounts of zinc, copper, iron and manganese in the shell and its associated membranes. It is clear, therefore, that provision of adequate levels of these minerals, which are key components of the shell matrix and play an essential role as co-enzymes, is essential if shell integrity is to be maintained. Vitamin D also plays an important role in the proper utilization of calcium and phosphorous and sufficient amounts of this vitamin should be included in the feed (Amiri Andi *et al.*, 2006).

Finally, care should be paid to the mixing of the diet. Thorough mixing of the feed is essential if each bird in the flock is to receive a similar amount of any given nutrient (Natalie, 2009). This is particularly true for layer hen diets which frequently contain raw materials with a wide range of different densities.

Mycotoxigenesis: Jewers (1990) reported that thin rubbery shells which break more readily than normal have been observed during field outbreaks of ochratoxigenesis. Similarly, in an outbreak of T-2 toxicosis, egg breakages increased from a normal 3 to 15% with a further 18% of eggs broken in transit to customers (Jewers, 1990). Zaghini *et al.* (2005) reported that birds fed diets containing 2.5 ppm of aflatoxin B1 had lower egg shell weights than those fed the control diet or diets supplemented with mannanoligosaccharides.

Genetics: Clunies *et al.* (1992) found that hens laying thick-shelled eggs retained more dietary calcium than those laying thin-shelled eggs. Although, there was no difference in egg production between thick and thin shell layers, both egg and shell weight were greater for the thick shelled eggs.

Disease: Infectious Bronchitis (IB), a viral disease caused by a coronavirus which attacks the mucus membranes of the respiratory and reproductive tracts (Butcher and Miles, 2003; Cavanagh and Naqi, 2003), may result in egg defects. These include pale shelled eggs and eggs with poor shell structure and integrity (Butcher and Miles, 2003; Khan *et al.*, 2004). Similarly, birds affected by Egg Drop Syndrome (EDS), caused by an adenovirus, initially produce pale eggs, quickly followed by thin-, soft-shelled or shell-less eggs (McFerran and Adair, 2003).

Texture: Rough or sandpaper shells, pimples, pinholes and mottled or glassy-shells are all egg shell defects associated with egg shell texture. These defects are frequently a result of bird age, but may also be caused by other factors (Coutts and Wilson, 1990; MAFF, 1976).

Disease: Certain diseases such as IB, Infectious Laryngotracheitis (ILT) (Spadbrow, 1993) and Avian Encephalomyelitis (AE) (Coutts and Wilson, 1990) have been implicated in the production of rough or sandpaper eggshells.

Mycotoxigenesis: As discussed above, ochratoxigenesis may result in rubbery-shelled eggs (Jewers, 1990).

Genetics: The production of eggs with calcium deposits on the shell (or pimples) is thought to be hereditary (Khan *et al.*, 2004).

Management: Overcrowding of birds, changes in the lighting programme, poor shed ventilation and inadequate water supply can contribute to increased incidence of shell defects associated with egg texture (Etuk *et al.*, 2004).

Shape: Misshapen eggs have a shape which differs from the smooth normal shape (for example, flat sided eggs and body checked eggs). This can be caused by a number of factors:

Age of bird: As with shell soundness, young birds with immature shell glands may produce misshapen eggs (Hess and Britton, 1989).

Stress: Body checked eggs, marked by grooves and ridges, occur when the shell of the egg breaks in the shell gland, during the formation process (i.e., 10-14 h before the egg is laid)

(Abanikannda *et al.*, 2007). Although, the damage can be partly repaired, a bulge forms around the egg (Solomon, 1991). Flat sided eggs occur where two eggs are in the shell gland at the same time (Solomon, 1991; Abanikannda *et al.*, 2007). Both defects may be caused by overcrowding, frights or other disturbances and poor lighting patterns (Koelkebeck, 1999; Coutts and Wilson, 1990). Jones (2006) stated that proper handling can reduce the incidence of body checks.

Disease: As the albumen of the egg and surrounding membranes provides the structure on which the egg shell is deposited, if the albumen quality is very poor, there is not sound foundation on which to build the shell (Abanikannda *et al.*, 2007). As a result, those diseases which result in poor albumen quality often cause an increase in the number of misshapen eggs. Examples of these are IB (Butcher and Miles, 2003; Cavanagh and Naqi, 2003), EDS (McFerran and Adair, 2003) and certain strains of Newcastle Disease (NCD) or Avian Influenza (AI) (Spadbrow, 1993).

Colour: The colour of an egg shell is determined primarily by the genetics of the hen, with white feathered hens laying white eggs and brown feathered hens laying brown eggs (Fairfull and Gowe, 1990). During the process of egg shell formation in brown egg layers, the epithelial cells lining the surface of the shell gland synthesise and accumulate pigments, such as biliverdin-IX, its zinc chelate and protoporphyrin-IX (Butcher and Miles, 2003). In the final three to four hours of shell formation these pigments are transferred to the viscous, protein rich cuticle. The quantity of pigment in the cuticle which determines the colour of the egg (Fairfull and Gowe, 1990). As the cuticle is deposited onto the eggshell at the same time that shell deposition reaches a plateau (approximately 90 min prior to oviposition), pigment distribution is not uniform throughout the shell, with very little pigment contained in the shell itself (Khan *et al.*, 2004). Thus, any factor which causes a disruption, either in the ability of the epithelial cells to synthesise pigment or in the deposition of the cuticle, will affect the colour of the egg shell. These factors include:

Stress: Epinephrine, a stress hormone, will cause a delay in oviposition and cessation of shell gland cuticle formation, which can cause pale shelled eggs to be produced. Stressors may include, amongst others, high cage density, loud noise and handling (Natalie, 2009).

Age of bird: As birds age increases, the intensity of pigment decreases. This may be due to decreasing production of pigment or increased surface area over which available pigment is distributed (Abdullah *et al.*, 2003).

Chemotherapeutic agents: Certain drugs have been shown to affect egg shell colour. For example, nicarbazin (an anticoccidial drug) fed at a level of 5 mg day⁻¹ can result in the production of pale eggs within 24 h, while higher doses can lead to complete depigmentation (Cavanagh and Naqi, 2003). Chlortetracycline (600-800 ppm) may also result in yellow egg shells (Cavanagh and Naqi, 2003).

Disease: Viruses, which affect the mucus membranes of the respiratory and reproductive tract, such as NCD and IB, not only cause a decrease in egg production, but also cause the shell to become abnormally thin and pale (Spadbrow, 1993).

Cleanliness: Cleanliness is probably the easiest aspect of egg shell quality control and good management plays an important role. Most eggs are clean when laid and subsequently become

Table 3: Summary of standards for interior quality of chicken eggs by candling

Interior quality factor	AA quality	A quality	B quality	Inedible
Air cell	1/8 inch or less in depth	3/16 inch or less in depth	More than 3/16 inch	Doesn't apply
White (albumen)	- Clear- Firm	- Clear-May bereasonably firm	- Clear-May be weak and watery	Doesn't apply
Yolk	Outline slightly defined	Outline may be fairly well-defined	Outline clearly visible	Doesn't apply
Spots (blood or meat)	None	None	Blood or meat spots aggregating not more than 1/8 inch in diameter	Blood or meat spots aggregating more than 1/8 inch in diameter

contaminated with faecal material or other contaminants. Defects which fall into this category, include cage marks, stained eggs and fly marks (EPF and NZFSA, 2002). Although, fungus or mildew on shells is listed as a defect, it is only likely to occur in poor conditions. Good management practices will help reduce the number of dirty eggs (Etuk *et al.*, 2004). These practices include frequent collection of eggs, as well as regular replacement of litter material in nest boxes or regular maintenance and cleaning of cage floors and rollout trays. Whilst less common, fly stains, water stains and grease or oil stains may occur and can be prevented by good shed and equipment maintenance or management (Etuk *et al.*, 2005). Moreover, factor which causes diarrhoea in the birds, (for example high dietary salt levels), will also result in an increase in the number of dirty eggs collected. Blood smears on eggs can be minimised by good pullet management, including lighting and beak trimming if necessary.

Internal egg quality: Unlike external (shell) quality, internal quality of the egg begins to decline as soon as the egg is laid (Okoli and Udedibie, 2001). Table 3 shows detail summary of standards for interior quality of chicken eggs by candling. Thus, although factors associated with the management and nutrition of the hen do play a role in internal egg quality, egg handling and storage practices do have a significant impact on the quality of the egg reaching the consumer (Okeudo *et al.*, 2003). Similarly, although the shell provides a unique package for the distribution of the egg contents to the consumer, it is in fact the internal quality of the egg that is most important to the consumer (Okeudo *et al.*, 2003). These aspects of internal quality are significantly more difficult to observe or evaluate in the intact egg, even with the use of candling. In addition to the obvious, nutritional quality of the egg, internal egg quality is extremely important because of its many functional and aesthetic properties. For example, eggs are used as thickening agents in custards and puddings, egg whites are used as smoothing agents to give icings a desirable texture and egg yolks add colour and richness to food (Okeudo *et al.*, 2003).

In recent years, much attention has been focused on increasing the omega 3:6 ratio and vitamin content of eggs, principally through manipulation of the diet. However, although these enhancements further complicate the issue of egg quality (Anyanwu *et al.*, 2008). The EPF Code of Practice (EPF and NZFSA, 2002) lists a total of nine internal defects and these can be broadly categorised into three groups; namely: defects affecting yolk quality, defects affecting albumin quality and defects affecting overall quality.

Yolk quality: Yolk quality is determined by the colour, texture, firmness and smell of the yolk.

Yolk colour: Although, yolk colour is a key factor in any consumer survey relating to egg quality (Okeudo *et al.*, 2003), consumer preferences for yolk colour are highly subjective and vary widely

from country to country. The primary determinant of yolk colour is the xanthophyll (plant pigment) content of the diet consumed (Silversides *et al.*, 2006). It is possible to manipulate the yolk colour of eggs by the addition of natural or synthetic xanthophylls to layer hen feeds. This ability to readily manipulate egg yolk colour can be an advantage in meeting market demands. However, the ease with which yolk colour can be manipulated can lead to unwanted colour changes. For example, the inclusion of higher than recommended levels or incorrect ratios of pigments can lead to orange-red yolks (Silversides *et al.*, 2006). Similarly, diphenyl-para-phenylenediamine (DPPD), an antioxidant, has been reported to cause excessive deposition of pigments in the egg yolk.

The inclusion of more than 5% cottonseed meal in a layer diet will result in olive or salmon coloured yolks (Esonu, 2006), while the inclusion of certain weeds or weed seeds may result in green yolks. Similarly, inadvertent omission of xanthophylls from the diet will lead to pale yolks (Esonu, 2006). Both inadequate mixing of the diet as well as excessive mixing of the diet will also result in a heterogeneous feed and subsequent variation in the amount of xanthophylls consumed by each hen in the flock. This will result in egg yolk colour not being uniform throughout the flock.

Pale yolks can result from any factor which alters or prevents the absorption of pigments from the diet or the deposition of these pigments in the yolk. These factors could include:

- Worms (Coutts and Wilson, 1990)
- Any factor which inhibits liver function, subsequent lipids metabolism and deposition of pigment in the yolk. For example, mycotoxicosis caused by aflatoxin B1 (Zaghini *et al.*, 2005)
- Coccidiosis, although this is rare in adult hens

Mottled yolks (with many pale spots and blotches which vary in colour, size and shape), occur when the contents of the albumen and yolk mix as a result of degeneration and increase permeability of the vitelline membrane (Amiri Andi *et al.*, 2006). Factors affecting mottling were reviewed in detail by Cunningham and Sanford (1974). Dietary factors which may cause mottled yolks include:

- The presence of nicarbazin (an anticoccidial agent) in the feed has shown by numerous authors to cause mottling (Jones *et al.*, 1990; Cunningham and Sanford, 1974)
- Worming drugs such as phenothiazine, dibutyltindialaurate (Berry *et al.*, 1968) and Piperazine (Amiri Andi *et al.*, 2006). However, Berry *et al.* (1968) did not observe yolk defects when Piperazine was fed at the manufacturer's recommendations. Similarly, they only observed defects when dibutyltindialaurate was fed at the recommended level but over a much longer period.
- Gossypol from cotton seed meal (Berry *et al.*, 1968; Esonu, 2006)
- Certain antioxidants such as gallic acid (from grapes, tea and oak bark) and tannic acid, or tannins from grains such as sorghum (Esonu, 2006)
- Calcium deficient diets (McCready *et al.*, 1972; Cunningham and Sanford, 1974)

Storage time and temperature has also been shown to affect the degree of egg yolk mottling (Okoli and Udedibie, 2001). Okeudo *et al.* (2003) stated that as the internal temperature of the egg increases above 7°C, the protein structures of the thick albumen and vitelline membrane breakdown faster. As the membrane degenerates during storage, water enters the yolk causing mottling and after prolonged storage, albumen proteins also enter the yolk increasing the severity

of mottling. In order to reduce the rate of breakdown of the vitelline membrane, eggs should be collected regularly, reducing the time they are exposed to higher environmental temperatures and contaminants and stored at temperatures of 7-13°C and humidity of 50-60%. Cunningham and Sanford (1974) also identified hen age, oil coating of eggs and movement of eggs as possible factors affecting mottling of eggs.

Yolk firmness: The yolk of a freshly laid egg is round and firm (Okoli and Udedibie, 2000). However, as the egg ages and the vitelline membrane degenerates, water from the albumen moves into the yolk and gives the yolk a flattened shape.

Yolk texture: Rubbery yolks may be caused by severe chilling or freezing of intact eggs, the consumption of crude cottonseed oil or the seeds of some weeds (Jacob *et al.*, 2000).

Albumin quality: Albumin quality is related to the consistency, appearance and the functional properties.

Consistency: Albumin quality is measured in terms of Haugh Units (HU) calculated from the height of the albumin and the weight of the egg (Haugh, 1993). A minimum measurement in HU for eggs reaching the consumer is 60. However most eggs leaving the farm should be between 75 and 85 HU (Zaman *et al.*, 2005). Albumin consistency is influenced by:

Age of the hen: HU will decrease with increasing bird age value, with HU decreasing by around 1.5 to 2 units (Awosanya *et al.*, 1998) for each month in lay. Doyon *et al.* (1986) stated that HU decreases at a fairly constant rate of 0.0458 units day⁻¹ of lay as the hen ages. Doyon *et al.* (1986) also noted that in an ideal situation, HU should be on average 102 at 20 weeks of age, falling to an average of 74 HU by 78 weeks of age.

Genetics: Strain of bird has also been shown to play a role in albumin consistency, with some strains consistently producing eggs with thin albumin. Rajkumar *et al.* (2009) reported that brown egg layers produced eggs with higher HU, while other authors (Sell *et al.*, 1982; Williams, 1992) reported that HU values were more variable within the brown egg layers compared with those that lay white eggs. High producing birds tend to lay eggs with relatively lower amounts of thick albumin and, although this can be influenced by selective breeding, egg numbers are usually considered more important.

Age and storage of the egg: As the egg ages and carbon dioxide (CO₂) is lost through the shell, the contents of the egg become more alkaline, causing the albumin to become transparent and increasingly watery (Okeudo *et al.*, 2003). At higher temperatures, loss of CO₂ is faster and the albumin quality deteriorates faster. Decreasing shed temperatures in the hotter months, combined with regular collection of eggs will help to reduce deterioration of the albumin before collection.

Eggs stored at ambient temperatures and humidity lower than 70% will lose 10-15 HU in a few days from point of lay. By 35 days, these eggs will lose up to 30 HU (Natalie, 2009). Storage of eggs at temperatures of 7-13°C and a humidity of 50-60% (as discussed under mottling), will reduce the rate of degeneration of thick albumen proteins and, consequently, egg albumin quality will be maintained for longer (Jones, 2006). Oiling of eggs can also help to reduce CO₂ losses and thus help

maintain internal egg quality (Okoli and Udedibie, 2000, 2001; Okeudo *et al.*, 2003) but is not a substitute for cool storage.

Vanadium: Henry and Miles (2001) reviewed the effects of vanadium on poultry performance. They noted that poorer albumin quality has been reported from laying hens consuming as little as 6 ppm. Sell *et al.* (1986) showed that the interior quality, of eggs decreased in two strains of laying hens fed 3 or 6 ppm added vanadium. Duyck *et al.* (1990) fed laying hens 10 ppm of vanadium for 30 days. HU from these hens averaged 71 HU after one day of storage (62°F (16.6°C) and 60% Relative Humidity) and 64 after seven days of storage. This was in contrast to the average of 82 and 74 HU after one and seven days storage, respectively, observed for hens fed the control diet. Henry and Miles (2001) reported that the negative effects of vanadium may be overcome by feeding cottonseed meal, ascorbic acid, vitamin E or carotene, although this is dose dependant.

Diseases: Diseases such as certain strains of EDS, IB, NCD and ILT (Jacob *et al.*, 2003) can all cause a decrease in albumin consistency.

Appearance: Normal albumin is transparent, with a slightly yellow green colour. Discolouration of the albumin may occur if the eggs are stored for an extended time in poor conditions, with the albumin becoming much yellower (Cavanagh and Naqi, 2003). Cyclopropene fatty acids from cottonseed meal and the certain weed seeds (Sell *et al.*, 1986) can cause albumin to turn pink after storage. Green whites are caused by excesses of riboflavin (vitamin B2) in the diet. Cloudy whites may be caused by the oiling of eggs within 6 h of lay (Silversides *et al.*, 2006).

Overall quality

Blood spots: Blood spots may vary from indistinguishable spots on the surface of the yolk to heavy contamination throughout the yolk (Cavanagh and Naqi, 2003). Although, blood spots are normally closely associated with the yolk, occasionally blood may be diffused through the albumin. Blood spots occur when small blood vessels in the ovary rupture when the yolk is released. Vitamin K plays an important role in blood clotting. Vitamin K deficiency can result in an increased occurrence of blood spots (Bains, 1999). Some strains of birds appear to be predisposed to blood spots although the incidence is low (Rajkumar *et al.*, 2009). Avian encephalomyelitis has been reported as a cause of blood spots (McFerran and Adair, 2003). Jewers (1990) reported an increase in blood spots from essentially 0 to 3% in birds affected with T-2 toxicosis. Bains (1999) suggested that mycotoxicosis may reduce vitamin K absorption and this may explain the elevated incidence of blood spots in hens affected by T-2 toxicosis.

Meat spots: These are usually associated with the albumin rather than the yolk and often consist of small pieces of body tissue (Curtis *et al.*, 1985). However, some may consist of partially broken down blood spots or pigments. The occurrence of blood spots varies with strain of bird, increases with age of bird and is reported to be higher in brown egg layers (Abdullah *et al.*, 2003).

Bacterial or fungal contamination: Solomon (1991) suggested that while pores on the surface of the egg do represent possible ports of entry for bacteria, particularly as the cuticle hardens just after oviposition, these are of secondary importance to the structural defects that may occur. Structural defects, because of their magnitude, offer a much more likely route for bacteria to enter

the egg contents. Bacterial and fungal contamination of eggs usually results in black, red or green rot. The egg looks and smells putrid when broken out of the shell. Bacterial and fungal contamination of eggs, resulting from faecal contamination of the egg, can be prevented by good management practices, including regular replacement of nesting materials or good cage maintenance as appropriate (Etuk *et al.*, 2004). Bacterial contamination of the egg contents may also occur as a result of an infection in the oviduct of the hen and any affected hens should be culled (Etuk *et al.*, 2005).

Proper handling and storage of eggs following collection will minimise the opportunity for bacterial or fungal contamination. However, improper washing procedures, high storage temperatures and humidity will increase the incidence of bacterial or fungal contamination (Etuk *et al.*, 2005). Careful attention should be paid to feed source, as *Salmonella* sp. can be transmitted through the feed.

Roundworms in eggs: Burley and Vadehra (1989) reported that where roundworm infestation of the intestinal tract occurs, worms may migrate from the cloaca into the oviduct and become enclosed in the egg. This can be prevented by good flock management.

Off odours/flavours: Although, off odours and flavours are rare if eggs are stored correctly (Okoli and Udedibie, 2000), eggs readily absorb strong odours or flavours. Storage of eggs in close proximity to fish oils and meals, sour milk, strongly scented or decaying fruit and vegetables, mould, disinfectants and kerosene is likely to result in the development of off odours or flavours (Okeudo *et al.*, 2003). However, eggs that have been oiled are less likely to absorb foreign odours. Old eggs and eggs stored at high temperatures are more likely to exhibit off odours or flavours (Okoli and Udedibie, 2000). Other causes of off odours or flavours include strongly flavoured feed ingredients such as fish meal or fish oil, some vegetables (including onions, turnips and excessive amounts of cabbage) and rapeseed or canola.

CONCLUSIONS

The consumer seldom sees many of the egg defects detailed above, as eggs are graded and most defects are normally removed prior to retail. Assuming adequate nutrition, the use of suitable raw materials, proper feed hygiene and a lack of contamination of the birds' diet with foreign matter and/or objects, there is little effect of nutrition on egg quality defects. Flock shed and feed management, egg storage and egg handling remain the three most important factors in determining the incidence of egg quality defects. However, the breeding companies should be aware of the market requirements for egg quality and should include these traits in their selection programmes, where possible. Finally, I will like to call on the Product Standard Organisation of Nigeria in collaboration with the Animal Product Researchers and Ministry of Agriculture in Nigeria to develop a standard and have a defined grade for shell eggs sold in Nigeria to meet the international standard and consumer's demand and aspiration.

REFERENCES

- Abanikannda, O.T.F., A.O. Leigh, L.O. Ojedapo and O.D. Kpossu, 2007. Interspecific variation in shell thickness of commercial poultry eggs. Proceedings of the 12th Annual Conference of Animal Science Association of Nigeria, Sept. 10-13, Nigeria, pp: 9-9.

- Abdullah, A.R., L.O. Ojedapo, T.A. Adedeji, T.B. Olayeni and O.S. Adedeji, 2003. Influence of hees age on egg quality parameters in Bovans Nera black layer strain. Proceedings of the 28th Annual Conferences of the Nigerian Society for Animal Producton, March 18-22, University of Calabar, Cross Rivers State, pp: 103-103.
- Ahmad, H.A. and R.J. Balander, 2004. Physiological responses of layers to alternative feeding regimen of calcium source and phosphorous level. *Int. J. Poult. Sci.*, 3: 100-111.
- Amiri Andi, M., M. Shivazad, S.A. Pourbakhsh, M. Afshar and H. Rokni *et al.*, 2006. Effects of vitamin E in broiler breeder diet on hatchability, egg quality and breeder and day old chick immunity. *Pak. J. Biol. Sci.*, 9: 789-794.
- Anyanwu, G.A., E.B. Etuk, I.C. Okoli and A.B.I. Udedibie, 2008. Performance and egg quality characteristics of layers fed different combinations of cassava root meal and bambara groundnut offal. *Asian J. Poult. Sci.*, 2: 36-41.
- Arima, Y., F.B. Mather and M.M. Ahmad, 1976. Response of egg production and shell quality, to increases in environmental temperature in two age groups of hens. *Poult. Sci.*, 55: 818-820.
- Arscott, G.H., P. Rachapaetayakom and P.E. Bernier, 1962. Influence of ascorbic acid, calcium and phosphorus on specific gravity of eggs. *Poult. Sci.*, 41: 485-488.
- Awosanya, B., J.K. Joseph and O.D. Olaosebikan, 1998. The effect of age of bird on shell quality and component yield of eggs. *Nig. J. Anim. Prod.*, 25: 68-70.
- Bains, B.S., 1999. A Guide to the Application of Vitamins in Commercial Poultry Feed. Rath Design Communications, Australia, pp:200.
- Balnave, D., I. Yoselewitz and R.J. Dixon, 1989. Physiological changes associated with the production of defective eggshells by hens receiving sodium chloride in the drinking water. *Br. J. Nutr.*, 61: 35-43.
- Bennett, C.D., 1992. The influence of shell thickness in commercial broiler breeder flocks. *J. Pure Applied Poult.*, 1: 61-65.
- Berry, J.O., G.W. Newell, D.P. Holder, G.V. Odell and D.E. Bee, 1968. The effect of cottonseed products and selected feed additives on egg yolk discoloration. *Poult. Sci.*, 47: 783-794.
- Boorman, K.N., J.G. Volynchook and C.G. Belyavin, 1989. Egg Shell Formation and Quality. In: *Recent Developments in Poultry Nutrition*, Cole, D.J.A. and W. Haresign (Eds.). Butterworths, Kent, England.
- Burley, R.W. and D.V. Vadehra, 1989. *The Avian Egg Chemistry and Biology*. John Wiley and Sons, New York, NY.
- Butcher, G.D. and R.D. Miles, 2003. Factors causing poor pigmentation of brown shelled eggs. University of Florida. <http://edis.ifas.ufl.edu/pdffiles/VM/VM04700.pdf>
- Cavanagh, D. and S.A. Naqi, 2003. Infectious Bronchitis. In: *Diseases of Poultry*, Saif, Y.M., H.J. Barnes, J.R. Glisson, A.M. Fadly, L.R. McDougald and D.E. Swayne (Eds.). Iowa State University Press, Ames, IA., USA., pp: 101-120.
- Clunies, M., D. Parks and S. Leeson, 1992. Calcium and phosphorus metabolism and eggshell thickness in laying hens producing thick or thin shells. *Poult. Sci.*, 71: 490-498.
- Coetzee, C., 2002. The effect of elevated Calcium Levels in drinking water on shell integrity. *Spesfeed News*, Summer 2002. <http://www.spesfeed.co.za/Summer2002.htm>
- Council of the European Union, 2006. Council Regulation (EC) No 1028/2006 of 19 June 2006 on Marketing Standards for Eggs. http://europa.eu.int/eurlex/en/consleg/pdf/2003/en_2003R2295_do_001.pdf

- Coutts, J.A. and G.C. Wilson, 1990. Egg Quality Handbook. Queensland Department of Primary Industries, Australia.
- Cunningham, F.E. and P.E. Sanford, 1974. A review of facts influencing egg yolk mottling. *World's Poult. Sci. J.*, 30: 103-114.
- Curtis, P.A., F.A. Gardner and D.B. Mellor, 1985. A comparison of selected quality and compositional characteristics of brown and white shell eggs. 2. interior quality. *Poult. Sci.*, 64: 302-306.
- Doyon, G., M. Bernier-Cardou, R.M.G. Hamilton, F. Castaigne and C.J. Randall, 1986. Egg quality. 2. Albumen quality of eggs from five commercial strains of White Leghorn hens during one year of lay. *Poult. Sci.*, 65: 63-66.
- Duyck, S.K., R.D. Miles, A.F. Rossi and P.R. Henry, 1990. Effect of time and storage conditions on interior egg quality from hens fed vanadium. *Poult. Sci.*, 69: 164-164.
- EPF and NZFSA, 2002. Egg producers federation of New Zealand (Inc) code of practice (Includes Requirements for Risk Management Programmes). <http://www.nzfsa.govt.nz/consultation/egg-cop/discussion-paper/egg-cop-ch1.pdf>
- Esonu, B.O., 2006. Animal Nutrition and Feeding: A Functional Approach. 2nd Edn., Rukzeal and Ruksons Associates Memory Press, Owerri, Imo State, Nigeria.
- Etuk, E.B., I.C. Okoli and A.B.I. Udedibie, 2005. Priority issues in tropical animal health management. *Anim. Prod. Res. Adv.*, 1: 83-92.
- Etuk, E.B., I.C. Okoli and M.U. Uko, 2004. Prevalence and management issues associated with poultry coccidiosis in Abak agricultural zone of Akwa Ibom state, Nigeria. *Int. J. Poult. Sci.*, 3: 135-139.
- European Commission, 2003. Commission regulation (EC) No 2295/2003 of 23 December 2003, introducing detailed rules for implementing Council Regulation (EEC) No 1907/90 on certain marketing standards for eggs. http://eurlex.europa.eu/LexUriServ/site/en/oj/2006/l_186/l_18620060707en00010005.pdf
- Fairfull, R.W. and R.S. Gowe, 1990. Genetics of Egg Production in Chickens. In: *Poultry Breeding and Genetics*, Crawford, R.D. (Ed.). Elsevier, New York, NY.
- Food Standards Australia New Zealand, 2006. Australia New Zealand Food Standards Code, Standard 2.2.2. http://www.foodstandards.gov.au/srcfiles/Standard_2_2_2_Egg_Products_v64.pdf
- Frost, T.J. and D.A. Roland, 1991. The influence of various calcium and phosphorus levels on tibia strength and eggshell quality of pullets during peak production. *Poult. Sci.*, 70: 963-969.
- Harms, R.H., 1982a. The influence of nutrition on eggshell quality. Part II. Phosphorous. *Feedstuffs*, May 10, pp: 25-26.
- Harms, R.H., 1982b. The influence of nutrition on eggshell quality. Part III. Electrolyte Balance. *Feedstuffs*, May 17, pp: 25-28.
- Haugh, R.R., 1993. The Haugh unit for measuring egg quality. *US Egg Poult. Magazine*, 43: 552-555.
- Henry, P.R. and R.D. Miles, 2001. Heavy metals-vanadium in poultry. *Ciência Anim. Brasileira*, 2: 11-26.
- Hess, J.B. and W.M. Britton, 1989. The effect of dietary chloride or protein changes on eggshell pimpling and shell quality in late production leghorn hens. *Nutr. Rep. Int.*, 40: 1107-1115.
- Jacob, J.P., R.D. Miles and F.B. Mather, 2000. Egg quality. University of Florida. <http://edis.ifas.ufl.edu/pdffiles/PS/PS02000.PDF>.

- Jacob, J.P., H.R. Wilson, R.D. Miles and F.B. Mather, 2003. Factors affecting egg production in backyard chicken flocks. University of Florida. <http://edis.ifas.ufl.edu/pdf/PS/PS02900.pdf>
- Jewers, K., 1990. Mycotoxins and their effect on poultry production. *Options Mediterraneennes Serie A.*, 7: 195-202.
- Jones, J.E., J. Solis, B.L. Hughes, D.J. Castaldo and J.E. Toler, 1990. Production and egg quality responses of White Leghorn layers to anticoccidial agents. *Poult. Sci.*, 69: 378-387.
- Jones, D.R., 2006. Conserving and monitoring shell egg quality. Proceedings of the 18th Annual Australian Poultry Science Symposium, Feb. 20-22, Australian, 157-165.
- Keshavarz, K. and R.E. Austic, 1990. Effects of dietary minerals on acid-base balance and eggshell quality in chickens. *J. Nutr.*, 120: 1360-1369.
- Kershavarz, K. and S. Nakajima, 1993. Re-evaluation of calcium and phosphorous requirements of laying hens for optimum performance and egg shell quality. *Poult. Sci.*, 72: 144-153.
- Khan, M.K.I., M.J. Khatun and A.K.M.G. Kibria, 2004. Study the quality of eggs of different genotypes of chickens under semi-scavenging system at Bangladesh. *Pak. J. Biol. Sci.*, 7: 2163-2166.
- Koelkebeck, K.W., 1999. What is egg quality and conserving it? University of Illinois. <http://www.traill.uiuc.edu/poultrynet/paperDisplay.cfm?ContentID=522>
- Kramer, A., 1951. What is quality and how can it be measured: From a food technology point of view. *Mktg. Research Workshop Report*, Michigan State College.
- MAFF, 1976. *Intensive Poultry Management for Egg Production*. MAFF, London.
- Mas, A. and L. Arola, 1985. Cadmium and lead toxicity effects on zinc, copper, nickel and iron distribution in the developing chick embryo. *Comparative Biochem. Physiol.*, 80: 185-188.
- McCready, S.T., D.A. Roland and J.L. Fry, 1972. External quality, composition and defects of eggs from hens fed a calcium deficient diet. *Poult. Sci.*, 51: 670-675.
- McFerran, J.B. and B.M. Adair, 2003. Egg Drop Syndrome. In: *Diseases of Poultry*, Saif, Y.M., H.J. Barnes, J.R. Glisson, A.M. Fadly, L.R. McDougald and D.E. Swayne (Eds.). Iowa State Press, Ames, Iowa, pp: 227-237.
- Miles, R.D. and R.H. Harms, 1982. Relationship between egg specific gravity and plasma phosphorus from hens fed different dietary calcium, phosphorus and sodium levels. *Poultry Science*, 61: 175-177.
- Miles, R.D., 2001. Trace minerals and avian embryo development. *Ciência Anim. Brasileira*, 2: 1-10.
- NRC (National Research Council), 1994. *Nutrient Requirements of Poultry*. 9th Rev. Edn., National Academy Press, Washington D.C.
- Natalie, G., 2009. Factors affecting egg quality in the commercial laying hen: A review. http://www.eggfarmers.co.nz/upload/A369F_Factors_affecting_egg_quality.pdf
- Noddegaard, F., 1992. Effect of different feeding and lighting programmes in rearing of pullets. Quality of eggs. *Arch. Geflugelk.*, 56: 271-282.
- Ojedapo, L.O., T.A. Adedeji, S.A. Ameen, T.B. Olayeni and S.R. Amao *et al.*, 2009. Effect of strain and age on egg quality characteristics of two different strains of layer chicken kept in cages in derived savanna zone of Nigeria. Proceedings of 14th Annual Conference of Animal Science Association of Nigeria (ASAN), Sept. 14-17, LAUTECH Ogbomoso, Nigeria, 1-3.
- Okeudo, N.J., C.I. Onwuchekwa and I.C. Okoli, 2003. Effect of oil treatment and length of storage on the internal quality, organoleptic attributes and microbial profile of chicken eggs. *Tropical Anim. Prod. Investigations*, 6: 63-70.

- Okoli, I.C. and A.B.I. Udedibie, 2000. Effect of oil treatment and storage temperature on egg quality. *J. Agric. Rural Develop.*, 1: 55-60.
- Okoli, I.C. and A.B.I. Udedibie, 2001. Effect of palm kernel oil treatment and method of storage on internal quality, viability and hatchability of fertile chicken eggs. *Trop. Agric. (Trinidad)*, 78: 137-140.
- Okoli, C.G., I.C. Okoli, U.V. Okorondu and M.N. Opara, 2006. Environmental and public health issues of animal food products delivery system in Imo State, Nigeria. *J. Health Allied Sci.*
- Peebles, E.D. and J. Brake, 1987. Egg shell quality and hatchability in broiler breeder eggs. *Poult. Sci.*, 66: 596-604.
- Rajkumar, U., R.P. Sharma, K.S. Rajaravindra, M. Niranjana, B.L.N. Reddy, T.K. Bhattacharya and R.N. Chatterjee, 2009. Effect of genotype and age on egg quality traits in naked neck chicken under tropical climate from India. *Int. J. Poult. Sci.*, 8: 115-1155.
- Roque, L. and M.C. Soares, 1994. Effects of eggshell quality and broiler breeder age on hatchability. *Poult. Sci.*, 73: 1838-1845.
- Sell, J.L., J.A. Arthur and I.L. Williams, 1982. Adverse effects of dietary vanadium contributed by dicalcium phosphate on egg albumin quality. *Poult. Sci.*, 61: 1541-1541.
- Sell, J.L., C.Y. Davis and S.E. Scheideler, 1986. Influence of cottonseed meal on vanadium toxicity and 48 vanadium distribution in body tissues of laying hens. *Poult. Sci.*, 65: 138-146.
- Silversides, F.G., T.A. Scott, D.R. Korver, M. Afsharmanesh and M. Hruby, 2006. A study on the interaction of xylanase and phytase enzymes in wheat-based diets fed to commercial white and brown egg laying hens. *Poult. Sci.*, 85: 297-305.
- Solomon, S.E., 1991. *Egg and Eggshell Quality*. Wolfe Publishing Ltd., London, England.
- Spadbrow, P.B., 1993. Newcastle disease in village chickens. *World's Poult. Sci. Rev.*, 5: 57-96.
- Taylor, T.G., 1965. Dietary phosphorus and egg shell thickness in the domestic fowl. *Br. Poult. Sci.*, 6: 79-87.
- Wideman, Jr. R.F. and A.L. Lent, 1991. Producer's field guide to kidney damage. *Egg Ind.*, 97: 24-30.
- Williams, K.C., 1992. Some factors affecting albumen quality with particular reference to Haugh unit score. *World's Poult. Sci. J.*, 48: 5-16.
- Yoruk, M.A., M. Gul, A. Hayirli and M. Karaoglu, 2004. Laying performance and egg quality of hens supplemented with sodium bicarbonate during the late laying period. *Int. J. Poult. Sci.*, 3: 272-278.
- Zaghini, A., G. Martelli, P. Roncada, M. Simioli and L. Rizzi, 2005. Mannan oligosaccharides and aflatoxin B1 in feed for laying hens: Effects on egg quality, aflatoxins B1 and M1 residues in eggs and aflatoxin B1 levels in liver. *Poult. Sci.*, 84: 825-832.
- Zaman, M.A., S. Ahmed and B.C. Sutradhar, 2005. Study on the egg quality of a breed and three crossbreds at various ages under semi scavenging system of management. *Pak. J. Biol. Sci.*, 8: 211-214.