

ISSN 1819-1894

Asian Journal of
Agricultural
Research

Experiment on Sand Incubator: An Alternative Mini-Hatchery Technique for Smallholder Poultry Farmers

Yonatan Kassu and Netsanet Beyero

Department of Animal and Range Science, Wolaita Sodo University, Ethiopia

Corresponding Author: Yonatan Kassu, Department of Animal and Range Science, Wolaita Sodo University, Ethiopia

ABSTRACT

The experiment was conducted to compare and evaluate the hatchability and survival rate of a sand incubator and natural hen incubation technique and to estimate the profitability of the two different incubation techniques. The results of this study indicated that fertility, hatchability and dead in shell there were no significant difference ($p>0.05$) among the treatments. Embryonic mortality, normal DOC, abnormal DOC and chick weight showed significant differences ($p<0.05$) among the treatments. Higher embryonic mortality (17%) in broody hen than (9.2%) in sand incubator was observed. On the first batch, the net return per 100 eggs in sand incubation was 177.83 ETB (\$9.3) which is higher net return/profit from that of the broody hen incubation -441.05 ETB or \$-23.0, which were not profitable for the first batch at 70.80% hatchability. On the 1st year run the net return was estimated and can incurred by sand incubator which make it sustainable in profitability for the smallholder farmers by making high income for the next seven years of production period. Finally in order to improve the performance of the hatchery traits before the technology adopt to the farmers intensive adoptive research should conduct and also training of smallholder poultry producers in the hatchery management of sand incubator is very crucial.

Key words: Sand incubator, broody hen, hatchability, net return

INTRODUCTION

Poultry production is an area of animal agriculture, where human food production is relatively fast, initial capital investment is low and use can be made of available household labour. In a developing country like Ethiopia, poultry production in rural areas is of great importance as a prime supplier of eggs and meat and as a source of income, especially to women (Demeke, 1996). Indigenous poultry contribute almost 99% of the national egg and poultry meat production (Tadelle *et al.*, 2003). However due to the low productivity (poor egg production performance, slow growth rate and late sexual maturity) and broodiness for an extended period limit the advantage obtain from this sector. Past attempts that have been made in the country to improve the production and productivity of local birds through upgrading (distribution of exotic birds) and cross breeding (Tadelle and Peters, 2003). The exotic breeds generally do have high productivity in terms of eggs and meat production performance. However, the limit for these breeds is that they did not hatch and produce day old chicken by their own due to lack of broadness and farmers forced to use the local broody hens, which are not enough to meet the actual demand of farmers and producers who want to carry out small-scale operations. A broody hen can hardly hatch 10 eggs in good management practice (Roy *et al.*, 2004).

The much known techniques to hatch improved chicken breed eggs is using modern electric incubator. Modern electric incubators are equipped with automatic devices and generally used by the commercial hatcheries for hatching chicken eggs to meet the huge demand for quality chicks all over the world particularly in the developed countries. This context is not applied in a rural part of many developing country where there is no or/intermittent electric supply and the price of modern incubator not easily afforded by the rural community and small-scale operation enterprises.

Mini-hatcheries can be constructed from cheap materials available locally, such as rice husk, quilts and sand, to retain heat. The incubators can be easily made using readily available skills and tools. Sand-based mini-hatcheries give the highest hatching percentages for both chicken and duck eggs 80-85 and 70-72%, respectively, compared with 70-75 and 65-68% for rice husk incubators and 75-80 and 60-62% for rice husk and quilt incubators (Rota *et al.*, 2011).

Mini-hatcheries are an incubator which is made from locally available materials to incubate chicken or duck eggs. Mini-hatchery incubators have been used to hatch chicken and duck eggs in Egypt and China and Bangladesh. However, the system was not widely adopted by most of developing countries (Hassan, 2011). Surprisingly little is known about the efficiency of mini-hatchery like sand incubation and rice husk incubation.

In Ethiopia improved chicken breed which have been distributed through packages to farmers for the purpose of improving the production status of local chicken and also to generate income by selling their products but the farmers do not have any kind of mini hatchery technology that could hatch a number of chicks other than use of local chicken to hatch egg, because the improved chicken do not incubate and hatch their egg due to low maternity instinct and broodiness. Despite increasing demand for meat, egg and qualities chicks, the mini-hatchery operation in the country is not well known and organized. Therefore, it has become necessary to find a method of egg incubation that is less risky and less expensive, which are affordable and be adopted equally by all sections of urban and rural people through better utilization of existing resources in the country. Therefore, the current study is aimed to evaluate and estimate the suitability, performance and profitability of the sand incubator technique by comparing with the local broody hen hatching technique.

MATERIALS AND METHODS

Precinct of study site: The experiment was conducted in Wolaita Sodo University. The University is located in the mid highlands of Southern Ethiopia at 330 km South of Addis Ababa at about 6049' N latitude and 37045' E longitudes and the farm is situated altitude of 1710 meter above sea level (m.a.s.l). The average mean maximum and minimum temperature are 26.8 and 11.4°C, respectively and the mean maximum and minimum relative humidity is 91.4 and 39.92%, respectively. The mean annual rainfall of the area is 1212 mm (NMSA., 2011).

Experimental animal and materials

Hatching eggs: A total of 747 fertile eggs from the breeds of White leghorn were purchased from Hawassa poultry breeding and distribution center. The hatching eggs were selected for medium and uniform in size with average weight of 54±1.6 g which also have strong shell. The eggs were stored in a room at 14-16°C and 70-80% relative humidity for 4 days. Before setting, all eggs were properly clean with mild luke-warm water solution of savlon (a mild disinfectant). Shape index of hatch able eggs were measured.

Selection and collection of broody hens: Twenty one broody hens with the same age and body size (852 ± 31 g) were purchased from local market. The broody hens were divided randomly into three replications each replication had 7 broody hens for incubation of seven hatching eggs. Before incubation, all broody hens were treated to remove both internal and external parasites. A complete layer grower ration containing 12.9 MJ kg^{-1} and 16.4% CP and fresh drinking water was provided as *ad-libitum*.

Sand incubation techniques: A sand incubator was modified and produced wood workshop which was constructed with materials such as wood, metal plates, jute sacks, black garment, cotton and sand. Before and after setting the eggs, the incubator was as thoroughly fumigated by using potassium permanganate and formaldehyde as per recommendations. The eggs were sat in the setting trays of the incubator with the small end down and the large end up at the optimum conditions of temperature ($37\text{-}39^\circ\text{C}$), relative humidity (60%) and frequent ventilation were maintained during the whole period of incubation using the topside opening and sometimes front door opening. Turning of eggs was done by rolling with a clean hand with gentle circular motion 6 times in 24 h at 4 h interval up to 18 days of incubation period. In both treatments, the candling of eggs was done by using torch-light on the 7th and 14th day of incubation. All clear and early dead embryos were removed after each candling (Fig. 1).

Data collection: Data was collected for fertility, embryonic mortality, dead-in-shells, abnormal chicks hatched, normal chicks hatched, day-old chick weight, survival of day-old chick, hatchability were considered in both methods of incubation techniques. All hatched chicks were weighed individually on a treatment basis with by using sensitive balance with 0.5 g accuracy.

Statistical analysis: All the data collected was subjected to statistical analysis of variance using the PROC GLM of statistically analysis software SAS 9.2 version (SAS Institute Inc., 2008). For the means those had a significant different, Least Significant Difference (LSD) procedure was used. Total estimation of cost and return for each incubation techniques budget analysis was made for each incubation techniques.




Fig. 1(a-b): (a) Hatching eggs and (b) Sand incubator preparation

RESULTS AND DISCUSSION

Hatchability trait performance: Number of egg set per batch, percentage of embryonic mortality, dead in shell, hatchability, normal/abnormal Day Old Chick (DOC), chick weight and survival rate are shown in Table 1. The number of egg set for sand incubator and broody hen is different due to the different holding capacity of eggs for broody hen and sand incubator.

The hatchability of fertile eggs had no significant different ($p > 0.05$) between the two incubation techniques. But numerically higher hatchability percentage was recorded from broody hen (70.8%) than the sand incubation (61.8%). Dolberg *et al.* (2002) and Sonaiya and Swan (2004) reported that the hatching rates of the mini-hatchery technique were in the range 50-70%, which is similar to the current finding (62%), which is very satisfactory, considering that in most of the mini-hatcheries operated with sand and kerosene fuel. In spite of the satisfactory hatchability rate from sand incubator the hatchability rate declines we goes up from the source of heat distributor (sand tray) to the top (2nd and 3rd) tray. This is due to ununiformed heat distribution through the incubator box which was at arrange of 36.5-39°C from top tray to the bottom, respectively (Fig. 2).

The embryonic mortality of broody hen is significantly ($p < 0.05$) higher than that of the sand incubator. Embryonic mortality caused by inadequate ventilation, heat, humidity and vice versa (Dafwang *et al.*, 2005). Broody hen leaves their nest to search for feed and water at this time it might be a fluctuation in temperature which lead to embryo mortality especially after the 13th day of incubation when the embryonic development become faster. Even if the percentage of the dead

Table 1: Hatchability traits of eggs incubated under different methods of incubation

Hatchability traits	Method of incubation		p-value	Sig.
	Broody hen	Sand incubation		
No. of eggs set per batch	147	600		
Fertile egg (%)	85.80±0.6227	87.3000±1.4530	0.406	NS
Embryonic mortality (%)	17.00±1.2282	9.1566±1.4183	0.014	*
Hatchability from total eggs (%)	60.43±1.1259	54.1700±5.8902	0.355	NS
Hatchability from fertile egg (%)	70.80±1.0800	61.8000±6.1941	0.205	NS
Dead in shell (%)	12.20±1.5716	29.0330±5.9839	0.052	NS
Normal DOC (%)	91.53±1.3667	96.8330±0.2603	0.019	*
Abnormal DOC (%)	8.47±1.3667	3.1667±0.2603	0.019	*
Chick weight (g)	35.20±0.4932	37.1933±0.4233	0.037	*
Survival rate (%)	50.77±7.2549	98.1400±0.9567	0.003	**

Values indicate Mean±SE, values within rows without superscript in common differ at $p < 0.05$, ***Significant differences at 5% ($p < 0.05$) and 1% ($p < 0.01$) level of significant, respectively, NS: Non-significant, DOC: Day old chick



Fig. 2(a-b): Chicks that hatched on sand incubator

in shell of broody hen incubation (12%) and sand incubation technique (29%) was not significantly different ($p>0.05$), the sand incubation technique had numerically higher percentage of dead-in-shell. However, in our case to maintain humidity in sand incubator was very difficult due to lack of appropriate hatchery room and provinces humidity condition may cause a severe increase in the proportion of embryo dead-in-shell. Broody hens are experienced enough to maintain optimum conditions of temperature and humidity. Roy *et al.* (2004) reported that dead-in-shell and embryonic mortality rates in the broody hen 2 and 4.80%, respectively. El-Ayadi (1956) reported that White Leghorn eggs dead-in-shell rates were 20.0%.

As shown in Table 1 that, the percentage of normal and abnormal day old chicks hatch had a significant difference ($p<0.05$) among the two incubation techniques. Abnormal chicks are produced, could be due to poor management of the parent flock, environmental conditions in the hatchery room and some genetical/breed factors may contribute for the abnormal chicks.

The implication from the findings of the present study is that the methods of incubation had significant effect on abnormal chicks hatched. The treatments also had influence on normal chicks hatched. Number of sound chicks is an indication of success of hatchability. Day old chicks weight found to be significantly higher than broody hen incubation technique ($p<0.05$). This difference might be due to inadequate supplementation of heat during the incubation period which result poor/retarded embryonic growth/development. The current study result is disagree with Roy *et al.* (2004) which indicate that the average chick weight for broody hen incubation and rice husk incubation were 27 g, which is lower than the current study; this could be the difference in chick strain most of the time the higher breed do have high chick weight whereas lower chicken breed do have lower chick weight.

The survival rate of the hatched chick for one week for broody hen incubation technique (50.7%) was significantly lower ($p<0.01$) than sand incubation technique (98.1%) as shown in Table 1. Among the indigenous local fowls, percentage of the chicks hatched died before they reached cockerel or pullet stage and annually percentage of the entire flock die, before the keeper could reap any benefits in terms of meat or eggs. The difference in survival rate between the two incubation techniques were due to the difference in the management practice. The sand incubated chicks were kept in hover, which they were supplemented by kerosene lamp shown a better survival rate whereas, the broody hen are the only source of heat for the chicks (Fig. 3).



Fig. 3(a-b): Day old chicks after transformed in to their Solomon brooder cage run

Cost and Return of the incubation techniques: The costs and returns of the two different incubation techniques are shown in Table 2 and 3. The total fixed cost per batch for sand incubation was 5255 ETB (\$273.7), which is about 72.60% of the gross cost and from which 66.60% of the fixed cost was spent for the production of the incubator.

From the study, the cost of production using broody hen was very less than that of sand incubation technique. The broody hen technique cost makes 94.5 and 50.3% of the total fixed cost and gross cost respectively. Natural incubation technique is attribute of the broody hen no further heating material are required rather than straw because the hen itself provide all the heat required by the egg in order to develop and hatch.

The total variable cost of sand incubation was higher than the broody hen incubation method, this is due to the difference in capacity and material difference in the two incubation techniques. Although, kerosene lamp were used as a source of heat supplementation for sand incubator in order to hatch a chick and after they transferred in to brooder house up to 2nd weeks of their age which replace the hen function in natural incubation technique. In this experiment, the gross return derived from the return of day-old-chick sale and unfertile egg sale. The gross return per batch attributed by sand incubator was four times higher than the natural incubation technique (Table 3). A well designed sand incubator with 800 eggs capacity can be used/serve as up to eight years with 10 successful batch per year. Whereas, a single hen only incubate an egg for two subsequent production years and can incubate, hatch and brood a chick three times per year.

In the first batch of incubation, which was including all fixed costs, the gross return per 100 eggs in sand incubation method was greater return from that of the broody hen incubation. While, the net return per 100 eggs in sand incubation was 177.83 ETB \$9.3, which is higher return from that of the broody hen incubation -441.05 ETB or \$-23.0, which were not profitable for the first batch at 70.80% hatchability. After the 1st batch of incubation, the net returns were increasing and sustained in both of incubation techniques this is due to the production costs not account the fixed costs. On the 1st year run, the net return was estimated and can incurred by sand incubator which make it sustainable in profitability for the smallholder farmers by making high income for the next seven years of production period. However, the hatchability

Table 2: Fixed and variable costs for hatcheries in two different incubation techniques

Sand incubator			Broody hen		
Parameters	Price		Parameters	Price	
	ETB	USD		ETB	USD
Fixed cost					
Sand incubator	3500.00	182.30	Hen	1470.00	76.60
Lamp	225.00	11.70	House rent	50.00	2.60
Jute sacks	90.00	4.70	Torch	35.00	1.80
Cloth	225.00	11.70			
Thermometer	180.00	9.40			
Hygrometer	950.00	49.50			
House rent	50.00	2.60			
Torch (candling)	35.00	1.80			
Total fixed cost	5255.00	273.70	Total fixed cost	1555.00	81.00
Variable cost					
Egg	1200.00	62.50	Egg	294.00	15.30
Kerosene	118.00	6.10	Feed	461.60	24.00
Disinfectant	140.00	7.30	Medicine	90.00	4.70
Labor	525.00	27.30	Labor	525.00	27.30
Total variable total	1983.00	103.30	Total variable cost	1370.60	71.40

USD: 19.20 ETB by the time of the study, USD: United states dollar , ETB: Ethiopian birr

of the sand incubator can be increase through time with improvement in managing before and after the egg incubation process/operation and on the availability of quality/fertile eggs which domino effect boosts the profitability the smallholder farmers through this technology.

CONCLUSION

Hatchability percentage were lower in sand incubation method than that of the natural incubation however, it is operated with kerosene fuel. Although, the sand incubation techniques from the point of number and cost of the incubator it could hatch a number of chicks per batch (600 eggs) and profitability was higher than that of broody hen incubation technique. In the first year run period earning \$3,339 amount money will be a significant addition to the income of the smallholder farmers.

Therefore, intensive adoptive research should be conduct in different agro climatic condition and seasons and training should be given to smallholder poultry producers on the hatchery management of sand incubator before the technology adopt to the farmers is very crucial and also to make it incentive for the farmers and to explore the profitability of the sand incubator techniques there is need to better DOC (day old Chick) price, low cost of hatching egg, technical support, built-in linkages between producers of fertile eggs and the mini-hatcheries and from the mini-hatcheries to DOCs rears are also necessary to assist farmers to improve their productivity and thus increasing households' income.

ACKNOWLEDGMENTS

The researchers gratefully acknowledge Wolaita Sodo University for providing a research grant for this research. We wish to express our special gratitude to Abay Shimelash for his technical support and Professor Solomon Demeke from Jimma University for their advice and encouragement. We are highly indebted to all staff members of Hawassa poultry breeding and distribution centre for their cooperation.

REFERENCES

- Dafwang, I.I., J.Y. Odiba and E.I. Ikani, 2005. Hatchery management practices in poultry. Extension Bulletin No. 86, Poultry Series No. 5, National Agricultural Extension and Research Liaison Services Ahmadu Bello University, Zaria.
- Demeke, S., 1996. Study on the egg production of White Leghorn under intensive, semi-intensive and rural household conditions in Ethiopia. *Livest. Res. Rural Dev.*, Vol. 8.
- Dolberg, F., E. Mallorie and N. Brett, 2002. Evolution of poultry model-a pathway out of poverty. *Proceeding of the Workshop on People Fight with Poultry: Learning from Bangladesh Experience*, October 20-24, 2002, Dhaka, pp: 18-21.
- El-Ayadi, M.N., 1956. Fertility and hatchability as related to quality in Baladi commercial eggs. M.Sc. Thesis, University of Alexandria, U.A.R.
- Hassan, J., 2011. Mini-hatchery: A low cost hatching technology. *Network for the Veterinarians of Bangladesh*, pp: 1-2.
- NMSA., 2011. Monthly report on temperature and rainfall distribution for Eastern Ethiopia metrological office. Addis Ababa, Ethiopia.
- Rota, A., N. Brett, J. Nahar, S.M.R. Rahman, Y. Ali, A. Sarwar and K.A. Fattah, 2011. Building and operating a mini-hatchery (Sand method)-video. *International Fund for Agricultural Development (IFAD)*.

- Roy, B.C., H. Ranvig, S.D. Chowdhury, M.M. Rashid and M.R. Faruque, 2004. Production of day-old chicks from crossbred chicken eggs by broody hens, rice husk incubator and electric incubator and their rearing up to 6 weeks. *Livest. Res. Rural Dev.*, Vol. 16.
- SAS Institute Inc., 2008. *Statistical Analysis Software Version 9.2*. SAS Institute Inc., Cary, North Carolina, USA.
- Sonaiya, E.B. and S.E.J. Swan, 2004. *Manual on Small-scale Poultry Production*. FAO., Rome.
- Tadelle, D. and K.J. Peters, 2003. *Indigenous chicken in Ethiopia: Their genetic potential, attempts made in the past for improvement and future areas of research*. *Animal Breeding for Tropics and Sub-tropics*, University of Berlin, Humboldt.
- Tadelle, D., T. Million, Y. Alemu and K.J. Peters, 2003. *Village chicken production systems in Ethiopia: 2. Use patterns and performance valuation and chicken products and socio-economic functions of chicken*. *Livestock Res. Rural Dev.*, Vol. 15.