

ISSN 1996-3351

Asian Journal of
Biological
Sciences



Research Article

Histological and Ultrastructure Alterations in the Midgut of *Blaps polycresta* and *Trachyderma hispida* (Coleoptera: Tenebrionidae) Induced by Heavy Metals Pollution

Dalia Abdel Moneim Kheirallah and Lamia Mostafa El-Samad

Department of Zoology, Faculty of Science, Alexandria University, 21511 Alexandria, Egypt

Abstract

Background and Objective: The present research focused on using two coleopteran insects *Blaps polycresta* and *Trachyderma hispida* as bioindicators for heavy metals pollution that resulted from ceramic industry in an urban area, Khorshed district, Alexandria, Egypt. **Materials and Methods:** Bioaccumulation of the heavy metals was quantified in the soil and midgut tissues of the two insects by using x-ray microprobe analysis. Histological and ultrastructure preparations were performed at the Electron Microscope Unit at Faculty of Science, Alexandria University, Egypt. **Results:** The x-ray analysis revealed high metal percentage in the soil and midgut tissues of the insects collected from the polluted site compared to the reference site (the garden of Faculty of Science, Alexandria University, Egypt). Histological and ultrastructure alterations were observed in the midgut cells of the two insects collected from the polluted site compared to the reference site. The most observed histological alteration was the distortion of the brush borders of the microvilli. Ultrastructure alterations included: Nuclear distortion of regenerative and digestive cells, lysis of the mitochondrial matrices, the appearance of electron-dense vesicles, the presence of myelin figures, vacuolated cytoplasm and dilation of the rough and smooth endoplasmic reticulum. **Conclusion:** The study validates that heavy metals pollution leads to cellular and sub-cellular alterations in insects inhabiting industrial districts.

Key words: Industrial pollution, heavy metals, beetles, histological and ultrastructure aberrations, x-ray analysis

Citation: Dalia Abdel Moneim Kheirallah and Lamia Mostafa El-Samad, 2019. Histological and ultrastructure alterations in the midgut of *Blaps polycresta* and *Trachyderma hispida* (coleoptera: tenebrionidae) induced by heavy metals pollution. Asian J. Biol. Sci., 12: 637-647.

Corresponding Author: Dalia Abdel Moneim Kheirallah, Department of Zoology, Faculty of Science, Alexandria University, 21511 Alexandria, Egypt
Tel: 002-01221775286 Fax: 002-034264455

Copyright: © 2019 Dalia Abdel Moneim Kheirallah and Lamia Mostafa El-Samad. This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Industrialization poses serious risks to human and ecosystems' health. Populations in developing countries are especially at risk because of the lacking of appropriate regulations and enforcement as well as barriers to accessing "cleaner" technologies. The undesirable outcomes from industrial plants such as the emission of toxic chemicals in the atmosphere resulted in environmental pollution¹⁻³. Ceramic industry is an important source of pollutants to the environment⁴. Many relevant aspects such as air pollution, human toxicity, global warming, acidic rain and ozone depletion from ceramic tile processes have been discussed by several authors^{2,3,5-11}. Residential plants and animals were found to be affected in ceramic industrial areas¹². Because of the usage of many chemicals and high energy which is essential in this industry soil pollution occurs¹³.

The chemicals that have been used in manufacturing ceramic contains heavy metals which are the primary cause of ecotoxicity¹¹. Variable fractions of the total concentration of the heavy metals are bioavailable to the soil animals¹⁴. The more accurate information is the measurements of the heavy metals concentrations within the inhabitants¹⁵. Heavy metals reside in the organisms' tissues and may pose health problems. Xenobiotic undergo a series of biotransformation

reactions catalyzed by different enzymes, their activation provides evidence for pollution exposure¹⁶.

Insects are good monitoring models to evaluate environmental alterations and heavy metals pollution¹⁷⁻²². The midgut in insects is directly subjected to pollutants as it acts as a physical and chemical barrier against the ingested toxic matters while feeding²³⁻²⁵. Histological and ultrastructure biomarkers deliver good surveillance for environmental health. Hence, the focal point of this survey is reporting structural abnormalities at the cellular and subcellular levels in the midgut of *B. polycresta* and *T. hispida* as biomarkers of ceramic industrial pollution and heavy metals toxicity. These anomalies will serve in the identification of different health risks that resulted from exposure to environmental stresses.

MATERIALS AND METHODS

Study sites: The two selected sites in the study were a rustic area, the garden of Faculty of Science Moharam Bek, Alexandria University, Alexandria, Egypt which considered as reference site (site A)¹⁸ and a densely populated area, Khorshed district which located in the suburbs of the eastern edge of Alexandria, Egypt (Latitude: 31.2018066, Longitude: 30.0300721), the polluted site (site B) (Fig. 1). Ceramic factories allocated through Khorshed district (Lecico and Khorshed ceramic factories).



Fig. 1: Map of Alexandria Governorate showing Khorshed district. Arrow pointed at ceramic factories. [https://www.google.com/maps/Alexandria, Egypt, Khorshed \(31.2018066, 30.0300721\)](https://www.google.com/maps/Alexandria, Egypt, Khorshed (31.2018066, 30.0300721))

Specimens identification: The coleopteran species *Blaps polycresta* and *Trachyderma hispida* were the superior beetles populated in the nominated sites. Beetles' identification was performed at Entomology Department, Faculty of Agriculture, Alexandria University. They belong to Coleoptera: Tenebrionidae.

Sampling procedure: The beetles, *B. polycresta* and *T. hispida* were collected randomly from ten sampling areas (1 m² each) at each site in August, 2018. Synchronously with the beetle collection, soil samples at a depth of 30 cm below the surface were gathered from the specific sites. The ten areas in Khorshed district were selected around the ceramic factories (five sites around each factory). Air temperature in July ranged from 27-35 °C and the mean relative humidity was 76% with nearly no differences between the two sites. About 100 beetles from each species were collected from each site. The specimens were sustained alive in local soil and plants in glass containers until processing. Beetles were anesthetized with absolute ethanol (95%), then dissected under a dissecting microscope in a drop of Ringer's physiological solution. The abdominal cavity was opened and the midgut was taken out. Maintenance of the insects was done in compliance with ethical guidelines for the protection and use of laboratory animals. The methodology was approved by the Ethics Committee of Alexandria University (protocol approval number is 0302440).

Bioaccumulation of metals in the soil and in the midgut tissues: Metals accumulation in sieved soil and in un-coated specimens of the midgut were detected using Jeol scanning electron microscope-5300 equipped with a Link-Isis energy dispersive X-ray micro-analyzer. A stationary spot (X500) was analyzed at random for 110 sec. Due to the divergent distribution of trace metals, four points were measured to check for the variability of trace metal composition.

Histological and ultrastructure investigations: The histological preparation followed Anderson and Gordon²⁶ methods of dehydration, clearing and paraffin embedding. Xylene was the clearing agent. Midgut was fixed in paraffin wax (65-60 °C) and 5 µm thick sections were stained with hematoxylin and eosin.

In ultrastructure preparation, midgut was fixed in OsO_4 in phosphate buffer solution (pH 7.2) at 4 °C for 3 h and post-fixed in 2% OsO_4 in the same buffer for 2 h. Samples were washed in the buffer and dehydrated at 4 °C through a series of ethanol. Specimens were immersed in Epon-Araldite mixture in labeled beam capsules. For semithin sections, LKB ultramicrotome was used (1 µm thick). Sections were mounted

on a glass slide, stained with toluidine blue and examined with the light microscope to determine the orientation and the structural features. Ultrathin sections (0.06-0.07 µm thick)²⁷ were cut for TEM and picked upon 200 mesh naked copper grids. Grids were stained with uranyl for half an hour and lead citrate for 20-30 min.

Statistical analysis: Data analysis was performed using the program SPSS²⁸. Data were analyzed statistically by using t-test²⁹ to determine the difference between the two studied sites.

RESULTS

X-ray microanalysis: Twelve elements were represented in the x-ray spectra including Na, Al, P, S, K, Pb, Cd, Ca, Fe, Ni, Cu and Zn. The soil samples and the midgut tissues of the two beetles collected from the polluted site showed a significant elevation in the percentages of Al, Cu and Zn with detection of Cd and Pb compared to the reference site (Table 1, 2). Proportions of Na, P, S, K and Fe were higher in the reference site than the polluted site.

Histological and ultrastructure archetypes observed in the midgut of *B. polycresta* and *T. hispida* collected from the reference site (site A)

Histological observations: The epithelial sheath is covered by two muscle layers, external longitudinal and internal circular (Fig. 2a, b, 3a). The epithelium composed of two types of cells: (1) The columnar digestive cells with a brush border of microvilli on the apical membrane, facing the peritrophic membrane and the gut lumen and (2) The regenerative cells occur singly or in small groups called "nidi", forming crypts, in contact with the basal lamina (Fig. 2a-c, 3a, b).

Table 1: Metal percentages in the soil collected from either reference or polluted site using energy dispersive X-ray micro-analysis (EDX)

Metals	Site A	Site B	p-value
Na	9.5±0.06	7.4±0.06	<0.001*
Al	2.0±0.06	8.5±0.06	<0.001*
P	37.1±0.06	41.2±0.06	<0.001*
S	35.7±0.06	11.6±0.06	<0.001*
K	1.2±0.06	ND	-
Pb	ND	7.3±0.06	-
Cd	ND	11.6±0.06	-
Ca	2.7±0.06	1.4±0.06	<0.001*
Fe	ND	1.2±0.06	-
Cu	1.9±0.06	16.9±0.06	<0.001*
Zn	1.1±0.06	13.8±0.06	<0.001*

For each metal, the percentage expressed by using minimum-maximum values and mean (n = 3) using Student t-test, p: p-value for comparing between site A and site B, *: Statistically significant at p≤0.05, ND: Not detected




Fig. 2(a-c): Semithin sections of the midgut tissues of *B. polycrsta* collected from the reference site (site A) (a) Epithelial sheath (Ep), longitudinal muscle (double head arrow), circular muscle (arrow), regenerative cells (RC) forming crypts (curved arrow), columnar digestive cells (CC), microvilli (Mv), peritrophic membrane (arrow head) evolving food particles (F). (b, c) Magnified parts of Fig. 2a

Table 2: Metal percentages in midgut tissues of *B. polycrsta* and *T. hispida* collected from either reference or polluted sites using energy dispersive X-ray micro-analysis (EDX)

Metals	<i>B. polycrsta</i>			<i>T. hispida</i>		
	Site A	Site B	p-value	Site A	Site B	p-value
Na	11.3±0.06	7.1±0.06	<0.001*	14.6±0.06	7.4±0.06	<0.001*
Al	5.8±0.06	11.8±0.06	<0.001*	6.9±0.06	16.5±0.06	<0.001*
P	26.2±0.06	43.4±0.06	<0.001*	26.5±0.06	41.1±0.06	<0.001*
S	34.2±0.06	26.5±0.06	<0.001*	31.4±0.06	20.4±0.06	<0.001*
K	8.0±0.06	ND	-	ND	ND	-
Pb	ND	4.3±0.06	-	ND	4.1±0.06	-
Cd	ND	7.6±0.06	-	ND	12.3±0.06	-
Ca	ND	0.3±0.06	-	4.9±0.06	ND	-
Fe	0.5±0.06	0.8±0.06	0.021*	0.3±0.06	0.2±0.06	0.288
Cu	2.9±0.06	7.9±0.06	<0.001*	3.4±0.06	6.7±0.06	<0.001*
Zn	2.1±0.06	4.2±0.06	<0.001*	2.6±0.06	5.47±0.03	<0.001*

For each metal, the percentage expressed by using minimum-maximum values and mean (n = 3) using Student t-test, p: p-value for comparing between site A and site B, *Statistically significant at p<0.05, ND: Not detected

Ultrastructure observations: The regenerative cells appeared with an oval nucleus, patches of heterochromatin and regular

nuclear envelope (Fig. 4a, 5a, b). In the cytoplasm, uniformly distributed rounded mitochondria, frequent cisterns of rough endoplasmic reticulum (RER) and free ribosomes were noticed (Fig. 4a, 5a, b). Smooth septate junctions were found between adjacent epithelial cells (desmosomes) (Fig. 4a, b, 5a, b). The columnar digestive cells appeared with an oval nucleus and regular nuclear envelope (Fig. 5c). The cytoplasm possesses numerous mitochondria, cisterns of the rough and smooth endoplasmic reticulum, glycogen granules and free ribosomes (Fig. 4b, 5c). The luminal border has a uniformly distributed microvilli (Fig. 4b, 5c).

Histological and ultrastructure archetypes observed in the midgut of *B. polycrsta* and *T. hispida* collected from the polluted site (site B)

Histological observations: Incisions between adjacent cells (Fig. 6a), vacuolation (Fig. 6a-c, 7a-c) and destruction of the brush border of the microvilli (Fig. 6c, 7c) were observed.




Fig. 3(a-b): Semithin sections of the midgut tissues of *T. hispida* collected from the reference site (site A) (a) Epithelial sheath (Ep), muscle layers: Longitudinal (double head arrow) and circular (arrow), regenerative cells (RC), columnar digestive cells (CC), microvilli (MV) and (b) Magnified part of Fig. 3a




Fig. 4(a-b): Electron micrographs midgut cells of *B. polycresta* collected from the reference site (site A) (a) Nucleus (N), nuclear envelope (Ne), chromatin (CH), mitochondria (M), rough endoplasmic reticulum (RER), free ribosomes (r), desmosomes (arrow head) and (b) Basal region of the digestive cells with numerous mitochondria (M), glycogen granules (GL) and rough endoplasmic reticulum (RER), luminal border with microvilli (MV), desmosomes (arrow head)

Ultrastructure observations: Signs of necrosis were observed in the nuclei of some regenerative cells which included: Formation of blebs (Fig. 8a, b, 9a) and pseudo-inclusions at the nuclear envelopes (Fig. 8c, 9a), globular inclusions in the nucleoplasm (Fig. 8a, b, 9a) and Karyorrhexis (Fig. 8a, 9a). Abnormal chromatin clumps were also observed in the nucleoplasm (Fig. 8a, b). Apoptotic regenerative and digestive cells were noticed (Fig. 8c, e).

Mitochondria appeared with lysed matrices (Fig. 8a, b). Lysosomes (Fig. 8b, c, e, 9a, b), electron-dense vesicles (Fig. 8d, e, 9b), myelin figures (Fig. 8c, e, 9b) and cytoplasmic vacuolation (Fig. 8a-e, 9a, b) were frequently observed. In addition, dilated rough and smooth endoplasmic reticulum (Fig. 8c) were remarked. Microvilli appeared with distorted brush borders (Fig. 8d, e, 9b).




Fig. 5(a-c): Electron micrographs of midgut cells of *T. hispida* collected from the reference site (site A) (a-b) Nucleus of regenerative cells (N), regular nuclear envelope (Ne), chromatin (CH), mitochondria (M), rough endoplasmic reticulum (RER), free ribosomes (r), desmosomes (arrow head). (c) Digestive cells with nucleus (N) regular nuclear envelope (Ne), mitochondria (M), rough endoplasmic reticulum (RER), smooth endoplasmic reticulum (SER), microvilli (MV), glycogen granules (GL), lysosomes (L)




Fig. 6(a-c): Semithin sections of the midgut tissues of *B. polycresta* collected from the polluted site (site B) (a-b) Lacerated (*) and vacuolated (V) epithelial sheath (Ep), (c) Vacuolated (V) epithelial sheath (Ep), distorted microvilli (MV) (curved arrow). Muscle layers: Longitudinal (double head arrow) and circular (arrow), regenerative cells (RC), columnar digestive cells (CC), peritrophic membrane (arrow head), food particles (F)




Fig. 7(a-c): Semithin sections of the midgut tissues of *T. hispida* collected from the polluted site (site B) (a) Vacuolated (V) epithelial sheath (Ep), (b-c) Vacuolated (V) epithelial sheath (Ep), distorted microvilli (MV) (arrow), regenerative cells (RC), columnar digestive cells (CC), peritrophic membrane (arrow head)




Fig. 8(a-e): Electron micrographs of abnormal midgut cells of *B. polycrsta* collected from the polluted site (site B) (a) Nucleus (N), abnormal chromatin clumps (CH), intended nuclear envelope (Ne), lysis of mitochondrial matrices (arrow), vacuoles (V), (b) Globular inclusions (arrow). Karyorrhexis (double head arrow), vacuoles (V), lysosomal vesicles (L), (c) Pseudo-inclusions (curved arrow), globular inclusions (arrow), lysed mitochondria (M), vacuoles (V), lysosomes (L), myelin figure (double head arrow), dilated rough (RER) and smooth endoplasmic reticulum (SER), apoptotic cell (*), (d) Distorted microvilli (arrow), vacuoles (V), electron dense vesicles (curved arrow), glycogen granules (GL) and (e) Karyolysed nucleus (N), myelin figure (double head arrow), vacuoles (V), lysosomes (L), glycogen granules (GL), free ribosomes®




Fig. 9(a-b): Electron micrographs of abnormal midgut cells of *T. hispidula* collected from the polluted site (site B) (a) Nucleus (N), chromatin (CH), irregular nuclear envelope (Ne), pseudo-inclusions and globular inclusions (arrow), Karyorrhexis (curved arrow), secretory vesicle (SV), lysosomes (L), rough endoplasmic reticulum (RER), microvilli (MV) and (b) Distorted microvilli (arrow), myelin figure (double head arrow), vacuoles (V), lysosomes (L), dense vesicles (curved arrow), mitochondria (M), glycogen granules (GL)

DISCUSSION

The current study illustrated the cellular anomalies that resulted from ceramic industrial pollution in two bioindicator beetles, *B. polycresta* and *T. hispidula*. Ceramic industry has been retreated from most of the countries and progressed in the developing countries. The problem is the management of the pollution control planet. Factories are required to eliminate the operational defects¹⁰. The chemicals that are contributed to the manufacturing process contain heavy metals that are released in the environment and threat the biota¹³. Many authors have reported the prominent act of insects as bioindicators^{18,20-22,25,30,31}. The x-ray microanalysis in the present study proclaimed significant high percentages of heavy metals in the soil and midgut tissues of beetles assembled from the polluted site. The midgut in insects is the main site of absorption and plays an essential role in ionic regulation and mineral³².

Ingestion of heavy metals by insects leads to histological and ultrastructure anomalies^{18,20-22,30,31,33}.

The observed alterations in the midgut tissues of the two beetles collected from the polluted site could be ascribed to metals accumulation in these tissues as reported previously by Xie *et al.*³⁴. Several studies recorded structural alteration in the midgut of insects exposed to pollutants^{24,25,35,36}.

A normal architecture of the midgut epithelium cells was noticed in the beetles collected from the reference site. An outer longitudinal and inner circular muscle layers covering the midgut epithelium, designed for slow and super contractions and an absorption surface, the microvilli³⁷.

Ultrastructurally, the nuclei, the rough endoplasmic reticulum and the mitochondria appeared in a healthy manner. Regenerative cells lack lysosomes, glycogen granules and secretory vesicles, they are found only in the digestive cells³⁸. The digestive cells have numerous mitochondria in the apical and basal thirds of the cells to support active transport role³⁸. The rough endoplasmic reticulum was arranged in spirals or vesicles throughout the cytoplasm, contrary to Silva-Olivares *et al.*³⁷. The abundance of the rough endoplasmic reticulum denotes that a large amount of proteins is synthesized in these cells³⁹.

Contrarily, the midgut of the beetles collected from the polluted site exhibited several histological and ultrastructure deformations. The earliest changes involve chromatin clumping suggesting a less efficient transcription⁴⁰. The formation of blebs at the nuclear envelopes, globular inclusions in the nucleoplasm and Karyorrhexis were signs of cell death pathway^{41,42}. Loeb *et al.*⁴³ found that Cd and Pb produce apoptosis in the midgut of lepidopteran larvae *Heliothis virescens*. They observed cellular shrinkage,

chromatin condensation, blebbing of the cytoplasm and DNA fragmentation. Several authors attributed apoptosis to metal accumulation⁴⁴⁻⁴⁷ which are in agreement with the current results. The vacuolated areas in the cytoplasm may be ascribed to the breakage of the mitochondria⁴⁸ or sometimes may be due to the enhanced endocytotic activity⁴⁹. The presence of electron-dense vesicles in the midgut tissues may be attributed to metal accumulations as a detoxification mechanism⁵⁰. The present results are in accordance with Polidori *et al.*⁴⁸ who detected metal precipitation in spherites in the midgut of paper wasps (*Polistes dominula*) collected from Urban environments. Also, a similar observation was noticed by Pigino *et al.*⁵¹, who reported that a large number of electron-dense granules, composed of a variety of heavy metals were accumulated in the epithelium of the midgut ventriculus of the mite *Xenillus tegeocranus* from a deserted mining and smelting area. Disruption of microvilli is one of the major findings in this study since it is the first site facing and interacting with the pollutants^{48,52}. Some changes in the cytoplasmic organelles were distinguished in the electron micrographs such as lysis of mitochondrial matrices, dilated rough and smooth endoplasmic reticulum and presence of myelin figures. It was found that heavy metals distort cytoplasmic membranes^{20,22,30,53}. Mitochondrial alteration is a reflection of the deregulation of mitochondrial membrane transport⁵⁴. Also, the interference of heavy metals with proper processing in the ER causes its dilation and activates the ER stress response⁵⁵. The proliferation of myelin figures in the current preparations has been interpreted as a symptom of intoxication triggered by xenobiotics which implies an adaptive mechanism in response to the high degradation of cellular organelles⁵⁶. Eventually, these results form a pathway of cell injury resulted from industrial pollution and diagnose the toxicological pathology accompanied manufacturing ceramic. Precautions against manufacturing ceramic should be taken to limit the operational defects that may result in a polluted environment.

CONCLUSION

The study validates that the beetles are sensible bioindicators for industrial pollution and reveals the ultrastructural damage to the midgut cells since they are the first sites of the action of heavy metals through ingestion. It also elucidates the threat resulted from ceramic industries in urban areas that may lead to a serious risk to human health.

SIGNIFICANCE STATEMENT

This study discovered the heavy metals toxicity that resulted from ceramic industrial planets by using ground beetles as bioindicators that can be beneficial for the detection of health problems. This study will help the researcher to uncover the critical areas of cell injuries that many researchers were not able to explore. Thus a new theory on "Hazards of ceramic industry" may be arrived at.

ACKNOWLEDGMENT

The authors are thankful to the Zoology Department and Electron microscope unit, Faculty of Science, Alexandria University.

REFERENCES

1. Shen, T.T., 1995. Industrial Pollution Prevention. Springer-Verlag, Berlin, Germany, ISBN-13: 978-3-662-03110-0, Pages: 373.
2. Shular, J., 1996. The emission factor documentation for AP-42, Section 11.7: Ceramic products manufacturing. EPA Contract 68-D2-0159, Work Assignment No. 4603-01, U.S. Environmental Protection Agency, USA., June 1996.
3. Nicoletti, G.M., B. Notarnicola and G. Tassielli, 2002. Comparative life cycle assessment of flooring materials: Ceramic versus marble tiles. J. Cleaner Prod., 10: 283-296.
4. Barros, C., P. Bello, E. Roca and J.J. Casares, 2007. Integrated pollution prevention and control for heavy ceramic industry in Galicia (NW Spain). J. Hazard. Mater., 141: 680-692.
5. Gavioli, G. and G. Timellini, 1981. Le prescrizioni della regione Emilia Romagna per la prevenzione dell'inquinamento atmosferico da industrie ceramiche. La Ceramica, 34: 37-41.
6. Palmonari, C. and G. Timellini, 1982. Pollutant emission factors for the ceramic floor and wall tile industry. J. Air Pollut. Control Assoc., 32: 1095-1100.
7. Palmonari, C. and G. Timellini, 1986. Air pollution from the ceramic industry, pollutant emission factors and chemical-physical characterization of gaseous emissions according to product type and manufacturing technology. Proceedings of the 7th World Clean Air Congress, August 25-19, 1986, Sydney, Australia, pp: 76-84.
8. Palmonari, C. and G. Timellini, 1989. Air pollution from the ceramic industry: Control experiences in the Italian ceramic-tile industry. Am. Ceram. Soc. Bull., 68: 1464-1469.
9. Palmonari, C. and G. Timellini, 1983. Reduction of pollution by new production technologies in the building ceramics industry. Proceedings of the European Symposium on Clean Technologies, November 4-7, 1980, The Hague, The Netherlands, pp: 523-536.

10. Timmellini G., F. Cremonini and C. Palmonari, 1993. Air pollution from ceramic tile processes. *Ceram. Eng. Sci. Proc.*, 14: 445-456.
11. Tikul, N. and P. Srichandr, 2010. Assessing the environmental impact of ceramic tile production in Thailand. *J. Ceram. Soc. Jpn.*, 118: 887-894.
12. Abrahao, R. and M. Carvalho, 2017. Environmental impacts of the red ceramics industry in Northeast Brazil. *Int. J. Emerg. Res. Manage. Technol.*, 6: 310-317.
13. Goldoni, S. and A. Bonoli, 2006. A case study about LCA of ceramic sector: application of life cycle analysis results to the environment management system adopted by the enterprise. University of Bologna, Italy.
14. Van Gestel, C.A.M., 1992. The Influence of Soil Characteristics on the Toxicity of Chemicals for Earthworms: A Review. In: *Ecotoxicology of Earthworms*, Greig-Smith, P.W., H. Becker, P.W. Edwards and F. Heimbach (Eds.). Intercept Ltd. andover, Hampshire, UK., ISBN: 0-946707-40-5, pp: 44-54.
15. Hensbergen, P.J., M.J. van Velzen, R.A. Nugroho, M.H. Donker and N.M. van Straalen, 2000. Metallothionein-bound cadmium in the gut of the insect *Orchesella cincta* (Collembola) in relation to dietary cadmium exposure. *Comp. Biochem. Physiol. Part C: Pharmacol. Toxicol. Endocrinol.*, 125: 17-24.
16. Mormede, S. and I.M. Davies, 2001. Heavy metal concentrations in commercial deep-sea fish from the Rockall Trough. *Continental Shelf Res.*, 21: 899-916.
17. Chen, T.B., Y.M. Zheng, M. Lei, Z.C. Huang and H.T. Wu *et al*, 2005. Assessment of heavy metal pollution in surface soils of urban parks in Beijing, China. *Chemosphere*, 60: 542-551.
18. Osman, W., L.M. El-Samad, E.H. Mokhamer, A. El-Touhamy and M. Shonouda, 2015. Ecological, morphological and histological studies on *Blaps polycresta* (Coleoptera: Tenebrionidae) as biomonitors of cadmium soil pollution. *Environ. Sci. Pollut. Res.*, 22: 14104-14115.
19. El-Moaty, Z.A., D.A. Kheirallah and D.A. Elgendy, 2016. Impact of cement dust on some biological parameters of *Trachyderma hispidula* (Coleoptera: Tenebrionidae) inhabiting the vicinity of a cement factory, Mariout region, Alexandria, Egypt. *J. Entomol. Zool. Stud.*, 4: 797-805.
20. Kheirallah, D.A.M., Z.A. El-Moaty and D.A. El-Gendy, 2016. Impact of cement dust on the testis of *Tachyderma hispidula* (Forsk., 1775) (Coleoptera: Tenebrionidae), inhabiting Mariout region (Alexandria, Egypt). *J. Entomol.*, 13: 55-71.
21. Osman, W. and M. Shonouda, 2017. X-ray metal assessment and ovarian ultrastructure alterations of the beetle, *Blaps polycresta* (Coleoptera, Tenebrionidae), inhabiting polluted soil. *Environ. Sci. Pollut. Res.*, 24: 14867-14876.
22. Shonouda, M. and W. Osman, 2018. Ultrastructural alterations in sperm formation of the beetle, *Blaps polycresta* (Coleoptera: Tenebrionidae) as a biomonitor of heavy metal soil pollution. *Environ. Sci. Pollut. Res.*, 25: 7896-7906.
23. Rawi, S.M., F.A. Bakry and M.A. Al-Hazm, 2011. Biochemical and histopathological effect of formulated and non formulated plant extracts on *Spodoptera littoralis* larvae. *Int. J. Plant Sci.*, 2: 107-118.
24. Decio, P., E.C.M. Silva-Zacarin, F.C. Bueno and O.C. Bueno, 2013. Toxicological and histopathological effects of hydramethylnon on *Atta sexdens rubropilosa* (Hymenoptera: Formicidae) workers. *Micron*, 45: 22-31.
25. Abu El-Saad, A.M., D.A. Kheirallah and L.M. El-Samad, 2017. Biochemical and histological biomarkers in the midgut of *Apis mellifera* from polluted environment at Beheira Governorate, Egypt. *Environ. Sci. Pollut. Res.*, 24: 3181-3193.
26. Anderson, G. and K. Gordon, 1996. *Tissue Processing, Microtomy and Paraffin Sections*. In: *Theory and Practice of Histological Techniques*, Bancroft, J.D. and A. Stevens (Eds.). Churchill Livingstone, New York, USA., ISBN-13: 9780443047602, pp: 47-80.
27. Reynolds, E.S., 1963. The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *J. Cell Biol.*, 17: 208-212.
28. Kirkpatrick, L.A. and B.C. Feeney, 2013. *A Simple Guide to IBM SPSS® Statistics for Version 20.0*. Cengage Learning, Wadsworth, OH., USA., ISBN-13: 9781285086019, Pages: 128.
29. Sokal, R.R. and F.J. Rohlf, 1981. *Biometry: The Principles and Practice of Statistics in Biological Research*. 2nd Edn., W.H. Freeman Publisher, San Francisco, USA., ISBN-13: 9780716712541, Pages: 859.
30. Kheirallah, D.A., 2006. Impact of pollution on the water bug *Sphaerodema urinator* (Dufour, 1833) inhabiting lakes Mariut and Edku. Ph.D. Thesis, Faculty of Science, Alexandria University, Egypt.
31. Kheirallah, D.A., 2015. Ultrastructure biomarker in *Anisops sardus* (Heteroptera: Notonectidae) for the assessment and monitoring of Water Quality of Al-Mahmoudia Canal, Western Part of Nile Delta, Egypt. *J. Biosci. Applied Res.*, 1: 326-334.
32. Ballan-Dufrencais, C., 2002. Localization of metals in cells of pterygote insects. *Microsc. Res. Tech.*, 56: 403-420.
33. Bednarska, A.J., I. Portka, P.E. Kramarz and R. Laskowski, 2009. Combined effect of environmental pollutants (nickel, chlorpyrifos) and temperature on the ground beetle, *Pterostichus oblongopunctatus* (Coleoptera: Carabidae). *Environ. Toxicol. Chem.*, 28: 864-872.
34. Xie, G., J. Zou, L. Zhao, M. Wu, S. Wang, F. Zhang and B. Tang, 2014. Inhibitional effects of metal Zn²⁺ on the reproduction of *Aphis medicaginis* and its predation by *Harmonia axyridis*. *PLoS ONE*, Vol. 9. 10.1371/journal.pone.0087639
35. Khan, I., A. Qamar, S.H. Mehdi and M. Shahid, 2011. Histopathological effects of *Datura alba* leaf extract on the midgut of *Periplaneta americana*. *Biol. Med.*, 3: 260-264.
36. Rawi, S.M., F.A. Bakry and M.A. Al-Hazmi, 2011. Biochemical and histopathological effect of crude extracts on *Spodoptera littoralis* larvae. *J. Evol. Biol. Res.*, 3: 67-78.

37. Silva-Olivares, A., E. Diaz, M. Shibayama, V. Tsutsumi, R. Cisneros and G. Zuniga, 2014. Ultrastructural study of the midgut and hindgut in eight species of the genus *Dendroctonus* Erichson (Coleoptera: Scolytidae). *Ann. Entomol. Soc. Am.*, 96: 883-900.
38. Billingsley, P.F. and M.J. Lehane, 1996. Structure and Ultrastructure of the Insect Midgut. In: *Biology of the Insect Midgut*, Lehane, M.J. and P.F. Billingsley (Eds.). Springer, Netherlands, ISBN: 978-0-412-61670-9, pp: 3-30.
39. Geneser, F., 2000. *Histologia: Sobre Bases Biomoleculares*. 3rd Edn., Medica Panamericana, Mexico, ISBN-13: 9789500608831, Pages: 813.
40. Grewal, S.I. and S. Jia, 2007. Heterochromatin revisited. *Nat. Rev. Genet.*, 8: 35-46.
41. Trump, B.E., I.K. Berezsky, S.H. Chang and P.C. Phelps, 1997. The pathways of cell death: Oncosis, apoptosis and necrosis. *Toxicol. Pathol.*, 25: 82-88.
42. Pazir, M.K., M. Afsharnasab, B.J. Jafari, I. Sharifpour, A.A. Motalebi and A. Dashtiannasab, 2011. Detection and identification of white spot syndrome virus (WSSV) and infectious hypodermal and hematopoietic necrosis virus (IHHNV) of *Litopenaus vannamei* from Bushehr and Sistan and Baloochestan Provinces (Iran), during 2009-2010. *Iran. J. Fish. Sci.*, 10: 708-726.
43. Loeb, M.J., R.S. Hakim, P. Martin, N. Narang, S. Goto and M. Takeda, 2000. Apoptosis in cultured midgut cells from *Heliothis virescens* larvae exposed to various conditions. *Arch. Insect Biochem. Physiol.*, 45: 12-23.
44. Hay, B.A., J.R. Huh and M. Guo, 2004. The genetics of cell death: Approaches, insights and opportunities in *Drosophila*. *Nat. Rev. Genet.*, 5: 911-922.
45. Xia, Q., H. Sun, X. Hu, Y. Shu, D. Gu and G. Zhang, 2005. Apoptosis of *Spodoptera litura* larval hemocytes induced by heavy metal zinc. *Chin. Sci. Bull.*, 50: 2856-2860.
46. Rodrigues, A., L. Cunha, A. Amaral, J. Medeiros and P. Garcia, 2008. Bioavailability of heavy metals and their effects on the midgut cells of a phytophagous insect inhabiting volcanic environments. *Sci. Total Environ.*, 406: 116-122.
47. Braeckman, B.P., 2011. Heavy Metal Toxicity in an Insect Cell Line (Methyl-HgCl, HgCl₂, CdCl₂ and CuSO₄). In: *Cellular Effects of Heavy Metals*, Banfalvi, G. (Ed.). Springer, Dordrecht, The Netherlands, ISBN: 978-94-007-0427-5, pp: 115-144.
48. Polidori, C., A. Pastor, A. Jorge and J. Pertusa, 2018. Ultrastructural alterations of midgut epithelium, but not greater wing fluctuating asymmetry, in paper wasps (*Polistes dominula*) from urban environments. *Microsc. Microanal.*, 24: 183-192.
49. Cavados, C.F.G., S. Majerowicz, J.Q. Chaves, C.J.P.C. Araujo-Coutinho and L. Rabinovitch, 2004. Histopathological and ultrastructural effects of δ -endotoxins of *Bacillus thuringiensis* serovar *israelensis* in the midgut of *Simulium pertinax* larvae (Diptera, Simuliidae). *Mem. Inst. Oswaldo Cruz*, 99: 493-498.
50. Kohler, H.R., 2002. Localization of metals in cells of saprophagous soil arthropods (Isopoda, Diplopoda, Collembola). *Microsc. Res. Tech.*, 56: 393-401.
51. Pigino, G., M. Migliorini, E. Paccagnini and F. Bernini, 2006. Localisation of heavy metals in the midgut epithelial cells of *Xenillus tegeocranus* (Hermann, 1804) (Acari: Oribatida). *Ecotoxicol. Environ. Saf.*, 64: 257-263.
52. Gregory, M.A., D.J. Marshall, R.C. George, A. Anandraj and T.P. McClurg, 2002. Correlations between metal uptake in the soft tissue of *Perna perna* and gill filament pathology after exposure to mercury. *Mar. Pollut. Bull.*, 45: 114-125.
53. Au, D.W.T., O.V. Yurchenko and A.A. Reunov, 2003. Sublethal effects of phenol on spermatogenesis in sea urchins (*Anthocidaris crassispina*). *Environ. Res.*, 93: 92-98.
54. Meyer, J.N., M.C.K. Leung, J.P. Rooney, A. Sendoel, M.O. Hengartner, G.E. Kisby and A.S. Bess, 2013. Mitochondria as a target of environmental toxicants. *Toxicol. Sci.*, 134: 1-17.
55. Schonthal, A.H., 2012. Endoplasmic reticulum stress: Its role in disease and novel prospects for therapy. *Scientifica*, Vol. 2012. 10.6064/2012/857516
56. Braunbeck, T. and A. Volkl, 1993. Toxicant-Induced Cytological Alterations in Fish Liver as Biomarkers of Environmental Pollution? A Case Study on Hepatocellular Effects of Dinitro-*O*-Cresolin Golden Ide (*Leuciscus idus melanotus*). In: *Fish: Ecotoxicology and Ecophysiology*, Braunbeck, T., W. Hanke and H. Segner (Eds). Wiley-VCH, Weinheim, Germany, ISBN-13: 9783527300105, pp: 55-80.