



# Research Journal of **Microbiology**

ISSN 1816-4935



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)

## Microbial Pectinases and Ruminant Nutrition

H.A. Murad and H.H. Azzaz

Department of Dairy Science, National Research Center, Dokki, Cairo, Egypt

*Corresponding Author: H.A. Murad, Department of Dairy Science, National Research Center, Dokki, Cairo, Egypt  
Tel: 202 33068626 Fax: 202 33370931*

### ABSTRACT

Pectinases are widely distributed in higher plants and microorganisms. They are today one of the upcoming enzymes of the commercial sector. It has been estimated that microbial pectinases account for 25% of the global food enzymes sales. Microbial pectinases can be produced from bacteria including actinomycetes, yeast and fungi. *Aspergillus niger* is the most commonly used fungal species for industrial production of pectinolytic enzymes. Among industrial applications of pectinases are using these enzymes as an animal feed supplementation. This usage of pectinases for ruminant's feed production can reduce the feed viscosity, which increases absorption of nutrients, liberates nutrients, either by hydrolysis of non-biodegradable fibers or by liberating nutrients blocked by these fibers and reduces the amount of faeces. This study deals with types and classification of pectinolytic enzymes, their mode of action, production techniques and the methods of activity assay. Furthermore, it provides a bird's eye view of the possible applications of these enzymes in ruminant nutrition.

**Key words:** Microbial pectinases production, submerged fermentation, solid state fermentation, pectinases activity assay, ruminant nutrition

### INTRODUCTION

Pectinases or pectinolytic enzymes are today one of the upcoming enzymes of the commercial sector. It has been reported that microbial pectinases account for 25% of the global food enzymes sales (Jayani *et al.*, 2005). Primarily, these enzymes are a heterogeneous group of related enzymes that hydrolyze the pectic substances, present mostly in plants.

Pectinolytic enzymes are widely distributed in higher plants and microorganisms (Whitaker, 1990). They are of prime importance for plants as they help in cell wall extension (Ward and Moo-Young, 1989) and softening of some plant tissues during maturation and storage (Sakai, 1992; Aguilar and Huirton, 1990). They also aid in maintaining ecological balance by causing decomposition and recycling of waste plant materials (Jayani *et al.*, 2005).

Microbial production of pectinases has been extensively studied during recent years (Kashyap *et al.*, 2001; Torres *et al.*, 2006). Pectinase production has been reported from bacteria including actinomycetes (Cao *et al.*, 1992; Bruhlmann *et al.*, 1994; Beg *et al.*, 2000), yeast (Blanco *et al.*, 1999; Reid and Ricard, 2000) and fungi (Elegado and Fujio, 1994; Huang and Mahoney, 1999). However, almost all the commercial preparations of pectinases are produced from fungal sources (Singh *et al.*, 1999). *Aspergillus niger* is the most commonly used fungal species for industrial production of pectinolytic enzymes (Naidu and Panda, 1998; Gummadi and Panda, 2003).




Fig. 1: Primary structure of pectic substances (Rexova'-Benkova' and Markovic', 1976)

Higher cost of the production is perhaps the major constraint in commercialization of new sources of enzymes. Though, using high yielding strains, optimal fermentation conditions and cheap raw materials as a carbon source can reduce the cost of enzyme production for subsequent applications in industrial processes. Among industrial applications of pectinases are using these enzymes as an animal feed supplementation. This usage of pectinases for ruminant's feed production can reduce the feed viscosity, which increases absorption of nutrients, liberates nutrients, either by hydrolysis of non-biodegradable fibers or by liberating nutrients blocked by these fibers and reduces the amount of faeces (Hoondal *et al.*, 2002).

This study mainly concentrates on the types and classification of pectinolytic enzymes, their mode of action, production techniques and the methods of activity assay. It also provides a bird's eye view of the possible applications of these enzymes in ruminant nutrition.

## THE SUBSTRATE

Pectic substance is the generic name used for the compounds that are acted upon by the pectinolytic enzymes. They are high molecular weight, negatively charged, acidic, complex glycosidic macromolecules (polysaccharides) that are present in the plant kingdom. They are present as the major components of middle lamella between the cells in the form of calcium pectate and magnesium pectate (Jayani *et al.*, 2005).

Chemically, pectic substances are essentially branched heteropolysaccharides containing between a few hundred and about one thousand building blocks per molecule, with a backbone consisting of galacturonic acid residues part of which are methylesterified. The pectin molecule (Fig. 1) is generally agreed to consist of a chain structure of axial-axial E-(1,4)-linked D-galacturonic acid units, containing blocks of L-rhamnose rich regions, with mainly arabinose, galactose and xylose as side chains. The carboxyl groups of the galacturonic acid are partly esterified by methyl groups and partly or completely neutralized by sodium, potassium or ammonium ions. Some of the hydroxyl groups on C<sub>2</sub> and C<sub>3</sub> may be acetylated (Pilnik and Voragen, 1970).

The American Chemical Society classified pectic substances into four main types as reported by Alkorta *et al.* (1998) as follows:

- Protopectin is the water insoluble pectic substance present in intact tissue. Protopectin on restricted hydrolysis yields pectin or pectic acids
- Pectic acid is the soluble polymer of galacturonans that contains negligible amount of methoxyl groups. Normal or acid salts of pectic acid are called pectates
- Pectinic acids are the polygalacturonan chain that contains >0 and <75% methylated galacturonate units. Normal or acid salts of pectinic acid are referred to as pectinates

- Pectin (Polymethyl galacturonate) is the polymeric material in which, at least, 75% of the carboxyl groups of the galacturonate units are esterified with methanol. It confers rigidity on cell wall when it is bound to cellulose in the cell wall

## THE PECTINASES


The pectinases may be divided in three broader groups as reported by Sakai (1992) and Palomaki and Saarilahti (1997) as follows:

- **Protopectinases:** Degrade the insoluble protopectin and give rise to highly polymerized soluble pectin
- **Esterases:** Catalyze the de-esterification of pectin by the removal of methoxy esters
- **Depolymerases:** Catalyze the hydrolytic cleavage of the  $\alpha$ -(1-4)-glycosidic bonds in the D-galacturonic acid moieties of the pectic substances

Depolymerases act on pectic substances by two different mechanisms, hydrolysis, in which they catalyze the hydrolytic cleavage with the introduction of water across the oxygen bridge and trans-elimination lysis, in which they break the glycosidic bond by a trans-elimination reaction without any participation of water molecule (Fig. 2) (Albersheim *et al.*, 1960; Codner, 2001). Depolymerases can be subdivided into four different categories, depending on the preference of enzyme for the substrate, the mechanism of cleavage and the splitting of the glycosidic bonds (Rexova'-Benkova and Markovic, 1976).

Polygalacturonase and polymethylgalacturonase breakdown pectate and pectin, respectively by the mechanism of hydrolysis. However, polygalacturonate lyase and polymethylgalacturonate lyase breakdown pectate and pectin by  $\beta$  elimination, respectively. Depending upon the pattern of action, i.e., random or terminal, these enzymes are termed as Endo or Exo enzymes, respectively. During the course of investigations and research conducted in recent years, the above classification of pectinolytic enzymes has slowly become obsolete. A more recent and elaborated classification of these enzymes has been provided in Table 1 (Rexova'-Benkova and Markovic, 1976; Ward and Moo-Young, 1989; Whitaker, 1990; Kashyap *et al.*, 2001).

**Protopectinases:** The enzyme that catalyzes the solubilization of protopectin was originally named protopectinase by Brinton *et al.* (1927). Pectinosinase is also synonymous with protopectinase (PPase). Protopectinase catalyzes the following reaction:



PPases are classified into two types, on the basis of their reaction mechanism (Sakai and Okushima, 1982; Sakamoto *et al.*, 1994). A-type PPases react with the inner site, i.e., the polygalacturonic acid region of protopectin, whereas B-type PPases react on the outer site, i.e., on the polysaccharide chains that may connect the polygalacturonic acid chain and cell wall constituents.

A-type PPase are found in the culture filtrates of yeast and yeast-like fungi. They have been isolated from *Kluyveromyces fragilis* IFO 0288, *Galactomyces reesei* L. and *Trichosporon penicillatum* SNO 3 and are referred to as PPase-F, -L and -S, respectively (Whitaker, 1990). B-type




Fig. 2: Enzymatic mode of action of the most frequent depolymerases on the pectin molecule. PMGL: Polymethylgalacturonate lyase (pectin lyase); PMG: Polymethylgalacturonase (pectin hydrolase); PMGE: Polymethylgalacturonate esterase (pectinesterase); PGL: Polygalacturonate lyase (pectate lyase); PG: Polygalacturonase (pectate hydrolase). (Serra *et al.*, 1992)

PPases have been reported in *Bacillus subtilis* IFO 12113 (Sakai and Ozaki, 1988), *B. subtilis* IFO3134 (Sakai and Sakamoto, 1990) and *Trametes* sp. (Sakai *et al.*, 1993) and are referred to as PPase-B, -C and -T, respectively. B-type PPases have also been found in the culture filtrate of a wide range of *Bacillus* sp. (Sakai, 1992).

Ppase activity is assayed by measuring the amount of pectic substance liberated from protopectin by the carbazole-sulphuric acid method (Siebert and Anto, 1946). The pectin concentration is measured as D-galacturonic acid from its standard curve. One unit of PPase activity is defined as the enzyme that liberates pectic substance corresponding to 1 mmol of Dgalacturonic acid per millilitre of reaction mixture under assay conditions.

**Polygalacturonases:** Polygalacturonases (PGases) are the pectinolytic enzymes that catalyze the hydrolytic cleavage of the polygalacturonic acid chain with the introduction of water across the oxygen bridge. They are the most extensively studied among the family of pectinolytic enzymes. The PGases involved in the hydrolysis of pectic substances are endo-PGase (E.C. 3.2.1.15) and exo-PGase (E.C. 3.2.1.67) (Table 1). PGases have the biological, functional and technical applications in food processing and plant-fungal interactions.

Endo-PGases are widely distributed among fungi, bacteria and many yeasts (Luh and Phaff, 1951). They are also found in higher plants and some plant parasitic nematodes (Sakai *et al.*, 1993). They have been reported in many microorganisms, including *Aureobasidium pullulans* (Sakai and Takaoka, 1984), *Rhizoctonia solani* Kuhn (Marcus *et al.*, 1986), *Fusarium moniliforme* (De Lorenzo *et al.*, 1987), *Neurospora crassa* (Polizeli *et al.*, 1991), *Rhizopus stolonifer* (Manachini *et al.*, 1987), *Aspergillus* sp. (Nagai *et al.*, 2000), *Thermomyces lanuginosus* (Kumar and Palanivelu, 1999), *Peacilomyces clavisporus* (Souza *et al.*, 2003). Endo-PGases have also been cloned and genetically studied in a large number of microbial species (Laing and Pretorius, 1993; Naumov *et al.*, 2001). In contrast, exo-PGases occur less frequently. They have been reported in *Erwinia carotovora* (Palomaki and Saarilahti, 1997), *Agrobacterium tumefaciens* (Rodrigues-Palenzuela *et al.*, 1991), *Bacteroides thetaiotamicron* (Tierny *et al.*, 1994), *E. chrysanthemi* (Kobayashi, *et al.*, 2001), *Alternaria mali* (Nozaki *et al.*, 1997), *Fusarium oxysporum* (Maceira *et al.*, 1997), *Ralstonia solanacearum* (Huang and Allen, 1997), *Bacillus* sp. (Kobayashi *et al.*, 2001). Exo-PGases can be distinguished into two types: fungal exo-PGases, which produce monogalacturonic acid as the main end product and the bacterial exo-PGases, which produce digalacturonic acid as the main end product (Sakai *et al.*, 1993). Occurrence of PGases in plants has also been reported (Pressey and Avants, 1973; Alonso *et al.*, 2003).

PGase activity is determined on the basis of measuring, during the course of the reaction: (1) the rate of increase in number of reducing groups and (2) the decrease in viscosity of the substrate solution (Rexova-Benkova and Markovic, 1976). The amount of reducing sugar can be readily measured by colorimetric methods like 3, 5-dinitrosalicylate reagent method (Miller, 1959) and the arsenomolybdate-copper reagent method (Nelson, 1944; Collmer *et al.*, 1988). One unit of enzyme activity is defined as the enzyme that releases 1 mmol/mL/min galacturonic acid under standard assay conditions.

Viscosity reduction measurements have also found widespread use in determining the PGase activity (Roboz *et al.*, 1952):

$$\% \text{ reduction in viscosity} = \frac{T_o - T_t}{T_o - T_w}$$

Table 1: An extensive classification of pectinolytic enzymes (Jayani *et al.*, 2005)

Enzyme	Action mechanism	Action pattern	Primary substrate	Product
<b>Esterase</b>				
1. Pectin methyl esterase	Hydrolysis	Random	Pectin	Pectic acid + methanol
<b>Depolymerizing enzymes</b>				
<b>a. Hydrolases</b>				
1. Protopectinases	Hydrolysis	Random	Protopectin	Pectin
2. Endopolygalacturonase	Hydrolysis	Random	Pectic acid	Oligogalacturonates
3. Exopolygalacturonase	Hydrolysis	Terminal	Pectic acid	Monogalacturonates
4. Exopolygalacturonan-digalacturono hydrolase	Hydrolysis	Penultimate bonds	Pectic acid	Digalacturonates
6. $\Delta 4:5$ Unsaturated oligogalacturonate hydrolases	Hydrolysis	Terminal	$\Delta 4:5$ Galacturonate	Unsaturated monogalacturonates and saturated (n-1)
8. Endopolymethyl-galacturonases	Hydrolysis	Terminal	Highly esterified pectin	Oligogalacturonates
<b>b. Lyases</b>				
1. Endopolygalacturonase lyase	Trans-elimination	Random	Pectic acid	Unsaturated oligogalacturonates
2. Exopolygalacturonase lyase	Trans-elimination	Penultimate bonds	Pectic acid	Unsaturated digalacturonates
3. Oligo-D-galactosiduronate lyase	Trans-elimination	Terminal	Unsaturated digalacturonates	Unsaturated monogalacturonates
4. Endopolymethyl-D-galactosiduronate lyase	Trans-elimination	Random	Unsaturated poly-(methyl-D-digalacturonates)	Unsaturated methyloligogalacturonates
5. Exopolymethyl-D-galactosiduronate lyase	Trans-elimination	Terminal	Unsaturated poly-(methyl-D-digalacturonates)	Unsaturated methylmonogalacturonates


where,  $T_o$ ,  $T_t$  and  $T_w$  represent the flow rate (in a capillary viscometer) in seconds for the reaction mixture without enzyme, the test mixture and water, respectively. The unit of enzyme activity is mostly selected as the amount of enzyme required for attaining a certain decrease of viscosity per unit time. However, this method has met with limited success. There is no direct correlation between viscosity reduction and number of glycosidic bonds hydrolyzed. PGase activity can also be determined by the cup-plate method (Dingle *et al.*, 1953). Cups are cut out from the solidified agar containing the substrate and are filled with the enzyme solution. After lapse of a certain period of time, the zones of degraded substrate are stained with iodine and quantified.

**Lyases:** Lyases (or transeliminases) perform non-hydrolytic breakdown of pectates or pectinates, characterized by a trans-eliminative split of the pectic polymer (Sakai *et al.*, 1993). The lyases break the glycosidic linkages at C-4 and simultaneously eliminate H from C-5, producing a D 4:5 unsaturated products (Codner, 2001; Albersheim *et al.*, 1960).

Lyases can be classified into following types on the basis of the pattern of action and the substrate acted upon by them (Table 1):

(I) Endopolygalacturonate lyase (EndoPGL, E.C. 4.2.2.2); (II) exopolygalacturonate lyase (ExoPGL, E.C. 4.2.2.9); (III) endopolymethylgalacturonate lyase (EndoPMGL, E.C. 4.2.2.10); (IV) exopolymethylgalacturonate lyase (ExoPMGL).

The reactions catalyzed by lyases can be illustrated as follows:




Polygalacturonate lyases (Pectate lyases or PGLs) are produced by many bacteria and some pathogenic fungi with endo-PGLs being more abundant than exo-PGLs. PGLs have been isolated from bacteria and fungi associated with food spoilage and soft rot. They have been reported in *Colletotrichum lindemuthionum* (Wijesundera *et al.*, 1984), *Bacteroides thetaiotaomicron* (McCarthy *et al.*, 1985), *Erwinia carotovora* (Kotoujansky, 1987), *Amucala* sp. (Bruhlmann, 1995), *Pseudomonas syringae* pv. Glycinea (Margo *et al.*, 1994), *Colletotrichum magna* (Wattad *et al.*, 1995), *E. chrysanthemi* (Favey *et al.*, 1992; Shevchik *et al.*, 1997), *Bacillus* sp. (Kobayashi *et al.*, 1999; Takao *et al.*, 2001), *Bacillus* sp. DT-7 (Kashyap *et al.*, 2000), *C. gloeosporioides* (Yakoby *et al.*, 2000; Drori *et al.*, 2003)

Very few reports on the production of polymethylgalacturonate lyases (pectin lyases or PMGLs) have been reported in literature. They have been reported to be produced by *Aspergillus japonicus* (Ishii and Yokotsuka, 1975), *Penicillium paxilli* (Szajer and Szajer, 1982, 1985), *Penicillium* sp. (Diaz Borrás *et al.*, 1987; Alana *et al.*, 1990; Sapunova *et al.*, 1995), *Pythium splendens* (Chen *et al.*, 1998) *Pichia pinus* (Moharib *et al.*, 2000), *Aspergillus* sp. (Sunnotel and Nigam, 2002), *Thermoascus auratniacus* (Martins *et al.*, 2002).

The most convenient method of following the activity of lyases is to measure the increase in absorbance at 235 nm due to formation of the  $\Delta 4:5$  double bonds produced at the non-reducing ends of the unsaturated products (Albersheim, 1966; Whitaker, 1990; Liao *et al.*, 1999). The molar extinction coefficients for PGL and PMGL are  $4.6 \times 10^3$  and  $5.5 \times 10^3$  M/cm, respectively. One unit of enzyme activity is defined as the amount of enzyme that releases 1 mmol of unsaturated product per minute under assay conditions. Reducing group methods (Miller, 1959; Collmer *et al.*, 1988) are also useful in determining the lyase activity. Viscosity reduction method (Roboz *et al.*, 1952), in conjunction with a reducing group method or along with intermediate product analysis by HPLC or GC, can be used to distinguish between endo-and exo-splitting enzymes (Albersheim, 1966).

**Pectinesterase:** Pectinesterase (PE, Pectin pectylhydrolase, E.C. 3.1.1.11), often referred to as pectinmethylesterase, pectase, pectin methoxylase, pectin demethoxylase and pectolipase, is a carboxylic acid esterase and belongs to the hydrolase group of enzymes (Whitaker, 1984). It catalyzes the deesterification of methyl ester linkages of galacturonan backbone of pectic substances to release acidic pectins and methanol (Cosgrove, 1997). The resulting pectin is then acted upon by polygalacturonases and lyases (Sakai *et al.*, 1993; Prade *et al.*, 1999). The mode of action of PE varies according to its origin (Micheli, 2001). Fungal PEs act by a multi-chain mechanism, removing the methyl groups at random. In contrast, plant PEs tends to act either at the non-reducing end or next to a free carboxyl group and proceed along the molecule by a single chain mechanism. The reaction catalyzed by PE can be represented as follows (Forster, 1988):



PE activity is implicated in cell wall metabolism including cell growth, fruit ripening, abscission, senescence and pathogenesis (Gaffe *et al.*, 1997; Dorokhov *et al.*, 1999). Commercially PE can be used for protecting and improving the texture and firmness of several processed fruits and vegetables as well as in the extraction and clarification of fruit juices (Fayyaz *et al.*, 1993). PE is



found in plants, plant pathogenic bacteria and fungi (Hasunuma *et al.*, 2003). It has been reported in *Rhodotorula* sp. (Vaughan *et al.*, 1969), *Phytophthora infestans* (Forster, 1988), *Erwinia chrysanthemi* B341 (Pitkanen *et al.*, 1992), *Saccharomyces cerevisiae* (Gainvors *et al.*, 1994), *Lachnospira pectinoschiza* (Cornick *et al.*, 1994), *Pseudomonas solanacearum* (Schell *et al.*, 1994), *Aspergillus niger* (Maldonado *et al.*, 1994; Maldonado and de Saad, 1998), *Lactobacillus lactis* subsp. *Cremoris* (Karam and Belarbi, 1995), *Penicillium frequentans* (Kawano *et al.*, 1999), *E. chrysanthemi* 3604 (Laurent *et al.*, 2000), *Penicillium occitanis* (Hadj-Taieb *et al.*, 2002), *A. japonicus* (Semenova *et al.*, 2003) and others. There are many reports of occurrence of PE in plants, viz., *Carica papaya* (Fayyaz *et al.*, 1993; Innocenzo and Lajalo, 2001), *Lycopersicum esculentum* (Warrilow *et al.*, 1994), *Prunus malus* (Macdonald and Evans, 1996), *Vitis vinifera* (Corredig *et al.*, 2000), *Citrus* sp. (Arias and Burns, 2002), *Pouteria sapota* (Arenas-Ocampo *et al.*, 2003) and *Malpighia glabra* L. (Assis *et al.*, 2004).

PE activity is most readily followed by gel diffusion assay, as described by Downie *et al.* (1998). Increased binding of ruthenium red to pectin, as the number of methyl esters attached to the pectin decreases, is used as the basis of the assay. The unit of activity in nano-or picokatal is calculated, based on the standard curve generated from the log-transformed commercial enzyme activity versus stained zone diameter. The sensitivity, specificity and simplicity of this PE assay are superior to all others. PE activity can also be measured by using a pH stat because ionization of the carboxyl group of the product releases a proton, which causes a change in pH (Whitaker, 1984).

## MICROBIAL PECTINASES PRODUCTION

In developing nations, there is noticeable growth of food and feed processing industries, leading to an increase in the demand for pectic enzymes. Unfortunately, many of these nations depend largely on imported pectinase, which has become more expensive because of higher cost of foreign exchange. Therefore, with a view of improving the utilization of horticultural product in the depressed economy of a nation like Egypt, it is necessary to carry out research on the production of pectic enzymes locally.

**Microbial pectinases production techniques:** There were two fermentation techniques we can use for pectinases production, as many other enzymes (Murad and Foda, 1992) these techniques are Solid State Fermentation (SSF) and submerged fermentation (SmF).

Solid state fermentation is defined as the cultivation of microorganisms on moist solid supports, either on inert carriers or on insoluble substrates that can be used as carbon and energy source. This process occurs in the absence or near absence of free water in the space between substrate particles. In this system, water is present in the solid substrate whose capacity for liquid retention varies with the type of material (Lonsane *et al.*, 1985; Pandey *et al.*, 2001). In contrast, in submerged fermentation (SmF) the nutrients and microorganisms are both submerged in water (Grigelmo-Migeul and Martin-Belloso, 1998).

Approximately 90% of all industrial enzymes are produced in SmF, frequently using specifically optimized, genetically manipulated microorganisms. In this respect SmF processing offers an insurmountable advantage over SSF. On the other hand, almost all these enzymes could also be produced in SSF using wild-type microorganisms (Filer, 2001; Pandey *et al.*, 2001). Interestingly, fungi, yeasts and bacteria that were tested in SSF in recent decades exhibited different metabolic strategies under conditions of solid state and submerged fermentation.

Table 2: Biotechnological advantages of Solid State Fermentation (SSF) against submerged technology (SmF) (Holker *et al.*, 2004)

Advantages	Consequences	Problems to be solved
<b>Biological advantages</b>		
Low water demand	Less waste water	Building of moisture gradients
High concentration of the end product	Lower downstream costs	
Catabolite repression significantly lower or missing	Fermentation in the presence of glucose	Building of substrate gradients Building of pH gradients
Utilisation of solid substrate	High concentration of the growth substrates	
Lower sterility demands	Mixed cultures of fermenting microorganisms	
Solid support for microorganism		
Simulation of the natural environment	Better performance of cultivated microorganisms	
Fermentation of water-insoluble solid substrates		
Mixed culture of microorganisms	Synergism of metabolic performance	
<b>Processing advantages</b>		
High-volume productivity	Smaller fermenter volumes	
Low energy demand for heating		Building of temperature gradients
Easy aeration		Building of oxygen gradients on a large scale
Utilisation of otherwise unusable carbon sources	Cheap and abundant carbon sources	
No anti-foam chemicals	No lost of microorganisms during fermentation	

The aim of SSF is to bring the cultivated fungi or bacteria into tight contact with the insoluble substrate and thus to achieve the highest substrate concentrations for fermentation. This technology results, only on a small scale, in several processing advantages of significant potential economic and ecological importance as compared with SmF (Table 2).

However, there are also several disadvantages of SSF, which have discouraged use of this technique for industrial production. The main obstructions are due mainly to the build-up of gradients of temperature, pH, moisture, substrate concentration or CO<sub>2</sub> during cultivation, which are difficult to control under limited water availability. It has become clear (as mentioned in nearly every review cited) that the cost-factor for the production of bulk-ware enzymes in most cases favors SSF over SmF. The low estimated costs of SSF are due to the rather traditional preferential claim of SSF, viz. SSF utilises complex, heterogenous agricultural wastes as substrates and uses low-cost technology regarding sterility and regulation demands. However, attempts to reduce costs progress in SSF because of the strongly increased diversity in SSF research.

There is no consensus on the methods, the microorganisms or the substrates used, that would allow comparison with other cultivation technologies. The broad spectrum of substrates used represents an especially severe problem. As already mentioned, one great advantage of SSF has always been the possibility of using substrates that are abundant, cheap and not applicable to SmF. However, regardless of the differences in process up-scaling, the scientific and technological impact of research data is difficult to compare when results are obtained with different microorganisms producing different products and using a vast variety of substrates (Rafae *et al.*, 2006; Abd El-Salam *et al.*, 1994). There are many substrates could be used as, pineapple, mixed fruit, maosmi waste, orange bagasse, mango waste, corn tegument, almond meal, apple pomace, molasses, banana peels, amaranth grain, eucalyptus kraft pulp, coffee residues and some food industry wastes (Holker *et al.*, 2004; Murad, 1998).

To facilitate comparison of results, the use of inert substrates as solid supports is becoming increasingly important (Gautam *et al.*, 2002; Ooijkaas *et al.*, 2000). Surprisingly, biological parameters, such as the stability of the produced enzymes at high temperature or extreme pH, have also been reported to be better in SSF (Deschamps and Huet, 1985; Acuna-Arguelles *et al.*, 1995). Capability repression or protein degradation by proteases severe problems in SmF were often reduced or absent in SSF (Pereira *et al.*, 1993; Aguilar *et al.*, 2001). In contrast, much less research has been carried out to evaluate the metabolic differences of microorganisms when cultivated in SSF or SmF.

**The factors affecting microbial pectinases production:** Environmental and nutritional factors are known to have marked effects on enzyme production by microorganisms. There are, therefore, variations in optimum conditions for pectic enzyme production. Some of the cultural factors that affect the production of pectic enzymes are presented in this study.

**Initial pH of growth medium:** According to Shoichi *et al.* (1985) the initial pH of the medium has a great effect on the growth of the organism, on the membrane permeability, also on the biosynthesis and stability of the enzymes (Murad, 1998; Murad and Salem, 2001). Optimum production of pectic enzymes from many moulds has been reported to be within the acidic pH range (Zetelaki-Horvath, 1980; Shin *et al.*, 1983). Zheng and Shetty (1999), reported that, polygalacturonase produced from *Lentinus edodes* has a relatively lower optimum pH (pH 5.0) in addition, Piccoli-Valle *et al.* (2001) observed that a high polygalacturonase and pectin esterase activity was showed by *P. griseoroseum* in more acid pH of 4.5 and 5. Also, Silva *et al.* (2002) found that *P. viridicatum* showed maximum production of polygalacturonase and pectinlyase at a pH of 4.5 and 5, respectively. Fawole and Odunfa (2003) reported that the optimum pectolytic activity observed was at pH 5. Phutela *et al.* (2005) concluded that the thermophilic fungi *A. fumigatus* Fres expressed maximum pectinase (1116 Ug-1) activity at pH 4.0 while polygalacturonase was active at pH 5.0 (1270 Ug-1). Also, Debing *et al.* (2005) found that the pH 6.5 was the optimal pH for pectinase production from *A. niger* by solid state fermentation. Reda *et al.* (2008) found that the polygalacturonase productivity by *Bacillus firmus*-I-10104 reached its maximum at initial pH 6.0 and 6.2. Rasheedha *et al.* (2010) found that *P. chrysogenum* exhibited maximum polygalacturonase production at initial pH of 6.5. However, the mechanism by which the pH acts on the production pectic enzyme is not known.

**Incubation period:** The time of fermentation had a profound effect on microbial product formation (Murad and Foda, 1992; Murad, 1998; Murad and Salem, 2001). Maximum production of pectic enzyme from different moulds varies from 1 to 6 days (Ghildyal *et al.*, 1981). Castilho *et al.* (2000) reported that the highest polygalacturonase activities were obtained by *A. niger* after 70 h of fermentation period. In addition, Fawole and Odunfa (2003) reported that optimum production of pectinmethylesterase was obtained after 4 days of fermentation under submerged fermentation condition. Moreover, Sarvamangala and Dayanand (2006) observed a gradual increase in the production of pectinase from deseeded sunflower head by *A. niger* after 72 h of fermentation period in submerged and up to 96 h in solid-state conditions. Reda *et al.* (2008) found that the level of polygalacturonase increased gradually with increasing the incubation period up to a maximum of 96 h by *Bacillus firmus*-I-10104 under solid state fermentation conditions.

**Nitrogen source:** The effects of organic and inorganic nitrogen sources on the production of pectinase were extensively studied. The observations of Hours *et al.* (1988) suggested that lower levels of  $(\text{NH}_4)_2\text{SO}_4$  (0.16%), or  $\text{K}_2\text{HPO}_4$  (0.1%) added to the growth medium as inorganic nitrogen sources did not influence pectinase yield. In addition Galiotou-Panayotou and Kapantai (1993) observed that ammonium phosphate and ammonium sulphate did influence production of pectinase positively but also recorded the inhibitory effects of ammonium nitrate and potassium nitrate on pectinase production. Moreover, Sarvamangala and Dayanand (2006) revealed that both ammonium phosphate and ammonium sulphate did influence production of pectinase positively in both submerged and solid-state conditions. In contrast, Sapunova (1990) found that ammonium salts stimulated the pectinolytic enzyme production in *A. alliaceus* BIM-83. Moreover, Sapunova *et al.* (1997) has also observed that  $(\text{NH}_4)_2\text{SO}_4$  stimulated pectinase synthesis, as in its absence fungus displayed a slight proteolytic activity and did not produce extracellular pectinases. In addition, Fawole and Odunfa (2003) found that ammonium sulphate and ammonium nitrate were good nitrogen sources for pectic enzyme production from *A. niger* while glycine and tryptophan did not support enzyme production. Also, Phutela *et al.* (2005) reported that  $(\text{NH}_4)_2\text{SO}_4$  stimulated pectinase production, as in its absence fungus displayed a slight proteolytic activity and did not produce extracellular pectinases. Phutela *et al.* (2005) found that the presence of yeast extract +  $(\text{NH}_4)_2\text{SO}_4$  in growth medium supported maximal production of pectinase ( $925 \text{ Ug}^{-1}$ ) followed by malt sprouts +  $(\text{NH}_4)_2\text{SO}_4$  ( $785 \text{ Ug}^{-1}$ ), which also supported maximal polygalacturonase activity ( $938 \text{ Ug}^{-1}$ ). In addition, Rasheedha *et al.* (2010) found that ammonium sulphate has enhanced the production of *P. chrysogenum* pectinase.

On the other hand, report of Aguilar *et al.* (1991) showed yeast extract (organic nitrogen source) as the best inducer of exopectinases by *Aspergillus* sp. Moreover Kashyap *et al.* (2003) found that, yeast extract, peptone and ammonium chloride were found to enhance pectinase production up to 24% and addition of glycine, urea and ammonium nitrate inhibited pectinase production. Also, Reda *et al.* (2008) found that the maximum value of polygalacturonase productivity by *Bacillus firmus*-I-10104 reached up to  $350 \text{ U mL}^{-1}$  in the presence of peptone as a nitrogen source in the growth medium. In addition, Vivek *et al.* (2010) found that organic nitrogen sources showed higher endo, exo pectinases activities than inorganic nitrogen sources. Also the increasing trend in the enzymes activity with the increase in nitrogen source content was observed in the case of organic nitrogen sources while decreasing trend observed for inorganic nitrogen sources. Soybean meal (4%) showed the maximum Exopectinase activity of  $5128 \text{ IU g}^{-1}$  and endo-pectinase activity of  $793 \text{ IU g}^{-1}$ .

**Carbon source:** An adequate supply of carbon as energy source is critical for optimum growth affecting the growth of organism and its metabolism. Aguilar and Huitron (1987) reported that the production of pectic enzymes from many moulds is known to be enhanced by the presence of pectic substrates in the medium. Fawole and Odunfa (2003) found that pectin and polygalacturonic acid promoted the production of pectic enzyme and they observed the lack of pectolytic activity in cultures with glucose as sole carbon source reflects the inducible nature of pectic enzyme from the strain of *A. niger*. However, when different concentrations of glucose were added to the medium containing pectin, production of pectic enzymes was inhibited at high glucose concentration while low glucose concentrations (0.5% w/v) stimulated enzyme production. Also, the reducing sugar content of the culture filtrate increased with increase in the amount of glucose added to the growth medium. The ability of high concentrations of glucose in the medium to meet growth requirement

of the organism probably made the breakdown of pectin in the medium unnecessary or minimal and thus the low pectic activities observed in cultures. Phutela *et al.* (2005) stated that wheat bran supported maximum pectinase production ( $589 \text{ U g}^{-1}$ ) while pure pectin give the maximum production of polygalacturonase ( $642 \text{ U g}^{-1}$ ). Sarvamangala and Dayanand (2006) reported that glucose (4-6%) increase the production of pectinase in submerged condition whereas 6-8% sucrose gives better yield of pectinase in solid-state condition. Reda *et al.* (2008) reported that *Solanum tuberosum* (ST) peels was the best carbon source for polygalacturonase production by *Bacillus firmus*-I-10104 under solid state condition.

## PURIFICATION OF MICROBIAL PECTINASES

In order to characterize and study the properties of microbial pectinases the enzymes must be purified. Important purification methods for the isolation of different pectinases are briefly summarized in this section. Pectinases from various sources of microorganisms have been purified to homogeneity. An exo-PG has been separated from mycelial extracts of *Aspergillus niger* by eluting from DEAE cellulose with 0.2 M sodium acetate buffer at pH 4.6. Purification was efficient with 209-fold increase in specific activity with a recovery of 8.6% and the enzyme displayed its full activity only in the presence of  $\text{Hg}_2^+$  ions (Mill, 1966a). A second PG was isolated with 205-fold increase in specific activity with a recovery of 1%. These two PGs are differentiated by their optimum pH and PG II was not inhibited by chelating agents and did not require  $\text{Hg}_2^+$  for activity (Mill, 1966b).

Benkova and Slezarik (1966) developed a purification strategy for the isolation of extracellular PMG, PG and PE. The enzyme was salted out with ammonium sulphate and precipitated with ethanol after gel filtration through Sephadex G-25. Repeated chromatography on DEAE-cellulose column yielded a homogeneous preparation of enzyme. Exo-PG, Endo-PG and pectinesterase have been separated from the culture filtrate of *Trichoderma reesei* by Sephadex chromatography (Markovic *et al.*, 1985). Polygalacturonase from *Rhizopus stolonifer* has been purified up to 10-fold by ethanol precipitation followed by CM-Sepharose 6B ion exchange chromatography and gel filtration by Sephadex G-100 (Manachini *et al.*, 1987). PG and PL (pectinlyase) from *Aureobasidium pullulans* LV10 have been separated by CM-Sepharose 6B followed by column chromatography (DEAE-cellulose column) and gel filtration on Sephadex G-100 (Manachini *et al.*, 1988). PG and PL (pectinlyase) have been separated into PG I and PG II and PL I and PL II, respectively. Pectatelyase (PGL) was synthesized by *Amycolata* species and the extracellular crude enzyme has been purified to homogeneity by both cation and anion exchange columns and hydrophobic interaction chromatography (Bruhlmann, 1995). It has been observed that purification resulted in a 4-fold increase in specific activity with 37% recovery. Pectinases from *Clostridium acetobutylicum* ID 91-36 a UV mutant, has been purified by cation exchange chromatography on a Sepharose column by eluting with NaCl (Seethaler and Hartmeier, 1992).

Endopectate lyase synthesized by *Bacillus macerans* has been purified by ammonium sulphate precipitation followed by DEAE-Sephadex A-50 chromatography and CM-cellulofine chromatography (Miyazaki, 1991). Similarly endopectate lyase I/IV have been isolated from the culture filtrate of *Erwinia carotovora* by CM Sepharose CL 6B chromatography, Sephadex S-200 gel filtration and isoelectric focusing (Tanabe *et al.*, 1984). Kobayashi *et al.* (2001) purified the first bacterial; exo-PG from *Bacillus* sp. strain KSM-P443 to homogeneity. This enzyme releases exclusively mono-galacturonic acid from polygalacturonic acid (PGA), di-, tri-, tetra- and penta-galacturonic acids. They also determined the N-terminal sequence and concluded that no sequence matched with other pectinases reported to-date.

An extracellular endo-PG produced by *Aspergillus awamori* IFO 4033 was purified homogeneity using cation-exchange and size-exclusion chromatographic columns (Nagai *et al.*, 2000). Sakamoto *et al.* (1994) isolated protopectinase-N (PPN) and protopectinase-R (PPR) from the culture filtrate of *Bacillus subtilis* IFO3134. These enzymes have been purified by hydrophobic interaction chromatography on butyl-toyopearl 650 M, cation exchange chromatography on CMtoyopearl650 M and gel filtration on sepharose 12HR. These enzymes have been found to be stable over a wide range of pH and temperature. Endopectate lyase produced by *Erwinia caratovara* FERM P-7576 has been selectively co-sedimented with an extracellularly produced lipopolysaccharide lipid complex (Fukoka *et al.*, 1990). The cell free broth was precipitated and the enzyme separated by gel chromatography with a specific activity of 710 U mg<sup>-1</sup> of protein. Co-sedimentation has been affected by pH and ionic strength. Denis *et al.* (1990) studied the effect of shear stress on purification of five isozymes of pectate lyase produced by *Erwinia chrysanthemi* 3937 in ultrafiltration equipment. Activity was not affected during 7 h of pumping and 36% activity was lost after 25 000 passes.

New affinity matrices have been developed for the purification of pectinases, which possess better mechanical and chemical stability than those cross-linked one with pectic acid (Lobarzewski *et al.* 1985). The culture filtrate was desalted on a Sephadex G-25 column. The supports used were silanized controlled pore glass, silica gel silanized with 5-aminopropyl triethoxysilane. All supports were activated with 3-(3-dimethylaminopropyl) carbodiimide and best results were obtained with silanized controlled pore glass. Gupta *et al.* (1996) developed an affinity precipitation technique for separation of selective proteins using heterobifunctional ligands. They used a soluble form of the ligand for affinity binding and then precipitation was induced for separating the protein complex. Alginate was used as successful ligand for pectinases. Wu *et al.* (2000) studied the partitioning behavior of endo-PG and total protein from *Kluyveromyces marxianus* culture broth in polyethylene glycol (PEG) and PEG-potassium phosphate aqueous two-phase systems (ATPS). Both enzyme and total protein partitioned into the bottom phase in both systems. Since, the enzyme protein content in the total protein of the fermentation broth was higher, they proposed that separating endo-PG was concentration rather than separation.

An efficient concentration was achieved using ATPS with polymer recycling and dialysis. Immobilized metal ion affinity polysulphone hollow-fiber membranes with a high capacity for protein adsorption were tested for commercial pectic enzyme fractionation (Camperi *et al.*, 2000). The flow through (unbound) fraction has higher activity for pectinase and PE was retained on the column. Similarly, Savary (2001) developed a rapid and simple method to separate PE from PG and other pectinases in potato enzyme preparations using perfusion chromatography (Poros HS). PE was eluted at low salt concentrations (-/80 mM NaCl) and PG at high salt concentrations (-/300 mM NaCl). The development of such columns looks promising for economical purification strategies for pectinases. A literature survey showed that extensive work has been carried out on the purification of PG, PE and PGL and less attention has been focused on PMG protopectinases and oligogalacturonases.

## RUMINANT AND MICROBIAL PECTINASES

Ruminants have an extensive array of microbial enzymes produced in the rumen and these enzymes play an important role in the ruminant digestive process. Pectinolytic activity in the rumen arises from protozoal, fungal and bacterial sources, one of the major pectinolytic bacterial species inhabiting the rumen, *Lachnospira multiparus*, produces a pectin lyase and a pectin

methylesterase (Silley, 1985). Ruminal fungi and protozoa also express one or more of these enzymes (Orpin, 1984; Chesson and Forsberg, 1988; Bonhomme, 1990).

On the other hand, specific enzyme preparations have become valuable tools for economically improving digestive processes in the ruminant (Gado *et al.*, 2009; Murad *et al.*, 2009; Azzaz, 2009). Use supplemental enzymes in the rumen may be important since the digestibility of organic matter in the rumen does not reach 100% and even small changes in digestibility can influence the efficiency of ruminal fermentations. As a result, many strategies have been developed to stimulate the digestion of the fibrous components in ruminant feeds. One particularly promising technology is supplementation of fibrolytic enzymes (Cellulases, xylanases and pectinases) to ruminant's diets (Selinger *et al.*, 1996). Although extensive studies were carried out to investigate the effects of cellulases and xylanases supplementation to ruminant diets on their productivity, study effects of pectinases supplementation to ruminant diets is lacking. Hoondal *et al.* (2002) reported that use of pectinases as ruminant's feed supplementation can reduces the feed viscosity, which increases absorption of nutrients, liberates nutrients, either by hydrolysis of non-biodegradable fibers or by liberating nutrients blocked by these fibers and reduces the amount of faeces. In addition, Jacob *et al.* (2008) reported that the reducing sugar level was increasing gradually for banana fiber treatment with polygalacturonase indicating the effectiveness of the treatment and it was evident from the photographs that the cells were separated after treatment as a result of pectin hydrolysis. This may be reflecting the importance of use pectinases as a fiber degrading enzymes in animal feed production.

To date, little is known about the way that exogenous fibrolytic enzymes improve feed by rumen microorganisms. Several potential modes of action have been proposed. These include: a) increase in microbial colonization of feed particles. Yang *et al.* (1999), enhancing attachment and/or improve access to the cell wall matrix by ruminal microorganisms and by doing so, accelerate the rate of digestion (Nsereko *et al.*, 2000) enhancing the hydrolytic capacity of the rumen due to added enzyme activities and *for* synergy with rumen microbial enzymes (Morgavi *et al.*, 2000). Also to date, it has not been possible to predict the effectiveness of enzyme products for ruminants based on their enzymatic activities because the effectiveness of enzyme products is highly variable (Colombatto *et al.*, 2003). Part of this variability may be due to the types and activities of enzyme, which can vary widely depending on the source organism, growth substrate and culture conditions employed, type of diet fed to animals, enzyme application method and even the level of animal productivity (Beauchemin *et al.*, 2003). Lineweaver and Jansen (1951) study effect of low and high concentration of pectic substances (citrus pulp) on polygalacturonase activity in steer's rumen fluid (*in vitro*). They found that polygalacturonase activity in steer's rumen fluid increased by ten folds compared with basal diet and there was no significant difference in polygalacturonase activity between the low and high level of citrus pulp. Both levels reached about the same maximum activity of 2,000 units mL<sup>-1</sup>) and returned to about 200 units on the basal diet. This ten fold increase in activity of polygalacturonase activity attributed to citrus pulp suggests that the formation of extracellular polygalacturonase was induced by the increase in total pectic substances in the diet.

Finally, we recommended in this study giving more attention to production of microbial pectinases and more attention to their application especially in animal feed production.

## REFERENCES

- Abd-El-Salam, M.H., M.A. Fadel, H.A. Murad, 1994. Bioconversion of sugarcane molasses into xanthan gum. *J. Biotechnol.*, 33: 103-106.

- Acuna-Arguelles, M.E., M. Gutierrez-Rojas, G. Viniegra-González and E. Favela-Torres, 1995. Production and properties of three pectinolytic activities produced by *Aspergillus niger* in submerged and solid-state fermentation. *Applied Microbiol. Biotechnol.*, 43: 808-814.
- Aguilar, G. and C. Huitron, 1987. Stimulation of the production of extracellular pectinolytic activities of *Aspergillus* sp. by galactouronic acid and glucose addition. *Enzyme Microb. Technol.*, 9: 690-696.
- Aguilar, G. and C. Huitron, 1990. Constitutive exo-pectinase produced by *Aspergillus* sp. CH-Y-1043 on different carbohydrate source. *Biotechnol. Lett.*, 12: 655-660.
- Aguilar, G., B. Trejo, J. Garcia and G. Huitron, 1991. Influence of pH on endo and exo-pectinase production by *Aspergillus* species CH-Y-1043. *Can. J. Microbiol.*, 37: 912-917.
- Aguilar, C.N., C. Augur, E. Favela-Torres and G. Viniegra-Gonzalez, 2001. Production of tannase by *Aspergillus niger* Aa-20 in submerged and solid-state fermentation: Influence of glucose and tannic acid. *J. Ind. Microbiol. Biotechnol.*, 26: 296-302.
- Alana, A., I. Alkorta, J.B. Dorminguez, M.J. Llama and J.L. Serra, 1990. Pectin lyase activity in *Penicillium italicum* strain. *Applied Environ. Microbiol.*, 56: 3755-3759.
- Albersheim, P., 1966. Pectin lyase from fungi. *Methods Enzymol.*, 8: 628-631.
- Albersheim, P., H. Neukom and H. Deuel, 1960. Splitting of pectin chain molecules in neutral solutions. *Arch. Biochem. Biophysiol.*, 90: 46-51.
- Alkorta, I., C. Garbisu, M.J. Liama and J.L. Serra, 1998. Industrial applications of pectic enzymes: A review. *Process Biochem.*, 33: 21-28.
- Alonso, J., W. Canet, N. Hawell and R. Alique, 2003. Purification and characterization of carrot (*Daucas carota* L.) pectinesterase. *J. Sci. Food. Agric.*, 83: 1600-1606.
- Arenas-Ocampo, M.L., S. Evangelista-Lozano, R. Arana-Errasquin, A. Jimenez-Aparicio and G. Davila-Ortíz, 2003. Softening and biochemical changes of sapote mamey fruit (*Pouteria sapota*) at different development and ripening stages. *J. Food. Chem.*, 27: 91-107.
- Arias, C. and J. Burns, 2002. A pectinmethylesterase gene associated with a heatstable extract from citrus. *J. Agric. Food. Chem.*, 50: 3465-3472.
- Assis, S.A., P. Fernandes, B.S. Ferreira, H.C. Trevisan, J.M.S. Cabral and O.M.M.F. Oliveira, 2004. Screening of supports for the immobilization of pectin-methylesterase from acerola (*Malpighia glabra* L.). *J. Chem. Technol. Biotechnol.*, 79: 277-280.
- Azzaz, H.H., 2009. Effect of cellulytic enzymes addition to diets on the productive performance of lactating goats. M.Sc. Thesis, Faculty of Agriculture, Cairo University, Egypt, pp: 141.
- Beauchemin, K.A., D. Colombatto, D.P. Morgavi and W.Z. Yang, 2003. Use of exogenous fibrolytic enzymes to improve feed utilization by ruminants. *J. Anim. Sci.*, 81: E37-47E.
- Beg, Q.K., B. Bhushan, M. Kapoor and G.S. Hoondal, 2000. Effect of amino acids on production of xylanase and pectinase from *Streptomyces* sp. QG-11-3. *World J. Microbiol. Biotechnol.*, 16: 211-213.
- Benkova, L.R. and A. Slezarik, 1966. Isolation of extracellular pectolytic enzymes produced by *Aspergillus niger*. *Colection Chechoslov Commun.*, 31: 122-129.
- Blanco, P., C. Sieiro and T.G. Villa, 1999. Production of pectic enzymes in yeast. *FEMS Microbiol. Lett.*, 175: 1-9.
- Bonhomme, A., 1990. Rumen ciliates: their metabolism and relationships with bacteria and their hosts. *Anim. Feed Sci. Technol.*, 30: 203-266.
- Brinton, C.S., W.H. Dore, H.J. Wichmann, J.J. Willaman and C.P. Wilson, 1927. Definitions written by the committee on nomenclature of pectin of the agriculture-food division. *J. Am. Chem. Soc.*, 49: 38-40.



- Bruhlmann, F., 1995. Production and characterization of an extracellular pectate lyase from an *Amycolata* sp. Applied Environ. Microbiol., 61: 3580-3585.
- Bruhlmann, F., K.S. Kim, W. Zimmerman and A. Fiechter, 1994. Pectinolytic enzymes from actinomycetes for the degumming of romie bast fibers. Applied Environ. Microbiol., 60: 2107-2112.
- Camperi, S.A., M. Grasselli and O. Cascone, 2000. High-speed pectic enzyme fractionation by immobilized metal ion affinity membranes. Bioseparation, 9: 173-177.
- Cao, J., L. Zheng and S. Chen, 1992. Screening of pectinase producer from alkalophilic bacteria and study on its potential application in degumming of ramie. Enzyme Microbial Technol., 14: 1013-1016.
- Castilho, L.R., R.A. Medronho and T.L.M. Alves, 2000. Production and extraction of pectinases obtained by solid state fermentation of agroindustrial residues with *Aspergillus niger*. Bioresour. Technol., 71: 45-50.
- Chen, W.C., H.J. Hsieh and T.C. Tseng, 1998. Purification and characterization of a pectin lyase from *Pythium splendens* infected cucumber fruits. Botanical. Bull. Academia. Sinica, 39: 181-186.
- Chesson, A. and C.W. Forsberg, 1988. Polysaccharide Degradation by Rumen Microorganisms. In: The Rumen Microbial Ecosystem, Hobson, P.N. (Ed.). Elsevier, New York, pp: 77.
- Codner, R.C., 2001. Pectinolytic and cellulolytic enzymes in the microbial modification of plant tissues. J. Applied Bacteriol., 84: 147-160.
- Collmer, A., J.L. Ried and M.S. Mount, 1988. Assay methods for pectic enzymes. Meth. Enzymol., 161: 329-335.
- Colombatto, D., D.P. Morgavi, A.F. Furtado and K.A. Beauchemin, 2003. Screening of exogenous enzymes for ruminant diet: relationship between biochemical characteristics and *in vitro* ruminal degradation. J. Anim. Sci., 81: 2628-2638.
- Cornick, N.A., N.S. Jensen, D.A. Stahl, P.A. Hartman and M.J. Allison, 1994. *Lachnospira pectinoschiza* sp. nov., an anaerobic pectinophile from the pig intestine. Int. J. Syst. Bacteriol., 44: 87-93.
- Corredig, M., W. Kerr and L. Wicker, 2000. Separation of thermostable pectinmethylesterase from marsh grapefruit pulp. J. Agric. Food. Chem., 48: 4918-4923.
- Cosgrove, D.J., 1997. Assembly and enlargement of the primary cell wall in plants. Ann. Rev. Cell Biol., 13: 171-201.
- De Lorenzo, G., G. Salvi, L. Degra, R. D'Ovidio and F. Cervone, 1987. Introduction of extracellular polygalacturonases and its mRNA in the phytopathogenic fungus *Fusarium moniliforme*. J. Gen. Microbiol., 133: 3365-3373.
- Debing, J., L. Peizun, F. Stagnitti, X. Xianzhe and L. Li, 2005. Pectinase production by solid fermentation from *Aspergillus niger* by a new prescription experiment. Ectotoxicol. Environ. Safety, 64: 244-250.
- Denis, S., S. Terre, Y. Bertheau and P. Boyaval, 1990. Factors affecting pectatelyase activity during membrane filtration. Biotechnol. Technol., 4: 127-132.
- Deschamps, F. and M.C. Huet, 1985. Xylanase production in solid-state fermentation: A study of its properties. Applied Microbiol. Biotechnol., 22: 177-180.
- Diaz Borrás, M.A., R. Vila Aguilar and E. Hernandez Gimenez, 1987. Comparacion de la actividad pectinolítica *in vitro* entre cepas de penicillium sensibles y resistentes a fungicidas comerciales. Alimentaria, 24: 57-58.

- Dingle, J., W.W. Reid and G.L. Solomons, 1953. The enzymatic degradation of pectin and other polysaccharides. II. Application of the 'Cup-plate' assay to the estimation of enzymes. J. Sci. Food. Agric., 4: 149-155.
- Dorokhov, Y.L., K. Mankinen, O.Y. Frolova, A. Mertis, J. Saarinen and N. Kalkkinen, 1999. A novel function for a ubiquitous plant enzyme pectin methylesterase: the host-cell receptor for the tobacco mosaic virus movement protein. FEBS Lett., 461: 233-238.
- Downie, B., L.M.A. Dirk, K.A. Hadfield, T.A. Wilkins, A.B. Bennet and K.J. Bradford, 1998. A gel diffusion assay for quantification of pectinmethylesterase activity. Anal. Biochem., 264: 149-157.
- Drori, N., H. Kramer-Haimovich, J. Rollins, A. Dinoor, Y. Okon and O. Pines, 2003. External pH and nitrogen source affect secretion of pectate lyase by *Colletotrichum gloeosporioides*. Applied Environ. Microbiol., 69: 3258-3262.
- Elegado, F.B. and Y. Fujio, 1994. Purification and some properties of polygalacturonase from *Rhizopus* sp. LKN. World J. Microbiol. Biotechnol., 10: 256-259.
- Favey, S., C. Bourson, Y. Bertheou, A. Kotoujansky and M. Boccara, 1992. Purification of the acidic pectate lyase of *Erwinia chrysanthemi* 2942 and sequence analysis of the corresponding gene. J. Gen. Microbiol., 138: 499-508.
- Fawole, O.B. and S.A. Odunfa, 2003. Some factors affecting production of pectic enzymes by *Aspergillus niger*. Int. Biodeterioration Biodegradation, 52: 223-227.
- Fayyaz, A., B.A. Asbi, H.M. Ghazali, T.B. Che Man and S. Jinap, 1993. Pectinesterase extraction from papaya. Food Chem., 47: 183-185.
- Filer, K., 2001. The newest old way to make enzymes. Feed Mix., 9: 27-29.
- Forster, H., 1988. Pectinesterase from *Phytophthora infestans*. Methods Enzymol., 161: 355-357.
- Fukoka, S., H. Kamishima, L. Sode and I. Karube, 1990. Facile isolation of endopectate lyase from *Erwinia carotovora* based on electrostatic interaction. Applied Biochem. Biotechnol., 26: 239-248.
- Gado, H.M., A.Z.M. Salem, P.H. Robinson and M. Hassan, 2009. Influence of exogenous enzymes on nutrient digestibility, extent of ruminal fermentation as well as milk production and composition in dairy cows. Anim. Feed Sci. Technol., 154: 36-46.
- Gaffe, J., M.E. Tizando and A.K. Handa, 1997. Characterization and functional expression of a ubiquitously expressed tomato pectin methylesterase. Plant Physiol., 114: 1547-1556.
- Gainvors, A., V. Frezier, H. Lemauresquier, C. Lequart, M. Aigle and A. Belarbi, 1994. Detection of polygalacturonase, pectin lyase and pectinesterase activities in *Saccharomyces cerevisiae* strain. Yeast, 10: 1311-1319.
- Galiotou-Panayotou, M.P.R. and M. Kapantai, 1993. Enhanced polygalacturonase production by *Aspergillus niger* NRRL-364 grown on supplemented citrus pectin. Lett. Applied Microbiol., 17: 145-148.
- Gautam, P., A. Sabu, A. Pandey, G. Szakacs and C.R. Socol, 2002. Microbial production of extra-cellular phytase using polystyrene as inert solid support. Bioresour. Technol., 83: 229-233.
- Ghildyal, N.P., S.V. Ramakrishna, P. Nirmala, B.K. Devi and H.A. Asthana, 1981. Large scale production of pectolytic enzyme by solid state fermentation. Food Sci. Technol., 18: 243-251.
- Grigelmo-Miguel, N. and O. Martin-Belloso, 1998. Characterization of dietary fiber from orange juice extraction. Food Res. Int., 31: 355-361.
- Gummadi, S.N. and T. Panda, 2003. Purification and biochemical properties of microbial pectinases: A review. Process Biochem., 38: 987-996.
- Gupta, M.N., R. Kaul, D. Guoqiang, C. Dissing and B. Mattiasson, 1996. Affinity precipitation of proteins. J. Mol. Recognit., 9: 356-359.

- Hadj-Taieb, N., M. Ayadi, S. Trigui, F. Bouabdollah and A. Gargouri, 2002. Hyper production of pectinase activities by fully constitutive mutant (CT 1) of *Penicillium occitanis*. *Enzyme Microbial Technol.*, 30: 662-666.
- Hasunuma, T., E.I. Fukusaki and A. Kobayashi, 2003. Methanol production is enhanced by expression of an *Aspergillus niger* pectin methylesterase in tobacco cells. *J. Biotechnol.*, 106: 45-52.
- Holker, U., M. Hofer and J. Lenz, 2004. Biotechnological advantages of laboratory-scale solid state fermentation with fungi. *Applied Microbiol. Biotechnol.*, 64: 175-186.
- Hoondal, G., R. Tiwari, R. Tiwari, N. Dahiya and Q.K. Beg, 2002. Microbial alkaline pectinases and their applications: a review. *Applied Microbiol. Biotechnol.*, 59: 409-418.
- Hours, R.A., C.E. Voget and R.J. Ertola, 1988. Some factors affecting pectin-ase production from apple pomace in solid-state-cultures. *Biol. Wastes*, 24: 147-157.
- Huang, Q. and C. Allen, 1997. An exo-poly-a-D-galacturonosidase, Peh B, is required for wild type virulence of *Ralstonia solanacearum*. *J. Bacteriol.*, 179: 7369-7378.
- Huang, L.K. and R.R. Mahoney, 1999. Purification and characterization of an endo-polygalacturonase from *Verticillium albo-atrum*. *J. Applied Microbiol.*, 86: 145-146.
- Innocenzo, M.D. and F.M. Lajalo, 2001. Effect of gamma irradiation on softening changes and enzyme activities during ripening of papaya fruit. *J. Food Biochem.*, 25: 425-438.
- Ishii, S. and T. Yokotsuka, 1975. Purification and properties of pectin lyase from *Aspergillus japonicus*. *Agric. Biol. Chem.*, 39: 313-321.
- Jacob, N., K.N. Niladevi, G.S. Anisha and P. Prema, 2008. Hydrolysis of pectin: An enzymatic approach and its application in banana fiber processing. *Microbiol. Res.*, 163: 538-544.
- Jayani, R.S., S. Saxena and R. Gupta, 2005. Microbial pectinolytic enzymes a review. *Process Biochem.*, 40: 2931-2944.
- Karam, N.E. and A. Belarbi, 1995. Detection of polygalacturonase and pectinesterases in lactic acid bacteria. *World J. Microbiol. Biotechnol.*, 11: 559-563.
- Kashyap, D.R., S. Chandra, A. Kaul and R. Tewari, 2000. Production, purification and characterization of pectinase from a *Bacillus* sp. DT7. *World J. Microbiol. Biotechnol.*, 16: 277-282.
- Kashyap, D.R., P.K. Vohra, S. Chopra and R. Tewari, 2001. Applications of pectinases in the commercial sector: A review. *Bioresour. Technol.*, 77: 215-227.
- Kashyap, R., K.S. Soni and R. Tewari, 2003. Enhanced production of pectinase by *Bacillus* sp. DT7 using solid-state fermentation. *Bioresour. Technol.*, 88: 251-254.
- Kawano, C.Y., M.A.S.C. Chellegatti, S. Said and M.J.V. Fonseca, 1999. Comparative study of intracellular and extracellular pectinases produced by *Penicillium frequentans*. *Biotechnol. Applied Biochem.*, 29: 133-140.
- Kobayashi, T., K. Koike, T. Yoshimatsu, N. Higaki, A. Suzumatsu and T. Ozawa, 1999. Purification and properties of a low-molecular weight, high-alkaline pectate lyase from an alkaliphilic strain of *Bacillus*. *Biosci. Biotechnol. Biochem.*, 63: 72-75.
- Kobayashi, T., N. Higaki, N. Yajima, A. Suzumatsu, H. Hagihara, S. Kawai and S. Ito, 2001. Purification and properties of a galacturonic acidreleasing exopolygalacturonase from a strain of *Bacillus*. *Biosci. Biotechnol. Biochem.*, 65: 842-847.
- Kotoujansky, A., 1987. Molecular genetics of pathogenesis by soft-rot *Erwinia*. *Annu. Rev. Phytopathol.*, 25: 405-430.

- Kumar, S.S. and P. Palanivelu, 1999. Purification and characterization of an exopolysaccharidase from the thermophilic fungus, *Thermomyces lanuginosus*. World. J. Microbiol. Biotechnol., 15: 643-646.
- Laing, E. and I.S. Pretorius, 1993. Co-expression of an *Erwinia chrysanthemi* pectate lyase-encoding gene (pel E) and an *E. carotovora* polygalacturonase- encoding gene (peh 1) in *Saccharomyces cerevisiae*. Applied Microbiol. Biotechnol., 75: 149-158.
- Laurent, F., A. Kotoujansky and Y. Bertheau, 2000. Overproduction in *Escherichia coli* of the pectin methylesterase A from *Erwinia chrysanthemi*3937: one-step purification, biochemical characterization and production of polyclonal antibodies. Can. J. Microbiol., 46: 474-480.
- Liao, C.H., L. Revear, A. Hotchkiss and B. Savary, 1999. Genetic and biochemical characterization of an exo polygalacturonase and a pectate lyase from *Yersinia enterocolitica*. Can. J. Microbiol., 45: 396-403.
- Lineweaver, I. and E.F. Jansen, 1951. Pectic enzymes. Adv. Enzymol., 11: 267-267.
- Lobarzewski, J., J. Fiedurek, G. Ginalska and J. Wolski, 1985. New matrices for the purification of pectinases by affinity chromatography. Biochem. Biophys. Res. Commun., 131: 66-74.
- Lonsane, B.K., N.P. Ghildyal, S. Budiatman and S.V. Ramakrishna, 1985. Engineering aspects of solid state fermentation. Enzyme Microb. Technol., 7: 258-265.
- Luh, B.S. and H.J. Phaff, 1951. Studies on polygalacturonase of certain yeasts. Arch. Biochem. Biophys., 33: 212-227.
- Macdonald, H.C. and R. Evans, 1996. Purification and properties of apple pectinesterase. J. Sci. Food. Agric., 70: 321-326.
- Maceira, F.I.G., A.D. Pietro and M.I.G. Roncero, 1997. Purification and characterization of a novel exopolysaccharidase from *Fusarium oxysporum* f. sp. *lycopersici*. FEMS Microbiol. Lett., 154: 37-43.
- Maldonado, M.C. and S.A.M. de Saad, 1998. Production of pectinesterase and polygalacturonase by *Aspergillus niger* in submerged and solid-state systems. J. Ind. Microbiol. Biotechnol., 20: 34-38.
- Maldonado, M.C., A.M.S. Saad and D.A.S. Callieri, 1994. Purification and characterization of pectinesterase produced by a strain of *Aspergillus niger*. Curr. Microbiol., 24: 193-196.
- Manachini, P.L., M.G. Fortina and C. Parini, 1987. Purification and properties of an endopolysaccharidase produced by *Rhizopus stolonifer*. Biotechnol. Lett., 9: 219-224.
- Manachini, P.L., C. Parini and M.C. Fortina, 1988. Pectic enzymes from *Aureobasidium pullulans*, LV10. Enzyme Microbiol. Biotechnol., 10: 682-685.
- Marcus, L., I. Barash, B. Sneh, Y. Koltin and A. Finkler, 1986. Purification and characterization of pectolytic enzymes produced by virulent and hypovirulent isolates of *Rhizoctonia solani* Kuhn. Physiol. Mol. Plant Pathol., 29: 325-336.
- Margo, P., L. Varvaro, G. Chilosi, C. Avanzo and G.M. Balestra, 1994. Pectinolytic enzymes produced by *Pseudomonas syringae* pv. *Glycinea*. FEMS Microbiol. Lett., 117: 1-6.
- Markovic, O., A. Slezarik and I. Labudova, 1985. Purification and characterization of pectinesterase and polygalacturonase from *Trichoderma reesei*. FEMS Microb. Lett., 27: 267-271.
- Martins, E.S., D. Silva, R. Da Silva and E. Gomes, 2002. Solid state production of thermostable pectinases from thermophilic *Thermoascus aurantiacus*. Process Biochem., 37: 949-954.
- McCarthy, R.E., S.F. Kotarski and A.A. Salyers, 1985. Location and characteristics of enzymes involved in the breakdown of polygalacturonic acid by *Bacteroides thetaiotaomicron*. J. Bacteriol., 161: 493-499.

- Micheli, F., 2001. Pectin methylesterase: cell wall enzymes with important roles in plant physiology. Trends Plant Sci., 6: 414-419.
- Mill, P.J., 1966a. The pectic, enzymes of *Aspergillus niger* mercury activated exopolygalacturonase. Biochem. J., 99: 557-561.
- Mill, P.J., 1966b. The peptic enzymes of *Aspergillus niger*: A second exopolygalacturonase. Biochem. J., 99: 562-565.
- Miller, G.L., 1959. Use of dinitrosalicylic acid reagent for determination of reducing sugar. Anal. Chem., 31: 426-428.
- Miyazaki, Y., 1991. Purification and characterization of an endo-pectate lyase from *Bacillus macerans*. Agric. Biol. Chem., 55: 25-30.
- Moharib, S.A., S.T. El-Sayed and E.W. Jwanny, 2000. Evaluation of enzymes produced by yeast. Nahrung, 44: 47-51.
- Morgavi, D.P., K.A. Beauchemin, V.L. Nsereko, L.M. Rode and A.D. Iwaasa *et al.*, 2000. Synergy between ruminal fibrolytic enzymes and enzymes from *Trichoderma longibrachiatum*. J. Dairy Sci., 83: 1310-1321.
- Murad, H.A and M.S. Foda, 1992. Production of yeast polygalacturonase on diary wastes. Bioresour. Technol., 41: 247-250.
- Murad, H.A., 1998. Utilization of ultrafiltration permeate for production of B. galactosidase from *Lactobacillus bulgaricus*. Milchwissenschaft, 53: 273-276.
- Murad, H.A. and M.M. Salem, 2001. Utilization of Uf-permeate for producing exo-polysaccharides from lactic acid bacteria. J. Agric. Mansoura Univ., 26: 2165-2173.
- Murad, H.H., M.A. Hanfy, A.M. Kholif, M.H. Abdel Gawad and H.A. Murad, 2009. Effect of cellulases supplementation to some low quality roughages on digestion and milk production by lactating goats. J. Biol. Chem. Environ. Sci., 4: 791-809.
- Nagai, M., T. Katsuragi, T. Terashita, K. Yoshikawa and T. Sakai, 2000. Purification and characterization of an endo-polygalacturonase from *Aspergillus awamori*. Biosci. Biotechnol. Biochem., 64: 1729-1732.
- Naidu, G.S.N. and T. Panda, 1998. Production of pectolytic enzymes: A review. Bioprocess Eng., 19: 355-361.
- Naumov, G.I., E.S. Naumova, M. Aigle, I. Masneuf and A. Belarbi, 2001. Genetic reidentification of the pectinolytic yeast strain SCPP as *Saccharomyces bayanus* var. *uvarum*. Applied Microbiol. Biotechnol., 55: 108-111.
- Nelson, N., 1944. A photometric adaptation of the Somogyi method for the determination of glucose. J. Biol. Chem., 153: 375-380.
- Nozaki, K., K. Miyairi, S. Hizumi, Y. Fukui and T. Okuno, 1997. Novel exopolygalacturonases produced by *Alternaria mali*. Biosci. Biotechnol. Biochem., 61: 75-80.
- Nsereko, V.L., D.P. Morgavi, L.M. Rode, K.A. Beauchemin and T.A. McAllister, 2000. Effects of fungal enzyme preparations on hydrolysis and subsequent degradation of alfalfa hay fiber by mixed rumen microorganisms *in vitro*. Anim. Feed Sci. Technol., 88: 153-170.
- Ooijskaas, L.P., F. Weber, R.M. Buitelaar, J. Tramper and A. Rinzema, 2000. Defined media and inert supports: Their potential as solid-state fermentation production system. Trends Biotechnol., 18: 356-360.
- Orpin, C.G., 1984. The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls. Anim. Feed Sci. Technol., 10: 121-143.

- Palomaki, T. and H.T. Saarilahti, 1997. Isolation and characterization of new C-terminal substitution mutation affecting secretion of polygalacturonases in *Erwinia carotovora* ssp. *carotovora*. FEBS Lett., 400: 122-126.
- Pandey, A., C.R. Soccol, J.A. Rodriguez-Leon and P. Nigam, 2001. Solid-State Fermentation in Biotechnology: Fundamentals and Applications. 1st Edn., Asiatech Publishers Inc., New Delhi, ISBN: 81-87680-06-7, pp: 221.
- Pereira, S.S., E.T. Torres, G.V. Gonzalez and M.G. Rojas, 1993. Effect of different carbon sources on the synthesis of pectinase by *Aspergillus niger* in submerged and solid state fermentation. Applied Microbiol. Biotechnol., 39: 36-41.
- Phutela, U., V. Dhuna, S. Sandhu and B.S. Chadha, 2005. Pectinase and polygalacturonase production by a thermophilic *Aspergillus fumigatus* isolated from decomposing orange peels. Braz. J. Microbiol., 36: 63-69.
- Piccoli-Valle, R.H., F.M.L. Passos, F.J.V. Passos and D.O. Silva, 2001. Production of pectin lyase by *Penicillium griseoroseum* in bioreactors in the absence of inducer. Braz. J. Microbiol., 32: 135-140.
- Pilnik, W. and A.G.J. Voragen, 1970. Pectic Substances and other Uronides. In: The Biochemistry of Fruits and their Products, Hulme, A.C. (Ed.). Academic Press, London, pp: 53-87.
- Pitkanen, K., R. Heikinheimo and R. Pakkanen, 1992. Purification and characterization of *Erwinia chrysanthemi* B374 pectin methylesterase produced by *Bacillus subtilis*. Enzyme Microbial. Technol., 14: 832-836.
- Polizeli, M.L.T.M., J.A. Jorge and H.F. Terenzi, 1991. Pectinase production by *Neurospora crassa*: purification and biochemical characterization of extracellular polygalacturonase activity. J. Gen. Microbiol., 137: 1815-1823.
- Prade, R.A., D. Zohan, P. Ayoubi and A.J. Mort, 1999. Pectins, pectinases and plant-microbe interactions. Biotechnol. Genet. Eng. Rev., 16: 361-391.
- Pressey, R. and J.K. Avants, 1973. Two forms of polygalacturonase in tomatoes. Biochim. Biophys. Acta, 309: 363-369.
- Rafae, R.J., G. Amin, H.A. Murad and H.M. Sahar, 2006. Exopolysaccharides production by dairy lactic acid bacteria. J. Agric. Sci. Mansoura Univ., 31: 5965-5998.
- Rasheedha, A.B., M.D. Kalpana, G.R. Gnanaprabhal, B.V. Pradeep and M. Palaniswamy, 2010. Production and characterization of pectinase enzyme from *Penicillium chrysogenum*. Indian J. Sci. Technol., 3: 377-381.
- Reda, A.B., M.Y. Hesham, A.S. Mahmoud and Z.A. Ebtsam, 2008. Production of bacterial pectinase(s) from agro-industrial wastes under solid state fermentation conditions. J. Applied Sci. Res., 4: 1708-1721.
- Reid, I. and M. Ricard, 2000. Pectinase in paper making, solving retention problems in mechanical pulps bleached with hydrogen peroxide. Enzyme Microbiol. Technol., 26: 115-123.
- Rexova-Benkova, L. and O. Markovic, 1976. Pectic enzymes. Adv. Carbohydrate Chem., 33: 323-385.
- Roboz, E., R.W. Barratt and E.L. Tatum, 1952. Breakdown of pectic substances by a new enzyme from *Neurospora*. J. Biol. Chem., 195: 459-471.
- Rodrigues-Palenzuela, P., T.J. Burr and A. Collmer, 1991. Polygalacturonase is a virulence factor in *Agrobacterium tumefaciens* biovar 3. J. Bacteriol., 173: 6547-6552.
- Sakai, T. and M. Okushima, 1982. Purification and crystallisation of a protopectin- solubilizing enzyme from *Trichosporon penicillatum*. Agric. Biol. Chem., 46: 667-776.

- Sakai, T. and A. Takaoka, 1984. Purification, crystallization and some properties of endo-polygalacturonase from *Aureobasidium pullulans*. Agric. Biol. Chem., 49: 449-458.
- Sakai, T. and Y. Ozaki, 1988. Protopectin solubilizing enzyme that does not catalyze the degradation of polygalacturonic acid. Agric. Biol. Chem., 52: 1091-1093.
- Sakai, T. and T. Sakamoto, 1990. Purification and some properties of a protopectin- solubilizing enzyme that has potent activity on sugar beet protopectin. Agric. Biol. Chem., 53: 1213-1223.
- Sakai, T., 1992. Degradation of Pectins. In: Microbial Degradation of Natural Products, Winkelmann, G. (Eds.). VCH, Weinheim, Germany, pp: 57-81.
- Sakai, T., T. Sakamoto, J. Hallaert and E.J. Vandamme, 1993. Pectin, pectinase and protopectinase: Production, properties and applications. Adv. Applied Microbiol., 39: 231-294.
- Sakamoto, T., R.A. Hours and T. Sakai, 1994. Purification, characterization and production of two pectic-transeliminases with protopectinase activity from *Bacillus subtilis*. Biosci. Biotechnol. Biochem., 58: 353-358.
- Sapunova, L.I., 1990. Pectinohydrolases from *Aspergillus alliaceus* Biosynthesis Characteristic Features and Applications. Institute of Microbiology Belarussian Academy of Science, Minsk.
- Sapunova, L.I., R.V. Mikhailova and A.G. Lobanok, 1995. Properties of pectin lyase preparations from the genus *Penicillium*. Applied Microbiol. Biochem., 31: 435-438.
- Sapunova, L.I., G. Lobanok and R.V. Mickhailova, 1997. Conditions of synthesis of pectinases and proteases by *Aspergillus alliaceus* and production of a complex macerating preparation. Applied Biotechnol. Microbiol., 33: 257-260.
- Sarvamangala, R.P. and A. Dayanand, 2006. Production of pectinase from deseeded sunflower head by *Aspergillus niger* in submerged and solid-state conditions. Bioresour. Technol., 97: 2054-2058.
- Savary, B.J., 2001. Perfusion chromatography separation of the tomato fruit-specific pectin methylesterase from a semi purified commercial enzyme preparation. Prep. Biochem. Biotechnol., 31: 241-258.
- Schell, M.A., T.P. Denny and J. Huang, 1994. Extracellular Virulence Factors of *Pseudomonas Solanacearum*: Role in Disease and their Regulation. In: Molecular Mechanisms of Bacterial Virulence, Kado, C.I. and J.H. Crosa (Eds). Kluwer Academic Press, The Netherlands pp: 311-324.
- Seethaler, D. and W. Hartmeier, 1992. Purification and properties of pectolytic enzymes of flocculent strain of *Clostridium acetobutylicum*. DECHEMA-Biotechnol. Conf., 5: 213-216.
- Selinger, L.B., C.W. Forsberg and K.J. Cheng, 1996. The rumen: A unique source of enzymes for enhancing livestock production. Anaerobe, 2: 263-284.
- Semenova, M.V., S.G. Grishutin, A.V. Gusakov, O.N. Okunev and A.P. Sinitsyn, 2003. Isolation and properties of pectinases from the fungus *Aspergillus japonicus*. Biochemistry, 68: 559-569.
- Serra, J.L., I. Alkorta, M.J. Llama and A. Alaiia, 1992. Aplicacion industrial de los enzimas pecticos. Production, purification, inmovilizacion y algunas propiedades de la pectina liasa de *Penicillium italicum*. Alimentacion Equiposy Tecnol., 11: 127-134.
- Shevchik, V.E, J. Robert-Baudouy and N. Hugouvieux-Cotte-Pattat, 1997. Pectate lyase Pel 1 of *Erwinia chrysanthemi* 3937 belongs to a new family. J. Bacteriol., 179: 7321-7330.
- Shin, I., S. Dowmeiz and O. Kilic, 1983. Study on pectolytic enzyme production from some agricultural wastes by fungi. Chem. Microbiol. Technol. Lebensm, 8: 87-90.
- Shoichi, T., K. Xoighi and S. Hiroshi, 1985. Cellulase production by *P. purpurogenum*. J. Ferment. Technol., 62: 127-127.

- Siebert, F.B. and J. Anto, 1946. Determination of polysaccharides in serum. J. Biol. Chem., 163: 511-522.
- Silley, P., 1985. A note on the pectinolytic enzymes of *Lachnospira multiparus*. J. Applied Bacteriol., 58: 145-149.
- Silva, D., E.D. Silva, R.D. Silva and E. Gomes, 2002. Pectinase production by *Penicillium viridicatum* RFC3 by solid state fermentation using agricultural wastes and agro industrial by-products. Braz. J. Microbiol., 33: 318-324.
- Singh, S.A., M. Ramakrishna and A.G.A. Rao, 1999. Optimization of downstream processing parameters for the recovery of pectinase from the fermented broth of *Aspergillus carbonarius*. Process Biochem., 35: 411-417.
- Souza, J.V.B., T.M. Silva, M.L.S. Maia and M.F.S. Teixeira, 2003. Screening of fungal strains for pectinolytic activity: endopolygalacturonase production by *Peacilomyces clavisporus* 2A.UMIDA.1. Process Biochem., 4: 455-458.
- Sunnotel, O. and P. Nigam, 2002. Pectinolytic activity of bacteria isolated from soil and two fungal strain during submerged fermentation. World J. Microbiol. Biotechnol., 18: 835-839.
- Szajer, I. and C. Szajer, 1982. Pectin lyase of *Penicillium paxilli*. Biotechnol. Lett., 4: 549-552.
- Szajer, I. and C. Szajer, 1985. Formation and release of pectin lyase during growth of *Penicillium paxilli*. Biotechnol. Lett., 7: 105-108.
- Takao, M., T. Nakaniwa, K. Yoshikawa, T. Terashita and T. Sakai, 2001. Molecular cloning, DNA sequence, and expression of the gene encoding for thermostable pectate lyase of thermophilic *Bacillus* sp. TS 47. Biosci. Biotechnol. Biochem., 65: 322-329.
- Tanabe, H., Y. Kobayashi, Y. Matuo, N. Nishi and F. Wada, 1984. Isolation and fundamental properties of exo pectate lyase p1 isozymes from *Erwinia carotovora*. Agric. Biol. Chem., 8: 2113-2120.
- Tierny, Y., M. Bechet, J.C. Joncquiert, H.C. Dubourguier and J.B. Guillaume, 1994. Molecular cloning and expression in *Escherichia coli* of genes encoding pectate lyase and pectin methylesterase activities from *Bacteroides thetaiotaomicron*. J. Applied Bacteriol., 76: 592-602.
- Torres, E.F., T.V. Sepulveda and G.V. Gonzalez, 2006. Production of hydrolytic depolymerising pectinases. Food. Technol. Biotechnol., 44: 221-227.
- Vaughan, I.L.H., T. Jakubczly, J.D. McMillan, T.E. Higgins, B.A. Dave and V.M. Crampton, 1969. Some pink yeast associated with softening of olives. Applied Microbiol., 18: 771-775.
- Vivek, R., M. Rajasekharan, R. Ravichandran, K. Sriganesh and V. Vaitheeswaran, 2010. Pectinase production from orange peel extract and dried orange peel solid as substrates using *Aspergillus niger*. Int. J. Biotechnol. Biochem., 6: 445-453.
- Ward, O.P. and M. Moo-Young, 1989. Enzymatic degradation of cell wall and related plant polysaccharides. CRC Crit. Rev. Biotechnol., 8: 237-274.
- Warrilow, A.G.S., R.J. Turner and M.G. Jones, 1994. A novel form of pectinesterase in tomato. Phytochemistry, 35: 863-868.
- Wattad, C., S. Freeman, A. Dinoor and D. Prusky, 1995. A nonpathogenic mutant of *Colletotrichum magna* is deficient in extracellular secretion of pectate lyase. Mol. Plant-Microbe Interac., 8: 621-626.
- Whitaker, J.R., 1984. Pectic substances, pectic enzymes and haze formation in fruit juices. Enzyme Microbial Technol., 6: 341-349.
- Whitaker, J.R., 1990. Microbial Pectinolytic Enzymes. In: Microbial Enzymes and Biotechnology, Fogarty, W.M. and C.T. Kelly (Eds.). 2nd Edn., Elsevier Science Ltd., London, pp: 133-176.



- Wijesundera, R.L.C., J.A. Bailey and R.J.W. Byrde, 1984. Production of pectin lyase by *Colletotrichum lindemuthionum* in culture and in infected bean (*Phaseolus vulgaris*) tissue. *J. Gen. Microbiol.*, 130: 285-290.
- Wu, Y.T., M. Pereira, A. Venoncio and J. Teixeira, 2000. Recovery of endopolygalacturonase using polyethylene glycol-salt aqueous two phase extraction with polymer recycling. *Bioseparation*, 9: 247-254.
- Yakoby, N., I. Kobiler, A. Dinoor and D. Prusky, 2000. pH regulation of pectate lyase secretion modulates the attack of *Colletotrichum gloeosporioides* in avocado fruits. *Applied Environ. Microbiol.*, 66: 1026-1030.
- Yang, W.Z., K.A. Beauchemin and L.M. Rode, 1999. Effects of an enzyme feed additive on extent of digestion and milk production of lactating dairy cows. *J. Dairy Sci.*, 82: 391-403.
- Zetelaki-Horvath, K., 1980. Factors affecting pectinase activity. *Acta Alimentaria*, 10: 371-378.
- Zheng, Z. and K. Shetty, 1999. Solid state production of polygalacturonase by *Lentinus edodes* using fruit processing wastes. *Process Biochem.*, 35: 825-830.