



Research Journal of
**Medicinal
Plant**

ISSN 1819-3455



Academic
Journals Inc.

www.academicjournals.com

Total Phenolic Content, Flavonoid Content and *in vitro* Antioxidant Activities of *Dendrophthoe falcata* (L.f.) Ettiingsh

¹S.P. Pattanayak, ¹P. Mitra Mazumder and ²P. Sunita

¹Division of Pharmacology, Department of Pharmaceutical Sciences, Birla Institute of Technology (BIT), A Deemed University u/s 3 of UGC Act, 1956, Mesra, Ranchi, Pin 835 215, India

²Department of Health and Family Welfare, Government Pharmacy Institute, Jharkhand Government, Bariatu, Ranchi, Pin 834 009, Jharkhand, India

*Corresponding Author: S.P. Pattanayak, Division of Pharmacology, Department of Pharmaceutical Sciences, Birla Institute of Technology (BIT), A Deemed University u/s 3 of UGC Act, 1956, Mesra, Ranchi, Pin 835215, India
Tel: +916512275444 Fax: +916512275209*

ABSTRACT

The Total Phenolic Content (TPC) and Total Flavonoid Content (TFC) of Chloroform (CEDF), Methanol (MEDF) and Hydroalcoholic (HEDF) extracts of an Indian ayurvedic plant *Dendrophthoe falcata* (L.f) Ettiingsh were measured using Folin-ciocalteau and aluminum chloride colorimetric methods, respectively. The *in vitro* antioxidant activities were also investigated by using *in vitro* antioxidant models including 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS⁺), 1,1-diphenyl-2-picrylhydrazyl (DPPH), Nitric Oxide (NO), hydroxyl radical (OH⁻), Super Oxide (SO) and reducing power (Fe³⁺ to Fe²⁺ transformation) assay. The TPC were 7.43±1.145, 10.783±0.702 and 19.833±0.550 mg gallic acid equivalents (GAE)/g extract, while TFC were 4.983±0.175, 9.346±0.610 and 14.933±1.033 µg quercetin equivalents (QRT)/g extract sample for CEDF, MEDF and HEDF, respectively. Analysis of the free radical scavenging activities of the fractions revealed a concentration-dependent antiradical activity resulting from reduction of ABTS⁺, DPPH, NO, OH⁻ and SO radicals to non-radical form. The scavenging activity of ascorbic acid, a known antioxidant used as positive control, was however higher and scavenging potential was in the order: Ascorbic acid>HEDF>MEDF>CEDF. The reducing power of ascorbic acid, CEDF, MEDF and HEDF increased gradually with increasing concentration. The order of the reduction potential was ascorbic acid>HEDF>MEDF>CEDF. These results obtained in the present study indicate that *D. falcata* extracts can be a potential source of natural antioxidant with strong antiradical capacity.

Key words: *Dendrophthoe falcata*, total phenolic content, total flavonoid content, antiradical activity, reducing power

INTRODUCTION

Oxidative stress depicts the existence of products called free radicals and Reactive Oxygen Species (ROS) which are formed under normal physiological conditions but become deleterious when not being eliminated by the endogenous systems. In fact, oxidative stress results from an imbalance between the generation of reactive oxygen species and endogenous antioxidant systems. ROS are major sources of primary catalysts that initiate oxidation *in vivo* and *in vitro* and create oxidative stress. These oxidative stress results in numerous diseases and disorders (Rackova *et al.*, 2007) such as cancer (Kinnula and Crapo, 2004), cardiovascular disease (Singh and Jialal, 2006),

Alzheimer's disease (Smith *et al.*, 2000), parkinsons disease (Bolton *et al.*, 2000) and atherosclerosis (Upston *et al.*, 2003). Oxygen derived free radicals such as superoxide anions, hydroxyl radicals and hydrogen peroxide are cytotoxic and give rise to tissue injuries (Jainu and Shyamala Devi, 2005). In addition, oxidative stress causes inadvertent enzyme activation and oxidative damage to cellular system (Pemp *et al.*, 2010).

Cells are equipped with different kinds of mechanisms to fight against ROS and to maintain the redox homeostasis of cell. For example, antioxidant enzymes such as Superoxide Dismutase (SOD), Catalase (CAT) and Glutathione Peroxidase (GPx) play important roles in scavenging the free radicals and preventing cell injury (Rakesh *et al.*, 2010). Molecules such as ascorbic acid and α -tocopherol inhibit lipid peroxydation in cell. When the mechanism of antioxidant protection becomes unbalanced in human body, antioxidant supplement may be used to help reduce oxidative damage.

Medicinal plants are an important source of antioxidants (Rice-Evans, 2004). Natural antioxidants increase the antioxidant capacity of the plasma and reduce the risk of certain diseases such as cancer, heart diseases and stroke (Prior and Cao, 2000). The secondary metabolites like phenolics and flavonoids from plants have been reported to be potent free radical scavengers. They are found in all parts of plants such as leaves, fruits, seeds, roots and bark (Mathew and Abraham, 2006). There are many synthetic antioxidants in use. It is reported, however, they have several side effects, such as risk of liver damage and carcinogenesis (Meenakshi *et al.*, 2009). There is a need for more effective, less toxic and cost effective antioxidants. Medicinal plants appear to have these desired comparative advantages, hence the growing interest in natural antioxidants from plants.

Dendrophthoe falcata (L.f) Ettingsh, commonly known as Banda (Hindi) is an evergreen hemiparasitic shrub with bark smooth grey, leaves opposite unequal, thick 1.6-25.4 cm long, flowers single, orange-red or scarlet softly pubescent, berries soft ovoid-oblong, 1.3 cm diameter and indigenous to India, Srilanka, Thailand, Indo-china, Australia. The tribal of India uses the aerial parts in wounds, menstrual troubles, asthma, psychic disorders, pulmonary tuberculosis, consumption and mania. Leaf paste is used in skin diseases and abortion. Its paste is applied on boils, setting dislocated bones and extracting pus. Traditional physicians of Korku, a tribe inhabiting the forest areas of Melghat region of Amravati district, Maharashtra state of India, use *D. falcata* as antifertility agent in women (Pattanayak *et al.*, 2008a). In addition to its medicinal value, the fruit of *D. falcata* tastes sweet and is consumed as a food (Mallavadhani *et al.*, 2006). The plant has been scientifically proved to have antilithiatic, diuretic, cytotoxic, immunomodulatory, wound healing and chemopreventive potentials (Pattanayak and Sunita, 2008; DKFZ, 2003). Recently our research group reported the effects of *Dendrophthoe falcata* in lowering the incidence of breast tumor growth in the experimental female wistar rats (Pattanayak *et al.*, 2008a; Pattanayak and Mazumder, 2010). The plant is also proved to have antifertility efficacy in wistar female rats (Pattanayak and Mazumder, 2009). The merit of the traditional use of *D. falcata* has also been reported by the isolation and identification of several possible active chemical constituents. The isolated compound from *D. falcata* are β -sitostirol, stigmasterol, kaempferol, quercetin-3-O-ramnoside, rutin, quercetin, myrecitin and their glycosides (+)-catechin, leucocyanidin, gallic acid, chebulinic acid (Pattanayak *et al.*, 2008b) and some pentacyclic triterpenes, kaempferol-3-O- α -l-rhamnopyranoside and quercetin-3-O- α -l-rhamnopyranoside, etc. (Mallavadhani *et al.*, 2006). It is evident from the above literature that *D. falcata* is therapeutically useful in oxidative stress induced diseases.

The aim of the present study was to evaluate the total phenolics content, flavonoid content of chloroform, methanol and hydroalcoholic extracts of *D. falcata* and to evaluate its *in vitro* antioxidant potential. Because different antioxidant compounds may act *in vivo* through different mechanisms, no single method can fully evaluate the total antioxidant capacity. For this reason, several complimentary test systems, including 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid), 1,1-diphenyl-2-picrylhydrazyl, nitric oxide, hydroxyl radical, super oxide and reducing power assay methods were used in this study.

MATERIALS AND METHODS

Plant material and extract preparation: Fresh aerial parts of *D. falcata*, were collected in the month of March in 2009 from the thick forest areas of Similipal biosphere reserve, Mayurbhanj district of Orissa, India. *Dendrophthoe falcata* (L.f) Ettingsh (Loranthaceae) was preliminarily authenticated by Dr. N.K. Dhal, Department of Natural products, Regional Research Laboratory (RRL), Bhubaneswar, India which was latter on confirmed from Botanical Survey of India, Hawrah, West Bengal, India (CNH/I-I/32/2010/Tech.II/237-2). One set of the herbarium has been preserved in our laboratory for future reference. The aerial parts were air-dried, pulverized to a coarse powder in a mechanical grinder, passed through a 40 mesh sieve and extracted with petroleum ether (60-80°C) and further with Chloroform (CEDF), Methanol (MEDF) and ethanol-water (8:2) (HEDF), separately. The extracts were decanted, filtered with Whatman No. 1 filter paper. The entire extraction process was repeated twice. The filtrates were individually pooled and solvents were removed from the filtrates under reduced pressure (Rotavapor RII, Buchi Labortechnik AG, Switzerland). Finally, *D. falcata* extracts were cooled in a dessicator for 30 min before the yield of each extract was calculated. Extracts from the aerial parts of *Dendrophthoe falcata* were kept at 4°C for further use.

Phytochemical screening: An attempt was also made to observe the presence and absence of different phytochemical constituents in CEDF/MEDF/HEDF. The tests for using the methods carried out flavonoid, sterols and tannins previously described by Tona *et al.* (1998). Two milligrams of each extract were separately dissolved in 2 mL of the adequate solvent. The detection of major chemical groups was carried out by Thin-layer Chromatography (TLC) on silica gel 60 F₂₅₄ from Merck (Dramstadt, Germany) (layer thickness, 0.25 mm). For flavonoid, TLC was developed in n-Butanol/acetic acid/water 4:1:5, then spots were visualized with 1% AlCl₃ solution in methanol under Ultraviolet (UV) 366 nm. Terpenes and steroids were detected with Libermann-Burchard as a reagent using n-hexane/CH₂Cl₂ 1:9 as a mobile phase. Ranges of colors are produced after heating sprayed plates for 10 min at 100°C. Tannins were detected with FeCl₃. Each class of tannins gave a specific coloration. And alkaloids (Dragendorff's test), glycosides (Molisch's reagent), fixed oil (spot test), proteins (Ninhydrine test), carbohydrates (Fehling's solution test) and polysaccharides (Iodine test) were detected according to standard methods (Soforowa, 1982; Trease and Evans, 1987).

Total phenolic content (Folin-Ciocalteu assay): Total phenolic contents of CEDF/MEDF/HEDF extracts were determined using Folin-Ciocalteu assay (Meda *et al.*, 2005).

Total flavonoid content: Total flavonoid content was determined by the aluminium calorimetric method (Quettier-Deleu *et al.*, 2000), using quercetin as a standard.

In vitro antioxidant assays

ABTS assay: The antioxidant potential was measured by 2,2'-azinobis - (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assay that measures the relative ability of antioxidant substances to scavenge the ABTS⁺ cation radical generated in the aqueous phase. 3.5 mL reaction mixture contained 0.17 mM ABTS, 25-250 µg mL⁻¹ CEDF/MEDF/HEDF/ascorbic acid and phosphate buffer (pH 7.4). The method used was based on Miller and Evans (1996) modified by Lister and Wilson (2001). The absorbance at 734 nm was measured using UV-vis spectrophotometer. The antioxidant capacities of samples were measured against the standard.

Determination of DPPH radical scavenging activity: The free radical scavenging activity was evaluated by the DPPH assay described by Blois (1958). In its radical form, DPPH absorbs at 517 nm but upon reduction by an antioxidant or a radical species, the absorption decreases. Briefly, 1 mL of 0.25 mM solution of DPPH in methanol was added to 1 mL of CEDF/MEDF/HEDF solution in methanol (25-250 µg mL⁻¹). After 20 min, the absorbance was measured at 517 nm. Ascorbic acid was used as a positive control. The percentage DPPH decolorization of the sample was calculated by the equation:

$$\% \text{ of DPPH scavenging} = \frac{(A_{\text{control}} - A_{\text{extract}})}{A_{\text{control}}} \times 100$$

where, A is the absorbance.

Nitric oxide radical scavenging (NO) assay: The nitric oxide radical inhibition activity was measured by the method of Garratt (1964) using Griess reagent.

Hydroxyl radical scavenging (OH) assay: Hydroxyl radical scavenging activity was determined by the method of Halliwell *et al.* (1987) based on the ability to compete with deoxyribose for hydroxyl radicals.

Superoxide radical scavenging (SO) assay: The super oxide anion scavenging activity was determined by the method of Nishikimi *et al.* (1972).

Ferric Reducing Antioxidant Power (FRAP) assay: The reductive potential was determined according to the method of Oyaizu (1986) based on the chemical reaction of Fe³⁺ to Fe²⁺.

Statistical analysis: The data are expressed as Mean±Standard Deviation (SD). All data were analysed by one-way Analysis of Variance (ANOVA) followed by Bonferroni's multiple comparison test (Sharma *et al.*, 2011) with equal sample size. The difference was considered significant when p-value<0.05.

RESULTS

Extraction yield: Figure 1 represents the yield of *D. falcata* extracts. The yield of the extracts varied from 8.94±0.164 to 20.493±0.469%. Among all the tested extracts the highest yield was obtained from HEDF (p<0.001). The yield of different extracts from the areal parts of *D. falcata* is presented in the following order: HEDF>MEDF>CEDF (p<0.01). Low extraction yield of the CEDF is probably due to the low solubility of the major components of the areal parts in chloroform.

Phytochemical screening: The results of the preliminary phytochemical screening of the HEDF revealed the presence of steroids, terpenes, glycosides, tannins, proteins, flavonoids, carbohydrates and polysaccharides. CEDF showed the positive results for steroids, terpenes, flavonoid and MEDF revealed the presence of steroids, tannins, terpenes, glycosides and flavonoids.

Total phenolic content and total flavonoid content: The content of phenolics compounds in the different extracts were determined through a linear gallic acid standard curve ($y = 0.133 + 0.019x$, $r^2 = 0.9980$). The total phenolic content of the extracts varied from 7.430 ± 1.145 to 19.833 ± 0.55 mg GAE/g extract. The highest content of total phenolics compounds was detected in HEDF (19.833 mg GAE/g extract) whereas the lowest content was measured in CEDF (7.43 mg GAE/g extract) ($p < 0.001$). Total phenolics content of the extracts is arranged in the following ascending order: CEDF < MEDF < HEDF ($p < 0.01$) (Table 1). In this study, the Total Flavonoid Content (TFC) of the extracts was evaluated by aluminium colorimetric assay. Quercetin (QRT) was used as a standard and the total flavonoid content of *D. falcata* extracts were expressed in microgram of quercetin equivalents per gram of extract ($\mu\text{g QRT/g extract}$) ($y = 0.005x + 0.156$, $r^2 = 0.9880$). The TFC of the extracts were varied from 4.983 ± 0.175 to 14.933 ± 1.033 $\mu\text{g WRT/g extract}$. The data presented in Table 1 indicates that the highest flavonoid content of

Table 1: Total phenolics content and total flavonoid content of *D. falcata* extracts (n = 3)

Samples (extracts)	Total phenolic content (mg GAE/g extract)	Total flavonoid content ($\mu\text{g QRT/g extract}$)
CEDF	7.430 ± 1.145	4.983 ± 0.175
MEDF	10.783 ± 0.702 a**	9.346 ± 0.61 a***
HEDF	19.833 ± 0.55 b***, c***	14.933 ± 1.033 b***, c***

Values are expressed as Mean \pm Standard Deviation (SD) from triplicate determination; a, b: MEDF and HEDF compared to CEDF; c: HEDF compared to MEDF; ** $p < 0.01$; *** $p < 0.001$




Fig. 1: Extraction yield of *D. falcata* extracts (n = 3); Data are represented as Mean \pm SD; * $p < 0.01$; ** $p < 0.001$




Fig. 2 (a-d): Inhibition of ABTS radical by (a) STD, (b) CEDF, (c) MEDF and (d) HEDF; Data are represented as Mean±SD of two independent experiments each

14.933 µg QRT/g extract was observed in the HEDF and the lowest content was observed in CEDF (4.983 µg QRT/g extract) ($p < 0.001$). TFC of the extracts is arranged in the following sequence: HEDF > MEDF > CEDF ($p < 0.01$).

In vitro antioxidant assay: Analysis of the free radical scavenging activities of the extracts revealed a concentration-dependent antiradical activity resulting from reduction of ABTS⁺ (Fig. 2), DPPH (Fig. 3), NO (Fig. 4), OH⁻ (Fig. 5) and SO (Fig. 6) radicals to non-radical form. The scavenging activity of ascorbic acid, a known antioxidant used as positive control, was however higher and scavenging potential was in the order: Ascorbic acid > HEDF > MEDF > CEDF. Figure 7 presents the reduction potential of CEDF, MEDF and HEDF. The reducing power of ascorbic acid, CEDF, MEDF and HEDF increased gradually with increasing concentration. The order of the reduction potential was ascorbic acid > HEDF > MEDF > CEDF.

DISCUSSION

These results suggest that the higher levels of antioxidant activity were due to the presence of phenolic components. These findings are in accordance with the earlier reports on total phenolic and antioxidant activity in roship extracts (Gao *et al.*, 2000). Phenols are very important plant constituents because of their scavenging ability due to their hydroxyl groups (Hatano *et al.*, 1989). The phenolic compounds may contribute directly to antioxidative action (Duh *et al.*, 1999). It is known that polyphenolic compounds have inhibitory effects on mutagenesis and carcinogenesis in humans when ingested up to 1 g daily from a diet rich in fruits and vegetables (Tanaka *et al.*, 1988). Phenolic compounds from plants are known to be good natural antioxidants. However, the activity of synthetic antioxidants was often observed to be higher than that of natural antioxidants




Fig. 3 (a-d): Inhibition of DPPH radical by (a) STD, (b) CEDF, (c) MEDF and (d) HEDF; Data are represented as Mean±SD of two independent experiments each

(Ningappa *et al.*, 2008). Phenolic compounds, at certain concentrations, markedly slowed down the rate of conjugated diene formation. The interests of phenolics are increasing in the food industry because they retard oxidative degradation of lipids and thereby improve the quality and nutritional value of food (Aneta *et al.*, 2007). The 2, 2'-azinobis (3-ethyl-benzothiazoline 6-sulfonate) (ABTS) formed from the reaction $ABTS \rightarrow ABTS^+$ reacts quickly with ethanol/hydrogen donors to form colorless ABTS. The reaction is pH-independent. A decrease of the $ABTS^+$ concentration is linearly dependent on the antioxidant concentration. All extracts at tested doses (25 - 250 µg mL⁻¹) revealed good scavenging activity for $ABTS^+$ in a dose dependent manner but the activity was higher in case of HEDF (IC₅₀ = 32.21 µg mL⁻¹) (Fig. 2). The DPPH radical is considered to be a model for a lipophilic radical. A chain in lipophilic radicals was initiated by the lipid autoxidation. DPPH is a stable free radical at room temperature and accepts an electron or hydrogen radical to become a stable diamagnetic molecule (Soares *et al.*, 1997). The reduction capability of DPPH was determined by the decrease in its absorbance at 517 nm which is induced by antioxidants. Positive DPPH test suggests that the samples were free radical scavengers. The scavenging effect of HEDF and ascorbic acid on DPPH radical was compared. On the DPPH radical, HEDF had significant scavenging effects with increasing concentration in the range of 25-250 µg mL⁻¹ and when compared with that of ascorbic acid, the scavenging effect of HEDF was lower. The IC₅₀ values were found to be 46.8, 81.4 and 100.07 µg mL⁻¹ for HEDF, MEDF and CEDF, respectively (Fig. 3). A higher DPPH radical-scavenging activity is associated with a lower IC₅₀ value. Nitric oxide plays an important role in various types of inflammatory processes in the body. In the present study the crude extracts of the *D. falcata* aerial parts (CEDF, MEDF and HEDF) were checked for its inhibitory effect on nitric oxide production. Nitric oxide radical generated from sodium nitroprusside at physiological pH was found to be inhibited by the extracts. The HEDF at varied concentrations




Fig. 4 (a-d): Inhibition of nitric oxide radical by (a) STD, (b) CEDF, (c) MEDF and (d) HEDF; Data are represented as Mean±SD of two independent experiments each

showed remarkable inhibitory effect of nitric oxide radical-scavenging activity compared to CEDF and MEDF (Fig. 4). Results showed the percentage of inhibition in a dose dependent manner for all the extracts tested. The concentration of HEDF needed for 50% inhibition (IC₅₀) was found to be 36.43 μg mL⁻¹, whereas, 92.87 and 98.43 μg mL⁻¹ was needed for MEDF and CEDF, respectively. The hydroxyl radical is an extremely reactive free radical formed in biological systems and has been implicated as a highly damaging species in free radical pathology, capable of damaging almost every molecule found in living cells (Hochstein and Atallah, 1988). This radical has the capacity to join nucleotides in DNA and cause strand breakage which contributes to carcinogenesis, mutagenesis and cytotoxicity. Hydroxyl radical scavenging capacity of an extract is directly related to its antioxidant activity (Babu *et al.*, 2001). The highly reactive hydroxyl radicals can cause oxidative damage to DNA, lipids and proteins (Trease and Evans, 1987). The effect of the extracts from *D. falcata* on the inhibition of free radical-mediated deoxyribose damage were assessed by means of the Iron (II)-dependent DNA damage assay. The Fentone reaction generates hydroxyl radicals (OH) which degrade DNA deoxyribose, using Fe²⁺ salts as an important catalytic component. Oxygen radicals may attack DNA either at the sugar or the base, giving rise to a large number of products. HEDF, MEDF and CEDF were also capable of reducing DNA damage at all concentrations used. Ascorbic acid was highly effective in inhibiting the oxidative DNA damage. As shown in Fig. 5, the extracts displayed potential inhibitory effect of hydroxyl radical-scavenging activity. All results showed hydroxyl radical scavenging activity in dose dependent manner. IC₅₀ values were found to be 41.31, 86.52 and 111.12 μg mL⁻¹ for HEDF, MEDF and CEDF, respectively (Fig. 5). The ability of the above mentioned extracts to quench




Fig. 5 (a-d): Inhibition of hydroxyl radical by (a) STD, (b) CEDF, (c) MEDF and (d) HEDF; Data are represented as mean±SD of two independent experiments each

hydroxyl radicals seems to be directly related to the prevention of propagation of the process of lipid peroxidation and seems to be good scavenger of active oxygen species, thus reducing the rate of the chain reaction. Ascorbic acid was used as reference standard. Hagerman *et al.* (1998) have also explained that high molecular weight and the proximity of many aromatic rings and hydroxyl groups are more important for the free radical-scavenging activity by phenolics than their specific functional groups. Super oxide is a reactive oxygen species which can cause damage to the cells and DNA leading to various diseases. It was therefore proposed to measure the comparative interceptive ability of the antioxidant extracts to scavenge the super oxide radical. Several *in vitro* methods are available for generation of super oxide radicals (Vani *et al.*, 1997). In our study super oxide radicals were generated by auto-oxidation of hydroxylamine in presence of NBT (Nitro blue tetrazolium). The reduction of NBT in presence of antioxidants was measured. The decrease of absorbance at 560 nm with antioxidants thus indicates the consumption of superoxide anion in the reaction mixture. As shown in Fig. 6, the all extracts at varied concentrations had superoxide-scavenging activity. IC₅₀ values of CEDF, MEDF, HEDF and ascorbic acid were found to be 144.19, 105.38 and 69.97 µg mL⁻¹, respectively. All of the extracts had a scavenging activity on the superoxide radicals in a dose dependent manner (25-250 µg mL⁻¹ in the reaction mixture). Nonetheless, when compared to ascorbic acid, the superoxide scavenging activity of the extract was found to be low. Generally, the reducing properties are associated with the presence of compounds which exert their action by breaking the free radical chain by donating a hydrogen atom (Duh *et al.*, 1999). The results of the ferric reducing assay indicated that HEDF had stronger reducing power than CEDF and MEDF (Fig. 7). This could be due to the presence of more reactive concentration of bioactive constituents and mixture of other constituents in HEDF than other two extracts.




Fig. 6 (a-d): Inhibition of super oxide radical by (a) STD, (b) CEDF, (c) MEDF and (d) HEDF; Data are represented as Mean±SD of two independent experiments each




Fig. 7: The reduction potential of CEDF, MEDF and HEDF (Mean±SD; n = 6); *, **, ***: significantly different from HEDF, MEDF and CEDF (p<0.05), (p<0.01), (p<0.001), respectively. a, b, c: HEDF significantly different from MEDF, CEDF (p<0.05), (p<0.01), (p<0.001), respectively

CONCLUSION

In the present study, there exists a positive correlation between the total phenolics content and the antioxidant activity which is in accordance with the earlier findings (Madsen *et al.*, 1996). We found higher *in vitro* antioxidant activity in HEDF with higher polyphenols. The higher radical scavenging efficacy of HEDF may be due to retention of antioxidant phytochemicals in this extract. A strong evidence supports these findings that HEDF enhance the antioxidant effects *in vivo* rodent models of carcinogenesis (Pattanayak and Mazumder, 2010). Moreover, these results suggest that *Dendrophthoe falcata* may offer effective protection from free radicals and support that *D. falcata*, is a promising source of natural antioxidant. However, further work is required on the isolation and identification of the antioxidant components present in it since it is also a precondition for a more extensive understanding of the mechanisms involved in the antioxidant capacity.

ACKNOWLEDGMENT

Authors are grateful to Birla Institute of Technology, Mesra, Ranchi for their financial support for the completion of this piece of research. Authors also express sincere gratitude to Department of Pharmaceutical Sciences and Department of Central Instrumentation Facility, BIT, Mesra for the infrastructural facilities provided to carry out this study.

REFERENCES

- Aneta, W., O. Jan and C. Renata, 2007. Antioxidant activity and phenolic compounds in 32 selected herbs. *Food Chem.*, 105: 940-949.
- Babu, B.H., B.S. Shylesh and J. Padikkala, 2001. Antioxidant and hepatoprotective effect of *Acanthus ilicifolius*. *Fitoterapia*, 72: 272-277.
- Blois, M.S., 1958. Antioxidant determinations by the use of a stable free radical. *Nature*, 181: 1199-1200.
- Bolton, J.L., M.A. Trush, T.M. Penning, G. Dryhurst and T.J. Monks, 2000. The role of quinones in toxicology. *Chem. Res. Toxicol.*, 13: 135-160.
- DKFZ, 2003. Research Report 2001/2002: Division of Toxicology and Cancer Risk Factors (C0200/C010). German Cancer Research Centre, Heidelberg.
- Duh, P.D., Y.Y. Tu and G.C. Yen, 1999. Antioxidative activity of water extracts of Hamg jyu (*Chrysanthemum morifolium*). *Lebensmittel-Wissenschaft Technol.*, 32: 269-277.
- Gao, X., M. Ohlander, N. Jeppsson, L. Bjork and V. Trajkovski, 2000. Changes in antioxidant effects and their relationship to phytonutrients in fruits of sea buckthorn (*Hippophae rhamnoides* L.) during maturation. *J. Agric. Food Chem.*, 48: 1485-1490.
- Garratt, C.J., 1964. Effect of iodination on the biological activity of insulin. *Nature*, 201: 1324-1325.
- Hagerman, A.E., K.M. Riedel, G.A. Jones, K.N. Sovik, N.T. Ritchard, P.W. Hartzfeld and T.L. Riechel, 1998. High molecular weight plant polyphenolics (tannins) as biological antioxidants. *J. Agric. Food Chem.*, 46: 1887-1892.
- Halliwell, B., J.M.C. Gutteridge and O.I. Aruoma, 1987. The deoxyribose method: A simple test-tube assay for determination of rate constant for reactions of hydroxyl radicals. *Anal. Biochem.*, 165: 215-219.
- Hatano, T., R. Edamatsu, M. Hiramatsu, A. Mori and Y. Fujita *et al.*, 1989. Effects of the interaction of tannins with co-existing substances. VI: Effects of tannins and related polyphenols on superoxide anion radical and on 1,1-diphenyl-2-picrylhydrazyl radical. *Chem. Pharm. Bull.*, 37: 2016-2021.

- Hochstein, P. and A.S. Atallah, 1988. The nature of oxidant and antioxidant systems in inhibition of mutation and cancer. *Mutat. Res.*, 202: 363-375.
- Jainu, M. and C.S. Shyamala Devi, 2005. *In vitro* and *in vivo* evaluation of free radical scavenging potential of *Cissus quadrangularis*. *Afr. J. Biomed. Res.*, 43: 773-779.
- Kinnula, V.L. and J.D. Crapo, 2004. Superoxide dismutases in malignant cells and human tumors. *Free Rad. Biol. Med.*, 36: 718-744.
- Lister, E. and P. Wilson, 2001. Measurement of Total Phenolics and ABTS Assay for Antioxidant Activity (Personal Communication). Crop Research Institute, Lincoln, New Zealand.
- Madsen, H.L., B.R. Nielsen, G. Bertelsen and L.H. Skibsted, 1996. Screening of antioxidative activity of spices. A comparison between assays based on ESR spin trapping and electrochemical measurement of oxygen consumption. *Food Chem.*, 57: 331-337.
- Mallavadhani, U.V., K. Narashimhan, A.V. Sudhakar, A. Mahapatra, W. Li and R.B. van Breemen, 2006. New pentacyclic triterpenes and some flavanoids from the fruits of Indian ayurvedic plant *Dendrophthoe falcate* and their receptor binding activity. *Chem. Pharm. Bull.*, 54: 740-744.
- Mathew, S. and T.E. Abraham, 2006. *In-vitro* antioxidant activity and scavenging effect of *Cinnamomum verum* leaf extract assayed by different methodologies. *Food. Chem. Toxicol.*, 44: 198-206.
- Meda, A., C.E. Lamien, M. Romito, J. Millogo and O.G. Nacoulma, 2005. Determination of the total phenolic, flavonoid and proline contents in Burkina Faso honey, as well as their radical scavenging activity. *Food Chem.*, 91: 571-577.
- Meenakshi, S., D.M. Gnanambigai, S.T. Mozhi, M. Arumugam and T. Balasubramanian, 2009. Total flavanoid and *in vitro* antioxidant activity of two seaweeds of rameshwaram coast. *Global J. Pharmacol.*, 3: 59-62.
- Miller, N.J. and C.R. Evans, 1996. Spectrophotometric determination of antioxidant activity. *Redox Rep.*, 2: 161-171.
- Ningappa, M.B., R. Dinesha and L. Srinivas, 2008. Antioxidant and free radical scavenging activities of polyphenol-enriched curry leaf extract (*Murraya koenigii* L.). *Food Chem.*, 106: 720-728.
- Nishikimi, M., N.A. Rao and K. Yagi, 1972. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochem. Biophys. Res. Commun.*, 46: 849-854.
- Oyaizu, M., 1986. Studies on products of browning reactions: Antioxidant activities of products of browning reaction prepared from glucosamine. *Jap. J. Nutr.*, 44: 307-315.
- Pattanayak, S.P. and P. Sunita, 2008. Wound healing, anti-microbial and antioxidant potential of *Dendrophthoe falcate* (L.f) Ettingsh. *J. Ethnopharmacol.*, 120: 241-247.
- Pattanayak, S.P., P. Sunita, P.M. Mazumder and P.K. Panda, 2008a. Evaluation of the antitumor activity of the aqueous extract of *Dendrophthoe falcate* on 7,12-dimethylbenz(a)anthracene-induced rat mammary tumor model. *Int. J. Biol. Sci.*, 2: 75-80.
- Pattanayak, S.P., P.M. Mazumder and P. Sunita, 2008b. *Dendrophthoe falcata* (L.f) ettingsh: A consensus review. *Pharmacog. Rev.*, 2: 359-368.
- Pattanayak, S.P. and P.M. Mazumder, 2009. Effect of *Dendrophthoe falcata* (L.f) Ettingsh on female reproductive system in Wistar rats: A focus on antifertility efficacy. *Contraception*, 80: 314-320.

- Pattanayak, S.P. and P.M. Mazumder, 2010. Therapeutic potential of *Dendrophthoefalcata* (L.f.) Ettingsh on 7,12-dimethylbenz(a)anthracene-induced mammary tumorigenesis in female rats: Effect on antioxidant system, lipid peroxidation and hepatic marker enzymes. *Comp. Clin. Pathology.*, (In press).
- Pemp, B., E. Polska, K. Karl, M. Lasta and G. Garhofer *et al.*, 2010. Effects of antioxidants (areds medication) on ocular blood flow and endothelial function in an endotoxin-induced model of oxidative stress in humans. *Invest. Ophthalmol. Visual Sci.*, 51: 2-6.
- Prior, R.L. and G. Cao, 2000. Antioxidant phytochemicals in fruits and vegetables: Diet and health implications. *Hortic. Sci.*, 35: 588-592.
- Quettier-Deleu, C., B. Gressier, J. Vasseur, T. Dine and C. Brunet *et al.*, 2000. Phenolic compounds and antioxidant activities of buckwheat (*Pagopyrum esculentum* moench), hulls and flour. *J. Ethnopharmacol.*, 72: 35-42.
- Rackova, L., M. Oblozinsky, D. Kostalova, V. Kettmann and L. Bezakova, 2007. Free radical scavenging activity and lipoxygenase inhibition of *Mahonia aquifolium* extract and isoquinoline alkaloids. *J. Inflammation*, 4: 15-22.
- Rakesh, S.U., P.R. Patil and S.R. Mane, 2010. Use of natural antioxidants to scavenge free radicals: A major cause of diseases. *Int. J. Pharm Tech. Res.*, 2: 1074-1081.
- Rice-Evans, C., 2004. Flavonoids and isoflavones: Absorption, metabolism and bioactivity. *Free Rad. Biol. Med.*, 36: 827-828.
- Sharma, P.V., R. Paliwal and S. Sharma, 2011. *In vitro* free radical scavenging and antioxidant potential of ethanolic extract of *Euphorbia neriifolia* Linn. *Int. J. Pharmacy Pharm. Sci.*, 3: 238-242.
- Singh, U. and I. Jialal, 2006. Oxidative stress and atherosclerosis. *Pathophysiol.*, 13: 129-142.
- Smith, M.A., C.A. Rottkamp, A. Nunomura, A.K. Raina and G. Perry, 2000. Oxidative stress in Alzheimer's disease. *Biochem. Biophys. Acta*, 1502: 139-144.
- Soares, J.R., T.C. Dinis, A.P. Cunha and L.M. Almeida, 1997. Antioxidant activities of some extracts of *Thymus zygis*. *Free Radic. Res.*, 26: 469-478.
- Soforowa, E.A., 1982. *Medicinal Plants and Traditional Medicine in Africa*. John Wiley and Sons, Chichester.
- Tanaka, M., C.W. Kuie, Y. Nagashima and T. Taguchi, 1988. Application of antioxidative Maillard reaction products from histidine and glucose to sardine products. *Bull. Jap. Soc. Scientific Fisheries*, 54: 1409-1414.
- Tona, L., K. Kambu, N. Ngimbi, K. Cimanga and A.J. Vlietinck, 1998. Antiamoebic and phytochemical screening of some congolese medicinal plants. *J. Ethnopharmacol.*, 61: 57-65.
- Trease, G.E. and W.C. Evans, 1987. *A Text Book of Pharmacognosy*. ELBS/Bailliere Tindal, Oxford, UK., pp: 1055.
- Upston, J.M., L. Kritharides and R. Stocker, 2003. The role of vitamin E in atherosclerosis. *Prog. Lipid Res.*, 42: 405-422.
- Vani, T., M. Rajani, S. Sarkar and C.J. Shishoo, 1997. Antioxidant properties of the ayurvedic formulation Triphala and its constituents. *Int. J. Pharm.*, 35: 313-317.