



Research Journal of
Phytochemistry

ISSN 1819-3471



Academic
Journals Inc.

www.academicjournals.com

Flavonoids of *Limoniastrum feei**

^{1,2}N. Belboukhari and ¹A. Cheriti

¹Phytochemistry and Organic Synthesis Laboratory,
University of Bechar, 08000 Bechar, Algeria

²Faculty of Sciences, Université de Paul Cézanne,
St-Jerome Av-escadrille N-Niemen F-13397 Marseille Cedex 20, France

Abstract: Phytochemistry investigation of the water-acetone extract of twig part of *Limoniastrum feei* led to isolation of four flavonoids. The structures of these compounds were identified as 6,3',4'-Tri-methoxy 3,5,5'-trihydroxy flavonol (1), 3-(6"-malonyl 2"-ramnosyl glucosil) 6,3',4'-tri-methoxy 5,5'-dihydroxy flavonol (2), Tetraacetate 7-dihydroxy-4'-Methoxy 8-O-β-glucopyranoside isoflavone (3) and Tetraacetate 7,4'-diMethoxy 8-O-β-glucopyranoside isoflavone (4) using spectroscopic analysis.

Key words: *Limoniastrum feei*, Plumbagenaceae, flavonol, flavonol glycoside, isoflavone

INTRODUCTION

One of the medicinal plants used to treat gastric infections is *Limoniastrum feei* (Plumbagenaceae). The plant is native to southeast of Algeria (Saoura, region of Bechar) northern Africa (Ozenda, 1983; Maire, 1953; Cheriti *et al.*, 2004).

The other uses of *Limoniastrum feei* are as an antibacterial, for treatment bronchitis, stomach infection (Cheriti, 2000). A previous investigations revealed that methanol extract from *Limoniastrum feei* leaves contained potential antifungal agent against *C. albican* and antibacterial agent against *E. coli* (Belboukhari *et al.*, 2005).

In this study, we describe the isolation of four flavonoids from *Limoniastrum feei* as well as the elucidation of their structures using spectroscopic analysis.

MATERIALS AND METHODS

General Experimental Procedure

UV spectra were obtained in MeOH solvent with UNICAM UV300 spectrophotometer. IR spectra were obtained with a AVATAR 320 FT-IR spectrophotometer. The NMR spectra were taken on a Bruker GP 250 (¹H, 250 MHz; ¹³C, 125 MHz) Spectrometer. EIMS spectra were obtained on a VG Trio-2 spectrometer. TLC was carried out on silica gel 60 F₂₅₄ plates (Merck, Germany). Column chromatography was performed over silica gel 60 (Merck, particle size 230-400 mesh).

Plant Materials


The whole plants of *Limoniastrum feei* were collected in March 2000 from kenadsa: (region of Bechar) Algeria. The botanical identification and a voucher specimen is conserved at the Phytochemical Herbarium of Phytochemistry and Organic Synthesis Laboratory of University Center of Bechar under to accession number CA99/14 (Belboukhari *et al.*, 2005). The leave, stem and twig were separated and dried, the twig part of plants were grounded into powder from using the grinder.

Corresponding Author: N. Belboukhari, Phytochemistry and Organic Synthesis Laboratory, University of Bechar, 08000 Bechar, Algeria Tel: 0668299305 Fax: 00213 49 81 52 44

*Originally Published in Research Journal of Phytochemistry, 2007

Extraction and Isolation


The dried twig part of plants (100 g) of *Limoniastrum feei* were extracted with acetone-water (70:30) using soxhlet apparatus, reflux for 3 h was performed. The residue was evaporated in vacuo apparatus until two third, the third of aqueous residue was partitioned sequentially with n-hexan, ethyl ether, EtOAc and n-BuOH (Hostettman *et al.*, 1998; Hamburger and Cordell, 1987). To purify and to identify the constituents of the fraction butanol (1.4 g) one achieved some separations by liquid chromatography on column, one using a column in glass of type, 20/300 mm (29/39) full with a stationary phase of silica gel (0.20 mm) and the mobile phase chosen for this separation is: Acetone/Toluene/Formic Acid. (60:80:10) (Belboukhari and Cheriti, 2006), the compound 1 correspond to the fraction 4 (8 mg) and the compound 2 with (56 mg) appears in the Fractions (32-55), the compound 3 (17 mg) in two fractions (56-57) and the compound 4 (87 mg) in fractions (70-97).



RESULTS AND DISCUSSION

Phytochemical investigation of twig part of *Limoniastrum feei* led to isolation of four flavonoids from the butanol fraction using column chromatography.

According to the results of IR analysis of all compounds, the absorption bands observed toward (3348-3448 cm^{-1}) correspond to the vibration of elongation $\hat{\text{O}}\text{-H}$ (valence vibration), the aliphatic links C-H is presented in the IR specter by fine and intense bands toward the 2951 cm^{-1} (asymmetric valance vibration of the CH_3). The frequency of vibrations situated between (2918-2934 cm^{-1}) corresponds to it: asymmetric valance vibration of the CH_2 , the bands of absorption to 2852 cm^{-1} associate has vibrations of symmetrical valence of the CH_2 (Cheriti, 2005). The frequency of vibration (989.98 cm^{-1}) corresponds to the distortion vibration out of the plan of the unsaturated hydrocarbons. The valence vibration of the cyclic ketones to six linkages or more, or aliphatic ketones ($\text{C}=\text{O}$) to be located toward 1716 cm^{-1} , one specified little that has 1738 cm^{-1} , the vibration are associate has some feature carbonyl for a saturated ester. The vibrations have 1765 and 1771 cm^{-1} corresponds to the ester of alcohol vinyl that possess the fragment of structure following:



this grouping entails an increase important of the frequency of vibration carbonyl (vinyl acetate absorb to 1776 cm^{-1} and the phenyl acetate absorb to 1770 cm^{-1}). The vibration of distortion outside of the plan of aromatic C-H depends mainly of the position of the various substituting to fix on the benzene and not of their nature (Harborne *et al.*, 1986).

Compound 1 and 2 was obtained as yellow amorphous powders. The compound 1 is a flavonol aglycone and represent the skeleton of basis of the compound 2, UV spectra of 1 and 2 in methanol showed λ_{max} 286, 312 for compound 1 and 281, 296, 334 for compound 2. the shift in the λ_{max} on the addition of diagnostic reagents confirmed the presence of free hydroxyl groups, suggesting that compound 1 are hydroxyl group in position 3 and 2 are each glycosylated at the 3-position of the aglycone. the results of ^1H NMR indicate the presence of three methoxyl groups in 7, 3', 4' position. The ^{13}C NMR spectra of 1 and 2 were very similar, except for the signal corresponding to the glycosyl group. The characteristic carbonyl carbon signals of a malonyl group were observed at δ 168.23 and 169.96 for compound 2 (Gohar, 2002; Mabry *et al.*, 1970; Bacon and Mabry, 1976). Compound 3 was a viscous gum and showed phenol characteristics in positive reaction with FeCl_3 reagents, it has the quasi molecular formula $\text{C}_{30}\text{H}_{31}\text{O}_{14}$, calculated by EIMS and ^{13}C NMR data, consistent with its $[\text{M}+\text{H}]^+$ at m/z 615. The UV spectrum with absorption bands at 211, 254, 306 nm suggested an isoflavone skeleton. The ^1H NMR spectrum displayed a characteristic singlet at δ_{H} 7.85 for H-2 of an isoflavone, two doublets with ortho coupling constant 9.0 Hz at δ_{H} 8.05 and 7.05 for H-5 and H-6, respectively and a pair of doublets of a p-disubstituted phenyl at δ_{H} 7.47 and 6.98. The methoxy group at δ_{H} 3.84 was assigned to C-4' in phenyl as irradiation of the aromatic protons (H-3', H-5') at δ_{H} 6.98. In addition, the proton signals ascribable to the glucose unit were observed together with the methyl protons of four acetyl groups (δ_{H} 2.20, 2.12, 2.07 and 2.06), indicating the presence of glucose as tetra-acetate derivative. The results of displacement proton at δ_{H} 4.99 and corresponding carbon at 131.1 indicated that the glucose unit was attached to the 8-hydroxyl of the isoflavone moiety.

There for, 3 was characterized as 7,8-dihydroxy-4'-Methoxy 8-O- β -glucopyranosyl isoflavone. And was confirmed by comparison to previously reported data (Rukachaisirikul *et al.*, 2002). The molecular formula of compound 4 was determined to be $\text{C}_{41}\text{H}_{47}\text{O}_{20}$ based on the ^{13}C NMR spectral data and EIMS [m/z 858, M^+]. A positive reaction with in FeCl_3 reagent also displayed the phenol characteristics of this compound. ^1H and ^{13}C NMR spectrums were similar to those of compound 3, except absence of hydroxyl group at the 7-position and presence a second unit in substituting glucosyl. The ^1H NMR suggested the presence of two methoxyl groups at δ_{H} 3.84 and 3.99, respectively in C-4' and C-7 positions. Comparison of its ^1H NMR spectrum of compound 3 indicated that compound 4 also contained β -glucopyranosyl and α -rhamnopyranosyl residues with anomeric protons at δ_{H} 5.22 (6.8 Hz) and 4.56 (1.6 Hz) which were attributed to H-1'' of the glucose unit and H-1''' of the rhamnose unit, respectively. The presence of an α -rhamnosyl residue was confirmed by a methyl doublet at δ_{H} 1.11 ($J = 6.3\text{ Hz}$). Therefore the compound 4 has been identified to: 8-hydroxy 4',7-dimethoxyisoflavone 8-O-[α -rhamnopyranosyl-(1-6)]- β -glucopyranoside (Markham and Geiger, 1994).

6,3',4'-Tri-methoxy 3,5,5'-trihydroxy flavonol (1)

$R_f = 0.8$, UV (MeOH): maxima at 286 and 312 nm, IR (KBr): 3410, 2934, 2847, 1689, 1558, 1430, 1377, 1115, 1033, 771 cm^{-1} . ^1H NMR: 6.70(H-6), 6.84(H-8), 7.35(H-2'), 6.93(H-6'), 3.12, 3.05, 3.23 (O- CH_3).

3-(6"-malonyl 2"-rhamnopyranosyl glucosil) 6,3',4'-tri-methoxy 5,5'-dihydroxy flavonol (2)

$R_f = 0.6$, UV (MeOH): 281, 296, 334, IR (KBr): 3393 (OH), 2951, 2923, 2862 (CH_3, CH_2), 1711 (CH_3COO), 1640 (C=O, C-4), 1601 (C=C) 1514 (arom), 1028, 1121 (C-O), ^1H NMR: 6.70(H-6), 6.84(H-8), 7.20(H-2'), 6.93(H-6'), 3.117, 3.057, 3.229 (O- CH_3), 3-glycosil: 5.55(H-1), 3.697(H-2),

3.57(H-3), 3.397(H-4), 3.292(H-5), 3.801(H-Ga), 3.397(H-G-b). 2"-ramnosyl: 5.21(H-1), 4.702(H-2), 3.729(H-3), 3.397(H-4), 4.1(H-5), 1.03(H-6, CH₃) 1.912, 1.988, 2.086, 2.177, 2.235, 2.027 (CH₃COO), ¹³C NMR: 162.24(C-2), 129.9(C-3), 174.06(C-4), 151.67(C-5), 129.9 (C-6), 140.1(C-7), 99.79 (C-8), 158.44 (C-9), 105.91(C-10), 122.53 (C-1'), 109.14 (C-2'), 140.1 (C-3'), 145.81 (C-4'), 137.81(C-5'), 109.14(C-6'), 47.77, 48.11, 48.44, 48.78 (CH₃-O), 3-glucosyl: 100.37(C-1), 79.17(C-2), 77.23(C-3), 71.49(C-4), 73.65(C-5), 63.37(C-6) 2"-ramnosyl: 102.62(C-1), 72.68(C-2), 72.35(C-3), 75.07(C-4), 70.43(C-5), 18.3 (C-6), 6"-malonyl: 41.4(CH₂), 28.99, 30.07, 34.9, 38.68, (CH₃COO).

Tetraacetate 7-dihydroxy-4'-Methoxy 8-O-β-glucopyranoside isoflavone (3)

R_f = 0.7, UV(MeOH): 211, 254, 306, IR(KBr): 3448, 2918, 2852, 1727, 1590, 1519, 1170, 1022, 1465, 716 cm⁻¹. ¹H NMR: 7.85(s, H-2), 8.05(d, 9.0, H-5), 7.05(d, 9.0, H-6), 7.47(d, 9.0, H-2', H-6'), 6.98(d, 9.0, H-3', H-5'), 4.99(d, 8.1, H-1''), 5.4(dd, 9.8, 8.1, H-2''), 5.33 (t, 9.8, H-3''), 5.2 (t, 9.8, H-4''), 3.82(ddd, 9.8, 5.6, 2.8, H-5''), 4.3(dd, 12.6, 5.6, H-6''), 3.84(s, 4'-OMe), 2.20, 2.12, 2.07, 2.06 (CH₃CO). ¹³C NMR: 151.3 (C-2), 125.1 (C-3), 175.6 (C-4), 118.7 (C-10), 124.3 (C-5), 115.3 (C-6), 154.4 (C-7), 131.1 (C-8), 150.2(C-9), 123.5 (C-1'), 130.2 (C-2', C-6'), 114.1(C-3', C-5'), 159.9(C-4'), 103.4 (C-1''), 71.0 (C-2''), 72.1 (C-3''), 67.9 (C-4''), 72.7 (C-5''), 61.3(C-6''), 55.4 (4'-OMe), 20.6, 20.7 (CH₃CO), 17.6, 170.1, 169.3, 169.2(COCH₃).

Hexacetate 8-Hydroxy-4',7-dimethoxy iso-flavone 8-O-[α-rhamnopyranosyl-(1-6)]-β-glucopyranoside, (4)

R_f = 0.4, C₄₁H₄₇O₂₀, [M+H]⁺ a m/z = 859, le spectre UV(MeOH) : 212, 253, 306 nm. IR (KBr): 3404, 3235, 2918, 2841, 1629, 1514, 1377, 1115, 1039 cm⁻¹. ¹H NMR: 8.25(s, H-2), 8.00(d, 9.0, H-5), 7.50 (d, 8.5, H-2', H-6'), 7.27(d, 9.0, H-6), 6.99(d, 8.5, H-3', H-5'), 5.07(d, 7.5, H-1''), 4.56(d, 1.5, H-1'''), 4.03(s, OMe), 3.82 (s, Ome) 3.84-3.37 (m, glucose and rhamnose protons), 1.06(d, 6.5, H-6'''). ¹³C NMR: 152.7(C-2), 124.7(C-3), 176.0(C-4), 119.4(C-10), 123.4(C-5), 110.3(C-6), 156.1(C-7), 132.4(C-8), 150.8(C-9), 124.2(C-1'), 130.5(C-2', C-6'), 114.2(C-3', C-5'), 159.9(C-4'), 101.2(C-1''), 72.1(C-2''), 72.9(C-3''), 69.2(C-4''), 74.4(C-5''), 66.5(C-6''), 97.9(C-1'''), 69.6(C-2'''), 69.3(C-3'''), 70.9(C-4'''), 66.9(C-5'''), 17.5(C-6'''), 56.8(7-OMe), 55.6(4'-OMe), 21.0, 20.9, 20.9, 20.8 (CH₃CO), 170.5, 170.3, 170.2, 170.1, 169.9, 169.6 (COCH₃).

REFERENCES

- Bacon, J.D. and T.J. Mabry, 1976. UV spectral procedures for distinguishing free and substituted 7-hydroxyl groups in flavones and flavonols. *Revista Latinoamericana De Quimica*, 7: 83-86.
- Belboukhari, N. Cheriti and A. Asian, 2005. Antimicrobial activity of aerial part crude extracts from *Limoniastrum feei*. *J. Plant Sci.*, 4: 496-498, 12.
- Belboukhari, N. and A. Cheriti, 2006. Phytochemical investigation of the bioactive extract from *Launaea arborescens*. *Pak. J. Biol. Sci.*, 9: 2930-2932.
- Cheriti, A., 2000. Report of the CRSTRA Project, Medicinal Plants of the Region of Bechar, (Ethnopharmacologie studies), Bechar: Algeria 2000.
- Cheriti, A., N. Belboukhari and S. Hacini, 2004. Ethnopharmacological survey and phytochemical screening of some medicinal plants of Algerian sahara, Iran. *J. Pharm. Res.*, pp: 51.
- Cheriti, A., N. Belboukhari and D. Hacini, El Abed, 2005. Medical plants of the region of Bechar: Contents in flavonoids and antibacterial assessment. *International Seminar on the Valorization of Medicinal Plants in the Arid Zones, Ouargla*, pp: 12.
- Gohar, A.A., 2002. Flavonol glycosides from *Cadaba glandulosa* Z. *Naturforsch*, 57c: 216.
- Hamburger, M.O. and G.A. Cordell, 1987. A direct bioautographic TLC assay for compounds possessing antibacterial activity. *J. Nat. Prod.*, 50: 19-22.

- Harborne, J.B., 1986. The Flavonoids. Chapman and Hall, London.
- Hostettman, K. *et al.*, 1998. Preparative Chromatography Techniques. Applications in Natural Product Isolation. 2nd Edn., Springer, Berlin Heidelberg.
- Mabry, T.J., K.R. Markham and H.B. Thomas, 1970. The Systematic Identification of Flavonoids. Springer Verlag, Heidelberg, New York.
- Maire, R., 1953. Flora of North Africa: Morocco, Algeria, Tunisia, Tripolitaine, Cyrenaïque and Sahara. Vol. 5, Paul Lechevallier: Paris.
- Markham, K.R. and H. Geiger, 1994. ¹H NMR Spectroscopy of Flavonoids. Advances in Research Since 1986. Harborne, J.B. (Ed.), Chapman and Hall, London, pp: 676.
- Ozenda, P., 1983. Flora of the Sahara. 2nd Edn., CNRS: Paris, pp: 270.
- Rukachaisirikul, V., Y. Sukpondma, C. Jansakul and W.C. Taylor, 2002. Isoflavone glycosides from *Derris scandens*. Phytochemistry, 60: 827-834.