



Research Journal of
Seed Science

ISSN 1819-3552



Academic
Journals Inc.

www.academicjournals.com

Effect of Exogenous Hormones and Chilling on Dormancy Breaking of Seeds of Asafoetida (*Ferula assafoetida* L.)

¹M. Otrshy, ²A. Zamani, ²M. Khodambashi, ¹M. Ebrahimi and ³P.C. Struik

¹Agricultural Biotechnology Research Institute of Iran,
P.O. Box 85135-487, Najafabad, Iran

²Shahr-e-kord University, Shahr-e-kord, Iran

³Centre for Crop Systems Analysis, Department of Plant Sciences,
Wageningen University, P.O. Box 430, 6700 AK Wageningen, The Netherlands

Abstract: Asafoetida (*Ferula assafoetida* L.) is a medicinal plant with a problematic seed germination. Seeds of this plant have a long dormancy. The present research was carried out to investigate whether exogenous application of the hormones Gibberellic Acid (GA₃), N6-furfurylaminopurine (kinetin) and 6-benzylaminopurine (BAP), alone or in combinations and chilling (for 0 or 28 days) could be used to alleviate the problem of dormancy. Germination was increased from 53.3% in a medium free of any hormones to 74.4% in a medium supplemented with 0.25 mg L⁻¹ BAP. This was the only single hormone treatment giving a statistically significant increase in germination. Chilling increased germination from 37.5% (0 days of chilling) to 82.9% (28 days of chilling). Combining exogenous application of 0.25 mg L⁻¹ BAP with 28 day chilling even gave more than 90% seed germination. Present findings suggest that seeds germination of Asafoetida can be greatly improved by combining chilling with BAP application.

Key words: Asafoetida, chilling, dormancy breaking, *Ferula assafoetida* L., exogenous application of hormones

INTRODUCTION

Asafoetida (*Ferula assafoetida* L.), also known as devil's dung, is an important specie of the Apiaceae family. It is a stout perennial, monocarpic herb (Singh Puri, 2003) reaching a height of 2-3 m with 2-4 pinnate pubescent leaves having cauline sheaths, yellow flowers in large terminal compound umbels and dorsally compressed cremocarps (Daniel, 2006). It is native to Iran.

This medicinal plant has important effects on the functioning of the digestive system and it is reported to be useful for breath, secretions, flatus and gastric eructation (Bown, 1995; Chevallier, 1996). The resin gum from the roots is vermifuge; it is antispasmodic, expectorant and appetizing (Chiej, 1984; Duke and Ayensu, 1985). The seeds of Asafoetida have a long dormancy. The dormant seeds do not germinate on the mother plant and they have enough time for dispersal (Baskin *et al.*, 1995). This characteristic is very useful. Seeds of most cultivated and garden plants lose their dormancy before or shortly after they are separated from the mother plant, but in contrast most wild plants have a long seed dormancy (Bryant, 1996). Usually a change in environmental conditions caused by seasonality of the

Corresponding Author: Mahmoud Otrshy, Agricultural Biotechnology Research Institute of Iran,
P.O. Box 85135-487, Isfahan, Najafabad, Iran

climate is a main factor determining the level of dormancy through their effects on the seed hormonal balance (Bewley and Black, 1994; Peng and Harberd, 2002; Thomas, 1990; Thomas and Sambrooks, 1985). Because seed dormancy is a nuisance in propagation programmes, removing or breaking dormancy has always been a challenge for plant physiologists (Baskin and Baskin, 1991).

Primary seed dormancy can be divided into two types: external and internal. Morpho-physiological dormancy belongs to the internal type of primary dormancy. It depends on plant species and seeds must be exposed to cold, heat, gibberellic acid or chemical materials for dormancy breaking (Bewley and Black, 1994). Seeds of many species of the Apiaceae family have different types of morpho-physiological dormancy that chilling can remove (Baskin and Baskin, 1991, 2004; Baskin *et al.*, 1999; Phillips *et al.*, 2003). In this family, different species of the genera *Erythronium*, *Osmorhiza* (Baskin and Baskin, 1984, 1989, 1991; Baskin *et al.*, 1995) and species like *Ferula gummosa*, *Teucrium polium* (Nadjafi *et al.*, 2006) and *Thaspium pinnatifidum* (Baskin *et al.*, 1992) need different periods of chilling to break seed dormancy.

GA₃ is one of the most important hormones that have been suggested to break primary dormancy (Nadjafi *et al.*, 2006). GA₃ can improve seed germination in Apiaceae which also requires light and cold (El-Dengawy, 2005; El-Nabawy *et al.*, 1980; Finch-Savage and Leubner-Metzger, 2006; Peng and Harberd, 2002; Thomas, 1990; Thomas and Sambrooks, 1985).

Asafoetida's seeds have a long dormancy and the aim of this research was to find efficient ways to break it. To the best of our knowledge no earlier studies on seed dormancy and dormancy breaking methods for *F. assafoetida* have been reported. For the first time we established *in vitro* culture systems for dormancy breaking of seeds of these plant species.

MATERIALS AND METHODS

The experiment was carried out at the tissue culture facility of the Agricultural Biotechnology Research Institute of Iran in October 2007.

Seeds of Asafoetida were obtained from the Medicinal Plants Collection of the Shahid Fozveh Research Center in July 2007. Seeds were washed under running tap water for 10 times to remove surface contamination. To eliminate probable pathogens, seeds were subsequently placed in a fungicide solution containing 2.5% (w/v) Benomyl and Carboxyn Tiram for 4 h. Seeds were rinsed again with sterile distilled water and then were soaked in ethanol 96% for 1 min. At the end of this sterilization process, seeds were placed in 25% chlorax supplemented with Tween 20 (1 drop) for 25 min. Seeds were rinsed again with sterile distilled water 3 times in a laminar flow cabinet. Fifteen seeds were put in a single Petri dish containing water agar and different concentrations of various hormones or hormone compositions.

In this study, 10 different hormone treatments and two chilling treatments (0 (control) and 28 days) were applied. Hormone treatments included (in addition to a control): 0.1 and 0.25 mg L⁻¹ BAP, 0.1 and 0.25 mg L⁻¹ kinetin, 5 mg L⁻¹ GA₃, plus the combinations of the BAP treatments or the kinetin treatments with the GA₃ treatment. Treatments are coded by the abbreviation of the substance followed by the concentration (without the unit).

For the chilling treatment Petri dishes containing seeds were placed in a dark refrigerator at 5°C temperature during 0 (control) or 28 days. After chilling these Petri dishes were transferred to a growth chamber at 24°C and with 18.5 μmol/m²/sec light intensity.

Germination rates were observed 2, 3 and 4 weeks after the transfer to the growth chamber. The experiment was conducted as a completely randomized factorial experiment with two factors and 3 replications. Data were statistically analyzed using the SAS package (Version 8 of the SAS System) (SAS, 1999) statistical computer program and Duncan's Multiple Range Tests were used to compare the Mean values.

RESULTS

The analysis of data showed that both experimental factors, chilling and hormone application, significantly affected seed germination. Chilling at 5°C for 28 days increased germination. For 0 day chilling treatment, only 37.5% of seeds were germinated, while chilling resulted in 82.9% germination.

The maximum positive effect of hormone application was an increase in germination by 21.1% (Table 1). Treatment 0.1 mg L⁻¹ kinetin showed a (non-significant) negative effect on germination (Table 1).

In Asafoetida, kinetin had no effect on germination while other cytokinin (BAP) showed a large positive effect on seed germination at the higher concentration. These results showed that different cytokinin may have different influences on dormancy break. Treatment 0.25 mg L⁻¹ BAP was more effective than the treatment with GA₃ and increased germination significantly. Table 2 shows the BAP effects averaged over the single treatment and the combination with GA₃.

It was observed that seed germination in the treatment without chilling was almost fixed from the third week onwards and hardly any further germination was observed. For the treatment without chilling hormone applications had an effect on the dynamics of germination. The highest value after two weeks was observed for the treatment 0.25 mg L⁻¹ BAP + 5 mg L⁻¹ GA₃ (Table 3). During the third week GA₃ had the highest additional germination. All hormone applications showed little germination in the fourth week (Table 3).

For the treatment with 28 days chilling, germination increased over time and considerable additional germination took place in the last week (Table 4). This late germination contributed much to the effect of chilling on germination.

Table 1: The effects of hormone application on germination

Treatments (mg L ⁻¹)	Germination (%)
BAP 0.1	61.1b
BAP 0.1+GA ₃ 5	60.0b
BAP 0.25	74.4a
BAP 0.25+GA ₃ 5	73.3a
Kin 0.1	50.0b
Kin 0.1+GA ₃ 5	53.3b
Kin 0.25	54.4b
Kin 0.25+GA ₃ 5	61.1b
WA (control)	53.3b

Mean values followed by the same letter(s) are not significantly different (p<0.05)

Table 2: Mean comparison of BAP concentrations

Concentration of BAP (mg L ⁻¹)	Mean of germinated seed (%)
0.00	59.0b
0.10	60.5b
0.25	73.9a

Mean values followed by the same letter(s) are not significantly different (p<0.05)

Table 3: The effects of hormones on percentage of additional germination at different intervals during germination without chilling

Treatments (mg L ⁻¹)	Germination (%)		
	2nd week	3rd week	4th week
BAP 0.1	40.0bc	2.2hi	6.7fghi
BAP 0.1 + GA ₃ 5	26.7d	4.4ghi	0.0i
BAP 0.25	42.2b	6.7fghi	8.9fghi
BAP 0.25 + GA ₃ 5	55.6a	0.0i	0.0i
Kin 0.1	13.3fgh	6.7fghi	0.0i
Kin 0.1 + GA ₃ 5	17.8def	6.7fghi	2.2hi
Kin 0.25	37.8bc	2.2hi	0.0i
Kin 0.25 + GA ₃ 5	26.7d	8.9fghi	0.0i
GA ₃ 5	22.2de	17.8defg	0.0i
WA (control)	15.6defg	6.7fghi	0.0i

Mean values followed by the same letter(s) are not significantly different (p<0.05)

Table 4: The effects of hormones on percentage of additional germination at different intervals during germination with chilling

Treatments (mg L ⁻¹)	Germination (%)		
	2nd week	3rd week	4th week
BAP 0.1	13.3ij	22.2h	37.8d
BAP 0.1+ GA ₃ 5	22.2h	44.4c	22.2h
BAP 0.25	13.3ij	62.2a	15.6i
BAP 0.25 + GA ₃ 5	15.6i	48.9b	26.7g
Kin 0.1	2.2kl	33.3e	44.4c
Kin 0.1 + GA ₃ 5	11.1j	37.8d	31.1ef
Kin 0.25	4.4kl	37.8d	26.7g
Kin 0.25 + GA ₃ 5	15.6i	42.2c	28.9fg
GA ₃ 5	11.1j	37.8d	33.3e
WA (control)	6.7kl	33.3e	44.4c

Mean values followed by the same letter(s) are not significantly different (p<0.05)

Germination showed different behaviours for treatments with chilling and without chilling. For 0 day chilling, most of the germination took place in the first two weeks, while after 28 days chilling most of the germination took place during the third week (Table 3, 4). Without chilling, all hormonal treatments, showed their maximum germination during the first two weeks and hardly any germination during the fourth week (Table 3). In contrast, after 28 days of chilling, all hormonal treatments showed most germination during the third or fourth week with relatively little germination during the first two weeks (Table 4).

The two-ways interaction of both factors (chilling and hormone application) was also significant regarding the final percentage of germinated seeds. Highest values for germination were obtained for 0.25 mg L⁻¹ BAP (91%) and 0.25 mg L⁻¹ BAP + 5 mg L⁻¹ GA₃ (93%) at 28 days of chilling while it was lowest with 0.1 mg L⁻¹ Kin without chilling (20%) (Table 5).

For chilling, highest and lowest percentage of germination was found with 0.25 mg L⁻¹ BAP + 0.5 mg L⁻¹ GA₃ and treatment 0.25 mg L⁻¹ kin, respectively. For non-chilling 0.25 mg L⁻¹ BAP + 5 mg L⁻¹ GA₃ resulted in the highest percentage. Of germination but 0.01 mg L⁻¹ kin showed lowest percentage of germination and there was no significant difference between this treatment and control.

It was interesting that on average 53.3% of final germination was obtained on a medium free of hormone (Table 1). When seeds were treated with chilling the control reached 84.4% but only 22.2% of germination was observed for the control without chilling (Table 5).

Table 5: Interaction between chilling and hormone treatment for germination

Treatments (mg L ⁻¹)	Germination (%)	
	0 day chilling	28 days chilling
BAP 0.1	48.9jkl	73.3bcdefgh
BAP 0.1 + GA ₃ 5	31.1mn	88.9abc
BAP 0.25	57.8hij	91.1ab
BAP 0.25 + GA ₃ 5	53.3ijk	93.3a
Kin 0.1	20.0mn	80.0bcdefg
Kin 0.1 + GA ₃ 5	26.7mn	80.0bcdefg
Kin 0.25	40.0klm	68.9defghi
Kin 0.25+ GA ₃ 5	35.6klmn	86.7abcd
GA ₃ 5	40.0klm	82.2abcdef
WA (control)	22.2mn	84.4abcde

Mean values followed by the same letter(s) are not significantly different (p<0.05)

DISCUSSION

The percentage of germination in *Asafoetida* can be increased with a chilling treatment. Walck and Hidayati (2004) reported that seed dormancy of *Osmorhiza depauperata* was broken by a 32 week incubation at 1-5°C and GA₃ was not a profitable substitute for cold. In *Thaspium pinnatifidum*, chilling was needed for breaking the physiological seed dormancy. It can stimulate immature embryos to grow and break their morphological dormancy (Baskin *et al.*, 1992). The natural habitats of *F. gummosa*, higher seed germination percentage occurred in colder regions with higher precipitation. In *Chaerophyllum temulum* growth of embryo also happened on the long term chilling at 5°C (Vandelook *et al.*, 2007). The main reason to explain the effect of chilling on breaking of seed dormancy is enhancing internal GA₃ accumulation (Phillips *et al.*, 2003). GA₃ is suggested to break primary dormancy and stimulate germination (Eastmond and Jones, 2005).

Dormancy is one of the most important physiological properties that are controlled by ABA/GA₃ ratio. At high levels of ABA and low levels of GA₃, the embryo remains dormant and with a reversed ABA:GA₃ ratio, the intensity of the embryo's dormancy is reduced (Finch-Savage and Leubner-Metzger, 2006). If the chilling enhances the concentration of internal GA₃ in seed, so exogenous application of GA₃ should also cause more or faster germination, but in the case of *Asafoetida*, contrary to most species of the Apiaceae, application of this hormone had no effect on germination of seeds. Therefore, cold treatment may have other effects on dormancy breaking that still require elucidation. Tsukamoto (1972) applied cytokinin to break dormancy of *Gladiolus*. Tsukamoto (1972) reported that cytokinin can break dormancy of many species and also dormancy of buds in trees but he could not find any reason for that. Cytokinin are not necessary for germination, but they can affect ABA which inhibits germination and overcome thermo-dormancy (Litwack, 2005). It has been reported that cytokinin enhance the effect of GA₃ in the seed system of Celery and cause faster germination (Biddington and Thomas, 1978; Thomas and Van Staden, 1995). In *Asafoetida*, kinetin had no significant effect on germination but presence of BAP resulted in significant effects on germination, so it can be suggested that cytokinin may affect dormancy breaking in different ways or that different cytokinin have different activities.

Present result shows that the seeds of *Asafoetida* have a deep and complex morpho-physiological dormancy and like many species of Apiaceae, they can germinate without GA₃ but are enhanced by chilling (at 5°C temperatures). Application of BAP is also a simulative factor for dormancy breaking.

CONCLUSION

From this study, it can be suggested that medium supplemented with BAP (0.25 mg L⁻¹) and without any nutrient can enhance dormancy breaking in Asafoetida seeds. GA₃ does not have sufficient effect on Asafoetida seed germination. Chilling can remove dormancy and enhances germination. With 28 days chilling and use of a medium containing 0.25 mg L⁻¹ BAP we could obtain more than 90% germination in seeds of Asafoetida. Longer chilling and higher concentrations of BAP may increase germination even further.

REFERENCES

- Baskin, C.C. and J.M. Baskin, 1984. Germination ecophysiology of the woodland herb *Osmorhiza longistylis* (Umbeliferae). Am. J. Bot., 71: 687-692.
- Baskin, C.C. and J.M. Baskin, 1989. Seed germination ecophysiology of *Jeffersonia diphylla*, a perennial herb of mesic deciduous forests. Am. J. Bot., 76: 1073-1080.
- Baskin, C.C. and J.M. Baskin, 1991. Nondeep complex morphophysiological dormancy in seeds of *Osmorhiza claytonii* (Apiaceae). Am. J. Bot., 78: 588-593.
- Baskin, C.C., E.W. Chester and J.M. Baskin, 1992. Deep complex morphophysiological dormancy in seeds of *Thaspium pinnatifidum* (Apiaceae). Int. J. Plant Sci., 153: 565-571.
- Baskin, C.C., E. Meyer and J.M. Baskin, 1995. Two type morphophysiological dormancy in seeds of two genera *Osmorhiza* and *Erythronium* with an Arcto-Tertiary distribution pattern. Am. J. Bot., 82: 293-298.
- Baskin, C.C., J.M. Baskin and E.W. Chester, 1999. Seed dormancy in the wetland winter annual *Pitilimnium nuttallii* (Apiaceae). Wetlands, 19: 359-364.
- Baskin, J.M. and C.C. Baskin, 2004. A classification system for seed dormancy. Seed Sci. Res., 14: 1-16.
- Bewley, J.D. and M. Black, 1994. Seeds: Physiology of Development and Germination. 2nd Edn., Plenum Press, London, ISBN: 0-306-44747-9, pp: 445.
- Biddington, N.L. and T.H. Thomas, 1978. Thermodormancy in celery seed and its removal by cytokinins and gibberellins. Physiol. Plantarum, 42: 401-405.
- Bown, D., 1995. Encyclopaedia of Herbs and their Uses. Dorling Kindersley, Ltd., London, ISBN: 0-7513-020-31, pp: 342.
- Bryant, G.A., 1996. Seed Physiology. Edward Arnold, London.
- Chevallier, A., 1996. The Encyclopedia of Medicinal Plants. 1st Edn., DK Publishing Inc., New York, USA., pp: 259.
- Chiej, R., 1984. Encyclopaedia of Medicinal Plants. Macdonald and Co., London, UK., pp: 1-139.
- Daniel, M., 2006. Medicinal Plants: Chemistry and Properties. Science Publishers, USA.
- Duke, J.A. and E.S. Ayensu, 1985. Medicinal Plants of China. 1st Edn., Reference Publisher, Algonac, MI.
- Eastmond, P.J. and R.L. Jones, 2005. Hormonal regulation of gluconeogenesis in cereal aleurone is strongly cultivar-dependent and gibberellin action involves SLENDER1 but not GAMYB. Plant J., 44: 483-493.
- El-Dengawy, E.F.A., 2005. Promotion of seed germination and subsequent seedling growth of loquat (*Eriobotrya japonica* Lindl.) by moist-chilling and GA₃ applications. Sci. Hort., 105: 331-342.
- El-Nabawy, S., M. Abou Rawash, A. El-Hamady, I. Desouky and F. Khalil, 1980. Effect of stratification and GA₃ on germination of pecan seeds and subsequent seedling growth. Ann. Agric. Sci., 25: 323-338.

- Finch-Savage, W.E. and G. Leubner-Metzger, 2006. Seed dormancy and the control of germination. *New Phytol.*, 171: 501-523.
- Litwack, G., 2005. *Plant Hormones* (ED). Academic Press, California, USA.
- Nadjafi, F., M. Bannayan, M. Rastgoo and L. Tabrizi, 2006. Seed germination and dormancy breaking techniques for *Ferula gummosa* and *Teucrium polium*. *J. Arid Environ.*, 64: 542-547.
- Peng, J. and N.P. Harberd, 2002. The role of GA-mediated signaling in the control of seed germination. *Curr. Opin. Plant Biol.*, 5: 376-381.
- Phillips, N., D. Drost and W. Varga, 2003. Chemical treatments enhanced seed germination in *Perideridia gairdneri*. *Acta Hort.*, 618: 477-482.
- SAS., 1999. Version 8 (TS MO) of the SAS System. SAS Institute Inc., Cary, NC, USA.
- Singh Puri, H., 2003. *Rasayana: Ayurvedic Herbs for Longevity and Rejuvenation*. CRC Press, USA.
- Thomas, T. and R. Sambrooks, 1985. Possible control of gibberellin-induced release of temperature dependent primary dormancy in seeds of celery (*Apium graveolens*) by transmembrane ion fluxes. *Plant Growth Regul.*, 3: 191-199.
- Thomas, T.H., 1990. Hormonal involvement in photoregulation of celery seed dormancy. *Br. Soc. Plant Growth Regul.*, 20: 51-58.
- Thomas, T.H. and J. Van Staden, 1995. Dormancy break of celery (*Apium graveolens* L.) seeds by plant derived smoke extract. *Plant Growth Regul.*, 17: 195-198.
- Tsukamoto, Y., 1972. Breaking dormancy in the gladiolus corm with cytokinins. *Proc. Japan Acad.*, 48: 34-39.
- Vandelook, F., N. Bolle and J.A. Van Assche, 2007. Seed dormancy and germination of the European *Chaerophyllum temulum* (Apiaceae) a member of a Trans-Atlantic Genus. *Ann. Bot.*, 100: 233-239.
- Walck, J.L. and S.N. Hidayati, 2004. Germination ecophysiology of the western North American species *Osmorhiza depauperata* (Apiaceae): Implications of preadaptation and phylogenetic niche conservatism in seed dormancy evolution. *Seed Sci. Res.*, 14: 387-394.