



Research Journal of  
**Seed Science**

ISSN 1819-3552



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)

## Effects of Hydropriming Duration and Limited Irrigation on Field Performance of Chickpea\*

K. Ghassemi-Golezani, P. Sheikhzadeh-Mosaddegh and M. Valizadeh  
Faculty of Agriculture, University of Tabriz, Tabriz, Iran

**Abstract:** Chickpea seeds (cv. ILC<sub>482</sub>) were soaked in distilled water at 17.3°C for 8(P<sub>2</sub>), 16(P<sub>3</sub>) and 24(P<sub>4</sub>) hours and subsequently dried back to about 30% moisture content. Germination percentage, germination rate and seedling dry weight for different treatments were determined in laboratory. The field experiment was conducted under different irrigation regimes (I<sub>1</sub>, I<sub>2</sub> and I<sub>3</sub>-irrigation after 80, 120 and 160 mm evaporation from class A pan) in 2006. In laboratory, seed germination rate was increased with increasing hydropriming duration. Maximum germination rate was obtained for P<sub>4</sub> treatment, which was not significantly different from P<sub>3</sub> treatment. Seedling dry weight of hydroprimed seeds was significantly higher than that of control (P<sub>1</sub>). In the field, ground cover, biological yield and grains per unit area were decreased, as water deficit severed. The lowest number of grains per unit area, 1000 grain weight and grain yield per unit area were obtained under I<sub>3</sub>. However, these traits were not significantly different between I<sub>1</sub> and I<sub>2</sub>. Mean seedling emergence rate and percentage improved, as a result of hydropriming. Consequently, ground cover of plants from primed seeds was higher than that from unprimed seeds, but this difference was not statistically significant. Mean biological yield, grains per unit area and grain yield of plants from primed seeds were significantly higher than those from non-primed seeds, but the effect of hydropriming duration on these traits was not significant. Mean 1000 grain weight of plants from P<sub>1</sub> and P<sub>4</sub> treatments decreased, with increasing water limitation. However, plants from P<sub>2</sub> and P<sub>3</sub> treatments produced the largest grains under I<sub>2</sub>. Yield improvement was 25-40%, depending on hydropriming duration. The highest yield increase was achieved with 16 h hydropriming. It was, therefore, suggested that this priming duration is the best treatment for invigoration of chickpea seeds.

**Key words:** Chickpea, hydropriming, seedling emergence, ground cover, grain yield

### INTRODUCTION

Chickpea (*Cicer arietinum* L.) is cultivated across a wide range of environments, from the subtropics to arid and semiarid Mediterranean regions and has considerable importance as food, feed and fodder (Singh, 1997; Saxena *et al.*, 1996). Although, chickpea is one of the major drought tolerant food legumes, but water deficit can limit growth and productivity of this crop (Ludlow and Muchow, 1990; Singh and Rao, 1993). Different techniques could be used to enhance chickpea yield, particularly under drought conditions. One of the simple techniques which can improve seedling vigor and establishment and consequently crop performance in the field is seed priming (Eira and Marcos Filho, 1990; Khan, 1992; McDonald, 2000).

Priming allows seed hydration to initiate the early events of germination but not to permit radicle emergence, followed by drying (McDonald, 2000; Ashraf and Foolad, 2005). Methods of seed priming include soaking seeds in water or osmotic solution and intermixture with matrix materials

---

**Corresponding Author:** K. Ghassemi-Golezani, Faculty of Agriculture, University of Tabriz, Tabriz, Iran  
Tel: 098411-3392028 Fax: 098411-3356003

\*Originally Published in *Research Journal of Seed Science*, 2008

(Khan, 1992; McDonald, 2000). There are reports that seed priming permits early DNA replication, increase RNA and Protein synthesis, enhances embryo growth, repairs deteriorated seed parts and reduces leakage of metabolites (McDouald, 2000). The early improvements may increase the rate and uniformity of seed germination and seedling emergence (Rowse *et al.*, 2001; Basra *et al.*, 2003, 2005; Farooq *et al.*, 2005; Abdurrahmani *et al.*, 2007). In this way, some of the deleterious effects of water limitation on crop performance may also be overcome by seed priming (Singh and Rao, 1993; Demir and Van de Venter, 1999; Bittencourt *et al.*, 2004; Demir *et al.*, 2006).

Hydropriming is a very simple, economical and environmental friendly type of seed priming in which seeds are soaked in water for a certain time and dried before sowing (Thornton and Powell, 1992). Since the beneficial effects of hydropriming on field performance of chickpea under water stress are poorly documented, this research was aimed to investigate the effects of hydropriming duration on seed invigoration, seedling emergence and grain yield of chickpea under different irrigation regimes.

## **MATERIALS AND METHODS**

Seeds of chickpea (cv. ILC<sub>482</sub>) were obtained from Dryland Agricultural Research Institute, Maragheh, Iran. These seeds were divided into four sub-samples, one of which was kept as control (non-primed, P<sub>0</sub>) and three other samples were soaked in distilled water at 17.3°C for 8 (P<sub>1</sub>), 16 (P<sub>2</sub>) and 24 (P<sub>3</sub>) hours and then dried back to about 30% moisture content at room temperature of 20-22°C. All the seeds were treated with Mancozebe at a rate of 2 g kg<sup>-1</sup>. Laboratory tests were carried out with CR design at Seed Technology Laboratory of Tabriz University, Tabriz, Iran. Four replicates of 25 seeds were placed between moist filter papers and germinated in an incubator adjusted on 10°C for 6 days. Germination (protrusion of radicle by 2 mm) was recorded in daily intervals. Seed germination rate for each treatment was calculated according to Ellis and Roberts (1980). At the end, percentage of normal seedlings and seedling dry weight were also determined.

The field experiment was conducted at the Research Farm of Tabriz University (Latitude 38° 05' N, Longitude 46° 17' E, Altitude 1360 m above sea level) in 2006. The climate is characterized by mean annual precipitation of 246 mm, mean annual temperature of 10°C, annual maximum temperature of 16.6°C and mean annual minimum temperature of 4.2°C. Soil type was sandy loam with EC of 0.68 dS m<sup>-1</sup> and pH of 8.1.

Seeds were hand sown in about 4 cm depth with a density of 68 seeds m<sup>-2</sup> on 11 April 2006. Each plot consisted of 6 rows with 5 cm length, spaced 25 cm apart. Experimental design was factorial, based on RCB design with four replications. All plots were irrigated immediately after sowing and then fertilized with 30 kg ha<sup>-1</sup> urea. Subsequent irrigations were carried out after 80 (I<sub>1</sub>), 120 (I<sub>2</sub>) and 160 (I<sub>3</sub>) mm evaporation from class A pan. Weeds were controlled by hand weeding during crop growth and development. Plants were protected from heliothis caterpillar attack by spraying Diazinon (2 ml L<sup>-1</sup>) at pod formation stage.

Seedling emergence was recorded in daily intervals up to the day, when maximum seedling establishment in each plot was achieved. Subsequently, rate and percentage of seedling emergence were calculated. Ground cover was measured every week, by viewing the canopy through wooden frame (50×50 cm dimensions) divided into 100 equal sections. The sections were counted when more than half filled with crop green area. At maturity, 10 plants were harvested from each plot and grains per unit area, 1000 grain weight and above ground biomass were determined. Finally, plants of 3 m<sup>2</sup> in the middle part of each plot were harvested and grain yield per unit area was recorded.

Analysis of variance of the data appropriate to the experimental design and comparison of means at p≤0.05 were carried out, using MSTATC software. EXCEL software was used to draw figures.

## RESULTS

Analysis of variance of the laboratory data showed significant effects of hydropriming duration on germination rate and seedling dry weight ( $p \leq 0.05$ ). The highest germination rate was achieved with  $P_4$  (hydropriming for 24 h), which was not significantly different from  $P_3$  (hydropriming for 16 h). Rate of germination for  $P_2$  (hydropriming for 8 h) was significantly lower than that for  $P_3$  and  $P_4$ , but it was significantly higher than that for  $P_1$  (non-primed seeds). Primed seeds ( $P_2$ ,  $P_3$  and  $P_4$ ) were also produced larger seedlings, compared to non-primed seeds ( $P_1$ ). However, no significant effect of hydropriming duration on seedling dry weight was observed (Table 1).

Seedling emergence rate and percentage in the field were significantly affected by hydropriming duration, but the effects of irrigation regimes and irrigation  $\times$  hydropriming interaction on these traits were not significant (Table 2). Rate and percentage of seedling emergence for primed seeds were significantly higher than those for non-primed seeds. However, seedling emergence did not differ significantly among  $P_2$ ,  $P_3$  and  $P_4$  (Table 4).

Effect of irrigation on ground cover in the field was significant (Table 2). Mean ground cover percentage decreased, as water limitation increased. However, differences between  $I_1$  and  $I_2$  and between  $I_2$  and  $I_3$  were not significant (Table 3). Although mean ground cover of plants from primed seeds was higher than that from unprimed seeds, but this difference was not statistically significant (Table 2, 4). Ground green cover of chickpea cultivars under different irrigation regimes increased with increasing growth period up to about 66-68 days after sowing and then started to decrease (Fig. 1). In general, ground cover of plants from  $P_2$  and  $P_3$  treatments was higher than that from  $P_1$  and  $P_4$  treatments (Fig. 1, Table 4).

Table 1: Effects of hydropriming on mean germination rate, germination percentage and seedling dry weight in laboratory

Treatments	Germination rate (per day)	Germination (%)	Seedling dry weight (mg)
$P_1$	0.340 <sup>c</sup>	96 <sup>c</sup>	21.2 <sup>b</sup>
$P_2$	0.453 <sup>b</sup>	97 <sup>a</sup>	27.0 <sup>a</sup>
$P_3$	0.502 <sup>a</sup>	94 <sup>a</sup>	27.5 <sup>a</sup>
$P_4$	0.515 <sup>a</sup>	95 <sup>a</sup>	28.2 <sup>a</sup>

Different letters indicating significant difference at  $p \leq 0.05$ .  $P_1$ ,  $P_2$ ,  $P_3$  and  $P_4$ : non-primed and hydroprimed seeds for 8, 16 and 24 h, respectively

Table 2: Analysis of variance of the effects of hydropriming on field performance of chickpea

Source	df	MS							
		Emergence rate	Emergence percentage	Ground cover	Grains per unit area	1000 grain weight	Biological yield	Harvest index	Grain yield
Replication	3	2.461 $\times 10^{-1}$ ns	304.267*	26.919ns	30608.892ns	220.971ns	4455.110ns	108.273ns	1508.422ns
Irrigation	2	1.631 $\times 10^{-3}$ ns	8.644ns	73.847*	125484.357*	2665.514*	13001.372*	84.906ns	2139.225*
Hydropriming	3	1.573 $\times 10^{-3}$ **	331.230*	40.914ns	79243.317*	696.528ns	12660.088**	226.236ns	3653.365*
Irrigation $\times$ Hydropriming	6	2.164 $\times 10^{-3}$ ns	59.376ns	16.748ns	4634.603ns	1967.886*	1363.194ns	75.605ns	832.126ns
Error	33	3.03 $\times 10^{-1}$	87.705	21.041	26146.316	785.250	2801.560	75.588	730.680

ns, \*, \*\* : No significant and significant at  $p \leq 0.05$  and  $p \leq 0.01$ , respectively

Table 3: Comparison of mean ground cover, grain yield and yield components of chickpea under different irrigation regimes

Treatments	Ground cover (%)	Biological yield ( $g m^{-2}$ )	Grains $m^{-2}$	1000 Grain weight (g)	Grain yield ( $g m^{-2}$ )
$I_1$	37.854 <sup>a</sup>	242.0 <sup>a</sup>	529.1 <sup>a</sup>	307.244 <sup>ab</sup>	78.85 <sup>a</sup>
$I_2$	35.970 <sup>ab</sup>	206.7 <sup>ab</sup>	434.1 <sup>ab</sup>	324.837 <sup>a</sup>	101.20 <sup>a</sup>
$I_3$	33.567 <sup>b</sup>	185.6 <sup>b</sup>	352.2 <sup>b</sup>	299.681 <sup>b</sup>	80.60 <sup>b</sup>

Different letters indicating significant difference at  $p \leq 0.05$ .  $I_1$ ,  $I_2$  and  $I_3$ : Irrigation after 80, 120 and 160 mm evaporation from class A pan, respectively

Table 4: Means of field traits for chickpea affected by hydropriming

Treatments	Emergence rate (per day)	Emergence (%)	Ground cover (%)	Biological yield ( $g m^{-2}$ )	Grains $m^{-2}$	Grain yield ( $g m^{-2}$ )	Yield improvement (%)
$P_1$	0.061 <sup>c</sup>	78.92 <sup>b</sup>	33.685a	168.9 <sup>b</sup>	324.5 <sup>b</sup>	78.805 <sup>b</sup>	0
$P_2$	0.068 <sup>b</sup>	89.46 <sup>a</sup>	36.942 <sup>a</sup>	223.8 <sup>a</sup>	460.3 <sup>ab</sup>	101.00 <sup>ab</sup>	28
$P_3$	0.068 <sup>b</sup>	89.95 <sup>a</sup>	37.692 <sup>a</sup>	237.3 <sup>a</sup>	517.0 <sup>a</sup>	110.70 <sup>a</sup>	40
$P_4$	0.068 <sup>b</sup>	88.73 <sup>a</sup>	34.868 <sup>a</sup>	220.7 <sup>a</sup>	452.1 <sup>ab</sup>	98.67 <sup>ab</sup>	25

Different letters indicating significant difference at  $p \leq 0.05$ .  $P_1$ ,  $P_2$ ,  $P_3$  and  $P_4$ : non-primed and hydroprimed seeds for 8, 16 and 24 h, respectively




Fig. 1: Changes in ground cover of chickpea affected by seed priming and water limitation during growth and development

P<sub>1</sub>, P<sub>2</sub>, P<sub>3</sub> and P<sub>4</sub>: non-primed and hydroprimed seeds for 8, 16 and 24 h, respectively

I<sub>1</sub>, I<sub>2</sub> and I<sub>3</sub>: Irrigation after 80, 120 and 160 mm evaporation from class A pan, respectively

Table 5: Mean 1000 grain weight of chickpea for hydropriming × irrigation interaction

Treatments	P <sub>1</sub>	P <sub>2</sub>	P <sub>3</sub>	P <sub>4</sub>
I <sub>1</sub>	323.15 <sup>ab</sup>	289.13 <sup>bc</sup>	300.50 <sup>bc</sup>	316.20 <sup>bc</sup>
I <sub>2</sub>	316.98 <sup>abc</sup>	332.05 <sup>ab</sup>	352.78 <sup>a</sup>	297.55 <sup>bc</sup>
I <sub>3</sub>	309.83 <sup>abc</sup>	321.95 <sup>abc</sup>	282.03 <sup>c</sup>	284.93 <sup>bc</sup>

Different letters indicating significant difference at p ≤ 0.05. P<sub>1</sub>, P<sub>2</sub>, P<sub>3</sub> and P<sub>4</sub>: non-primed and hydroprimed seeds for 8, 16 and 24 h, respectively. I<sub>1</sub>, I<sub>2</sub> and I<sub>3</sub>: Irrigation after 80, 120 and 160 mm evaporation from class A pan, respectively

Thousand grain weight was only affected by irrigation, but grains per unit area and grain yield were affected by both irrigation and hydropriming. Irrigation × hydropriming interaction for 1000 grain weight was also significant. Neither priming nor irrigation had significant effect on harvest index (Table 2). Mean number of grains per unit area, 1000 grain weight and grain yield were significantly decreased under severe water deficit (I<sub>3</sub>), compared to other irrigation treatments (Table 3). Grains per unit area and grain yield for P<sub>1</sub> were lower than those for P<sub>2</sub>, P<sub>3</sub> and P<sub>4</sub>, but these traits were statistically similar for primed seeds (Table 4). Mean 1000 grain weight of plants from P<sub>1</sub> and P<sub>4</sub> treatments decreased, with increasing water limitation. However, plants from P<sub>2</sub> and P<sub>3</sub> treatments produced the largest grains under I<sub>2</sub> (Table 5).

## DISCUSSION

Hydropriming improved seedling vigor of chickpea as indicated by seed germination rate and seedling dry weight in laboratory tests (Table 1). The earlier germination might be attributed

to increased metabolic activities in the hydroprimed seeds (Soon *et al.*, 2000; Basra *et al.*, 2002). It is obvious that higher rate of seed germination can lead to the production of large and uniform seedlings. According to McDonald (2000), primed seeds can rapidly imbibe and revive the seed metabolism, enhancing germination rate and uniformity. In many crops, seed germination and early seedling growth are the most sensitive stages to water limitation. Water deficit may delay the onset and reduce the rate and uniformity of germination, leading to poor crop performance and yield (Demir *et al.*, 2006). Therefore, the beneficial effects of priming may be more evident under unfavorable rather than favorable conditions (Parera and Cantliffe, 1994; Bradford, 1995).

Since plots irrigated immediately after sowing, seeds did not experience drought stress during germination and seedling emergence in the field. So, rate and percentage of seedling emergence for irrigation treatments were not statistically different (Table 2). Therefore, reductions in ground cover and biological yield of chickpea under limited irrigation (Table 3) were not influenced by rate and percentage of seedling emergence. These reductions were then the consequence of growth retardation, due to water deficit (Sing, 1993; Kamara *et al.*, 2003). Poor ground cover and biological yield under severe water deficit ( $I_3$ ) resulted in production of less and smaller grains and finally lower grain yield per unit area (Table 3), compared to well-watering ( $I_1$ ) and mild water stress ( $I_2$ ). These traits did not differ significantly between  $I_1$  and  $I_2$ , suggesting that when water resources are limited, irrigation after 120 mm evaporation from class A pan could be a favorite irrigation regime for chickpea crop.

Enhanced emergence rate and percentage of chickpea seedlings as a result of hydropriming, without any effect of priming duration (Table 4), suggest that hydration of chickpea seeds is an effective method to improve seedling vigor and establishment in the field. The rate and percentage of seedling emergence are very important in determining yield potential of annual crops (Kahlon *et al.*, 1992; Musa *et al.*, 1999; Hussain *et al.*, 2006). Rapid emergence of seedlings could lead to the production of vigorous plants. In addition, improvement of seedling emergence percentage could help to establish optimum plant population density under a wide range of environmental conditions (Harris *et al.*, 1999; Podlaski *et al.*, 2003; Hussain *et al.*, 2006).

Some improvement in ground cover of plants from primed seeds (Fig. 1) was clearly related with the early emergence and superior establishment of their seedlings (Table 4). Since, there is a linear relationship between ground cover and light interception (Burstall and Harris, 1983), it is, therefore a reliable index to estimate crop performance and yield, particularly under adverse conditions. Also, it has the practical advantage of quick, simple and non-destructive measurement, allowing frequent sampling.

The beneficial effects of hydropriming on grain yield were previously reported in sunflower (Hussain *et al.*, 2006), rice (Farooq *et al.*, 2006) and wheat (Kahlon *et al.*, 1992). In this research, superiorities of hydroprimed seeds in stimulation of seedling emergence and establishment continued to improve biological yield, grains  $m^{-2}$  and grain yield per unit area. Plants from  $P_2$  and  $P_3$  also produced the largest grains under  $I_2$  (Table 5).

## CONCLUSION

Hydropriming improved field performance and grain yield of chickpea. Yield improvement was 25-40%, depending on hydropriming duration. The highest yield increase was obtained with 16 h hydropriming (Table 4). Thus, it is conceivable to suggest that this priming duration is the best treatment for invigoration of chickpea seeds.

## REFERENCES

- Abdurrahmani, B., K. Ghassemi-Golezani, M. Valizadeh and V. Feizi asl, 2007. Seed priming and seedling establishment of barley (*Hordeum vulgare* L.). J. Food Agric. Environ., 5: 179-184.

- Ashraf, M. and M.R. Foolad, 2005. Pre-sowing seed treatment-a shotgun approach to improve germination, plant growth and crop yield under saline and non-saline conditions. *Adv. Agron.*, 88: 223-271.
- Basra, S.M.A., M.N. Zia, T. Mehmood, I. Afzal and A. Khaliq, 2002. Comparison of different invigoration techniques in wheat (*Triticum aestivum* L.) seeds. *Pak. J. Arid Agric.*, 5: 11-16.
- Basra, S.M.A., M. Farooq and A. Khaliq, 2003. Comparative study of pre-sowing seed enhancement treatment in indica rice (*Oryza sativa* L.). *Pak. J. life Soc. Sci.*, 1: 5-9.
- Basra, S.M.A., I. Afzal, A.R. Rashid and M. Farooq, 2005. Pre-sowing seed treatment to improve germination and seedling growth in wheat (*Triticum aestivum* L.). *Cadernode Pesquisa Sér. Bio., Santa Cruz de Sul.*, 17: 155-164.
- Bittencourt, M.L.C., D.C.F.S. Dias, L.A.S. Dias and E.F. Araújo, 2004. Effects of priming on asparagus seed germination and vigour under water and temperature stress. *Seed Sci. Technol.*, 32: 607-616.
- Bradford, K.J., 1995. Water Relations in Seed Germination. In: *Seed Development and Germination*, Kigel, J. and G. Galili (Eds.). Marcel Dekker, New York, pp: 351-396.
- Burstall, L. and P.M. Harris, 1983. The estimation of percentage light interception from leaf area index and percentage ground cover in potatoes. *J. Agric. Sci.*, 100: 241-244.
- Demir, I. and H.A. Van de Venter, 1999. The effect of priming treatments on the performance of watermelon (*Citrullus lanatus* (Thunb.) Matsum. and Nakai) seeds under temperature and osmotic stress. *Seed Sci. Technol.*, 27: 871-875.
- Demir, K., G. Okçu, M. Atak, Y. Çıkılı and O. Kolsarici, 2006. Seed treatments to overcome salt and drought stress during germination in sunflower (*Helianthus annuus* L.). *Eur. J. Agron.*, 24: 291-295.
- Eira, M.T.S. and J. Marcos-Filho, 1990. Osmotic conditioning of lettuce seeds: II. Performance under moisture, saline and temperature stresses. *Revista Brasileira de Sementes*, 12: 28-45.
- Ellis, R.H. and E.H. Roberts, 1980. Towards a Rational Basis for Testing Seed Quality. In: *Seed Production*, Hebblethwaite, P. D. (Ed.). Butterworths, London, pp: 605-635.
- Farooq, M., S.M.A. Basra, K. Hafeez and N. Ahmad, 2005. Thermal hardening: A new seed vigour enhancement tool in rice. *J. Integrative Plant Biol.*, 47: 187-193.
- Farooq, M., S.M.A. Basra and Hafeez-ur-Rehman, 2006. Seed priming enhances emergence, yield and Quality of direct-seeded rice. *Crop Manage. Physiol.*, 3: 42-44.
- Harris, D., A. Joshi, P.A. Khan, P. Gothkar and P.S. Sodhi, 1999. On-farm seed priming in semi-arid agriculture: Development and evaluation in maize, rice and chickpea in India, using participatory methods. *Exp. Agric.*, 35: 15-29.
- Hussain, M., M. Farooq, S. M.A. Basra and N. Ahmad, 2006. Influence of seed priming techniques on the seedling establishment, yield and quality of hybrid sunflower. *Int. Agric. Biol.*, 8:14-18.
- Kahlon, P.S., H.S. Dhaliwal, S.K. Sharma and A.S. Randawa, 1992. Effect of pre-sowing seed soaking on yield of wheat (*Triticum aestivum*) under late sown irrigated conditions. *Indian J. Agric. Sci.*, 62: 276-277.
- Kamara, A.Y., A. Menkir, B. Badu-Apraku and O. Ibikunle, 2003. The influence of drought stress on growth, yield and yield components of selected maize genotypes. *J. Agric. Sci.*, 141: 43-50.
- Khan, A.A., 1992. Preplant physiological seed conditioning. *Hortic. Rev.*, 13: 131-181.
- Ludlow, M.M. and R.C. Muchow, 1990. A critical evaluation of traits for improving crop yields in winter limited environments. *Adv. Agron.*, 43: 107-153.
- McDonald, M.B., 2000. Seed Priming. In: *Seed Technology and Its Biological Basis*, Black, M. and J.D. Bewley (Eds.). Sheffield Academic Press. England. Chapter 9, pp: 287-325.
- Musa, A.M., C. Johansen, J. Knmar and D. Harris, 1999. Response of chickpea to seed priming in the high Barind Tract of Bangladesh. *Int. Chickpea Pigeonpea Newslett.*, 6: 20-22.

- Parera, C.A. and D.J. Cantliffe, 1994. Presowing seed priming. *Hortic. Rev.*, 16: 109-141.
- Podlaski, S., Z. Chrobak, Z. Wyszowska, 2003. Effect of parsley seed treatment on root yield. *Plant Soil Environ.*, 5: 213-217.
- Rowse, H.R., J.M.T. Mckee and W.E. Finch-Savage, 2001. Membranes priming-A method for small samples of high value seeds. *Seed Sci. Technol.*, 29: 587-597.
- Saxena, N.P., M.C. Saxena, C. Johansen, S.M. Virmani and H. Harris, 1996. Future Research Priorities for Chickpea in Wana and Sat. In: *Adaptation of Chickpea in the West Asia and North Africa Region*, Saxena, N.P., M.C. Saxena, C. Johansen, S.M. Virmani and H. Harris (Eds.). ICARDA, Aleppo, Syria, pp: 257-263.
- Sing, K.B., 1993. Problems and Prospects of Stress Resistance Breeding in Chickpea. In: *Breeding for Stress Tolerance in Cool-Season Food Legumes*, Sing, K.B. and M.C. Saxena, (Eds.). Willey, Chichester, pp: 17-35.
- Singh, B.G. and G. Rao, 1993. Effect of Chemical Soaking of Sunflower (*Helianthus annuus* L.) Seed on Vigour Index. *Indian J. Agric. Sci.*, 63: 232-233.
- Singh, K.B., 1997. Chickpea (*Cicer arietinum* L.). *Field Crops Res.*, 53: 161-170.
- Soon, K.J., C.Y. Whan, S.B. Gu, A.C. Kil and C.J. Lia, 2000. Effect of hydropriming to enhance the germination of gourd seeds. *J. Korean Soc. Hortic. Sci.*, 41: 559-564.
- Thornton, J.M. and A.A. Powell, 1992. Short-term aerated hydration for the improvement of seed quality in *Brassica oleracea*. *Seed Sci. Res.*, 2: 41-49.