



Trends in  
**Applied Sciences  
Research**

ISSN 1819-3579



Academic  
Journals Inc.

[www.academicjournals.com](http://www.academicjournals.com)

## Efficiency of Universal Barcode Gene (Coxi) on Morphologically Cryptic Mugilidae Fishes Delineation

<sup>1</sup>C. Prasanna Kumar, <sup>2</sup>B. Akbar John, <sup>1</sup>S. Ajmal Khan, <sup>1</sup>P.S. Lyla, <sup>1</sup>S. Murugan, <sup>3</sup>M. Rozihan and <sup>2</sup>K.C.A. Jalal

<sup>1</sup>Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, Tamil Nadu, India

<sup>2</sup>Institute of Oceanography and Maritime Studies, International Islamic University Malaysia, Bandar Indera Mahkota, 25200 Kuantan, Pahang, Malaysia

<sup>3</sup>Department of Aquaculture, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

*Corresponding Author: P.S. Lyla, Centre of Advanced Study in Marine Biology, Annamalai University, Parangipettai, Tamil Nadu, India*

### ABSTRACT

An effort was made to assess the utility of 650 bp partial Cytochrome C oxidase subunit I (DNA barcode) gene in delineating the members of taxonomically ambiguous marine fin fishes (Family: Mugilidae). To address the issue we used all the 95 barcode sequences of Mugilidae family available at NCBI (National Centre for Biotechnological Information) along with the barcode data generated from Mugilidae fishes of Parangipettai coastal waters. The average GC content of Mugilidae was found to be 46.46%. *Crenimugil crenilabis* showed less GC content (44.55%) whereas *Liza macrolepis* showed high GC content (48.53%) among the mullet species studied. The phylogenetic and genetic distance data showed that *Mugil platanus* and *M. liza* represent the continuum of same species. Among the members of family Mugilidae, the genus *Mugil* might possibly contains more haplotype diversity as revealed by intra-species genetic distance data. Species within genera of Mugilidae family invariably clustered in single clade with high bootstrap value. We conclude that partial COI sequencing (barcoding) in identifying the members of the family and that way has resolved the taxonomic ambiguity among the members of the family Mugilidae.

**Key words:** Molecular phylogene, mugilidae taxonomy, intra-species variations, DNA barcoding, GC content

### INTRODUCTION

The family Mugilidae includes marine fin fishes belonging to 17 genera and more than 60 species of grey mullets (Papasotiropoulos *et al.*, 2007). This family was previously classified in the order Perciformes but is now considered the sole representative of the order Mugiliformes. They are most commonly found in the coastal waters and estuaries of the tropical and subtropical zones of the world. Mulletts are generally confined to shallow inshore waters; some of them are stenohaline and others exhibit different degrees of euryhalinity (Papasotiropoulos *et al.*, 2001). The grey mullets are of considerable importance in the capture and culture fisheries in many parts of the world but to Ichthyologists mugilids are one of the most difficult taxonomic groups due to its

morphologic conservity. Number of researchers has experienced difficulties with animal groups owing to highly overlapping morphological characters which are of taxonomic value and to explore the evolutionary relationship (Blasco-Costa *et al.*, 2008; Li *et al.*, 2011; Liu and Zhao, 2010; Ghajarieh *et al.*, 2006). These characters undergo marked changes during growth. Due to its high degree of conservative morphology, classifying them using classical morphometry and morphology has proved to be complex and difficult (Gilbert, 1993; Thomson, 1997). Since the early 19th century adipose eyelid, thickness of the lips with the presence and absence of papillae or lamellae, scales, teeth on palate and tongue and lateral scale numbers are being used as key characters for distinguishing genera and species in this family (Heras *et al.*, 2007). The results obtained were in most cases controversial, failing to provide any conclusive answers. The general morphological uniformity displayed by the members of Mugilidae family, restricts the number of suitable characters that can be used to answer phylogenetic questions unambiguously. As a result, the phylogenetic status of the family remains particularly obscure, especially at the interspecific level (Rossi *et al.*, 1998). This difficulty is due to very few suitable characters key characters to establish unambiguously the phylogenetic relationships among species (Caldara *et al.*, 1996). The distribution pattern of the species also compounds confusion among conventional taxonomists in classifying the species to the species level. The taxonomy of *Mugil liza* (Valenciennes, 1836), *M. platanus* (Günther, 1880) and *M. cephalus* (Linnaeus, 1758) is confounded in Western Atlantic. *M. liza* has been reported as a valid species with a distribution ranging from southern Florida to Rio de Janeiro (Thomson, 1997). A substitution of *M. liza* for *M. platanus* with more southerly latitudes represents a parapatric distribution involving separate but adjacent habitats. However, *M. liza* and *M. cephalus* are particularly difficult to distinguish morphologically (Heras *et al.*, 2007).

The close relationship between *M. liza* and *M. platanus* based on their shared haplotypes was proven by the mtDNA data (Fraga *et al.*, 2007). Their morphological similarities including gill rakers besides overlapping values of lateral series scales (LT) counts justifies combined consideration of these taxa (Eiras-Stofella *et al.*, 2001; Cousseau *et al.*, 2005). Despite the fact that the systematics of different Mugilid species has been revised many times, there are still substantial disagreements in foreign literature including Russian. Specifically, in Russian scientific literature *Liza aurata* is often named *Mugil auratus* (Papasotiropoulos *et al.*, 2001). The situation with *Liza haematocheila* is even more complicated. In Russia this species was described as *Mugil soiuy* while in Korea and Japan waters this species was initially described as *Mugil haematocheilus* later it was renamed *Liza haematocheila* (Papasotiropoulos *et al.*, 2001).

As regards the phylogeny of the Mugilidae family, it appears particularly obscure at both the intra- and inter-specific levels; it is extremely difficult to distinguish among species, especially in the juvenile stages because their morphological and physiological characters frequently do not exhibit significant differences (Papasotiropoulos *et al.*, 2002). It was also observed that due to close conservative morphology displayed by all mullets, many investigations based on various morphological characters did not elucidate this problem (Gilbert, 1993) while the use of the pharyngobranchial organ as a key character to address the taxonomy and the phylogeny of grey mullets were also proven futile (Harrison and Howes, 1991). Moreover karyotypic studies on several mullet species did not result in a clear phylogenetic figure (Rossi *et al.*, 2000, 1997). That was despite all these major revisions, the systematic status of some species and genera within the family Mugilidae is still confused (Rossi *et al.*, 1998).

Recently the efficacy of 648 bp of Cytochrome C oxidase subunit I gene in mitochondrial DNA (barcode) has been found to be useful in delineating morphologically cryptic organisms including

fishes to their species level (Ward *et al.*, 2005; Khan *et al.*, 2010; Kamaruzzaman *et al.*, 2011). However, comprehensive study on its utility towards mugilidae members is still scanty. Hence present study was aimed to explore (1) the phylogenetic status of mugilidae fishes, (2) species congruence within Mugilidae family and (3) efficiency of COXI gene in delineating the members of mugilidae fishes to its lowest possible taxon level beyond various geographical boundary.

## MATERIALS AND METHODS

**Sampling, DNA extraction, amplification and sequencing:** Mulletts from Parangipettai coastal waters (*Mugil cephalus*, *Liza tade* and *Liza parsia*) were collected alive and transported to the laboratory where the right side of the fishes was photographed and a cube of lateral muscle (5-7 mm) from left side of the fish was exercised for DNA isolation. The fishes were preserved in 95% ethanol for future references. Salting out protocol was adopted for precise and quick DNA isolation from the fish tissue.

The tissue was placed in 1.5 mL eppendorf tube and 500  $\mu$ L of solution I (50 mM Tris-HCl pH 8, 20mM EDTA pH8 and 2% SDS) was added. The tissue was homogenized with sterile homogenizer and 5  $\mu$ L of Proteinase K (20 mg mL<sup>-1</sup>) was added and quick vortexed. The sample was incubated at 55°C in water bath for 2 h with occasional mixing. Following incubation the sample was chilled over ice for 10 min and 250  $\mu$ L of solution II (6 M NaCl) was added and inverted several times for thorough mixing. The tube was chilled on ice for 5 min and centrifuged at 8000 rpm for 15 min. About 500  $\mu$ L of supernatant was carefully collected in to new-labeled 1.5 mL tube and twice the volume (i.e., 1 mL) of 100% AR grade ethanol was added to precipitate the DNA. The precipitate was pellet down at 8000rpm for 5 min and the supernatant was removed without touching the pellet. The DNA pellet was rinsed with 500  $\mu$ L of cold ethanol and centrifuged at 11000 rpm for 5 min. The supernatant was carefully removed and the excess liquid was drained using pipette. The pellet was partially dried (devoid of Ethanol) with lid off at 55°C on heating block. The pellet was re-suspended with 50-200  $\mu$ L of fresh sterile H<sub>2</sub>O depending on size of pellet (100  $\mu$ L average) by gently pipetting sample with wide-bore filter tip until dissolved. This dissolved DNA acted as a template for Polymerase Chain Reaction (PCR).

The fragment of COI was amplified by GeneAmp PCR system 9700. PCR was carried out in 25  $\mu$ L volumes [2.5  $\mu$ L of 10X PCR buffer, 1.5  $\mu$ L of MgCl<sub>2</sub> (2mM  $\mu$ L<sup>-1</sup>), 1  $\mu$ L of DNA template, 1  $\mu$ L of each primer (10 pmoles  $\mu$ L<sup>-1</sup>), 2 dNTPs (1 mM  $\mu$ L<sup>-1</sup>), 10 U of 1  $\mu$ L of Taq polymerase (Bioserve biotechnologies Pvt, Ltd, Hyderabad, India) and 15  $\mu$ L of sterile Mill Q water]. Fish F1 [5'-TCAACCAACCACAAAGACATTGGCAC-3'] and Fish R1 [5'-TAGACTTCTGGGTGGCCAAAGAATCA-3'] primers were employed for COI amplification (Ward *et al.*, 2005; Khan *et al.*, 2010). The thermocyclic conditions for PCR included the initial denaturation at 94°C for 1 min, five cycles of 94°C for 30 sec, annealing at 45°C for 40 sec and extension at 72°C for 1 min, with a final extension at 72°C for 10 min, followed by indefinite hold at 4°C.

Following PCR, about 10  $\mu$ L of PCR product with 2  $\mu$ L of bromo thymol blue were added to 2% agarose gel, prepared with 2.5  $\mu$ L of 1% Ethidium bromide and electrophorized at 90 V until the dye moved for 6 cm in the gel. The gel was moved to gel doc system for viewing the amplicons with the aid of UV trans-illuminator. Sequencing PCR was carried out using Dye terminator mix v3.1 and quantified in Euro bio-agarose gel. The samples were loaded onto MegaBace sequencer (MB 1000) at Bioserve Biotechnologies, Pvt. Ltd. Hyderabad, India.

**Sequence data analysis:** The electropherogram generated by automated DNA sequencer was read by Chromas Pro v1.42 and the sequences were carefully checked for mis-calls and base spacing. 95 barcode sequences of Mugilidae were extracted via FASTA format from NCBI. ClustalX 2.0.6 was used to align the nucleotide sequences (Thomson, 1997). The GC content of all 98 barcodes was estimated by BioEdit sequence alignment editor (Hall, 1999). MEGA 4.1 was used to construct phylogenetic trees via Neighbourhood joining method using Kimura 2-parameter and to calculate genetic distance of the given set of sequences (Tamura *et al.*, 2007). Barcode sequence of *Lates calcarifer* (IOBML45) sampled from Parangipettai coastal waters was used as an out group in Phylogenetic tree construction.

## RESULTS AND DISCUSSION

**Sequence features:** Ninety five sequences (collected from NCBI) belonging to various species of mullets (Table 1) disseminated around the world along with the barcode data (651 bp) generated from this work (constitutes 98 sequences belonging to 17 species representing 6 genera of mullets) was considered for phylogram construction and genetic distance analysis. Uniformity in GC content in the barcode region (5' cytochrome oxidase C subunit I) of the family Mugilidae was noted and it ranged between 44.55 and 48.53% while the maximum GC content was found in *Liza macrolepis* and the minimum was observed in *Crenimugil creilabis*. The average GC content of mullets was found to be 46.46%.

**Phylogenetic analysis:** Two phylogenetic trees were constructed to verify the following hypothesis. The first phylogenetic tree (Fig. 1) was constructed to test the efficacy of COI in delineating the members of mugilidae fishes to its species level. Barcode sequence of *Lates calcarifer* from Parangipettai coastal waters was used as an out group and this has been clearly distinguished as an out group in the phylogenetic tree. *Crenimugil* sp. shows more genetic relatedness to *Mugil* sp. as they were placed among the clusters of *Mugil* sp. (evident from Clade A). Interestingly, *Liza parsia* and *Liza tade* were placed among the clade A with members of genus *Mugil* sp. These

Table 1: Sequences of various species of mullets from different countries used to construct the phylogram in the present study

Species	No. of sequences used	Country
<i>Aldrichetta forsteri</i>	1	Australia
<i>Chelon haematocheilus</i>	3	Russia
<i>Chelon labrosus</i>	4	Spain
<i>Crenimugil crenilabis</i>	2	Canada
<i>Liza aurata</i>	8	Spain, Russia, Greece
<i>Liza parsia</i>	2	India
<i>Liza subviridis</i>	1	Russia
<i>Liza macrolepis</i>	7	India, Russia
<i>Liza saliens</i>	4	Spain, Russia, Greece
<i>Liza tade</i>	6	India
<i>Liza ramado</i>	4	Spain, Russia, Greece
<i>Mugil liza</i>	2	Spain
<i>Mugil cephalus</i>	17	Canada, Spain, Russia, Mexico, India, Greece
<i>Mugil curema</i>	15	Spain, USA,
<i>Mugil platanus</i>	10	Canada, Spain,
<i>Valamugil seheli</i>	1	Australia
<i>Valamugil cunnesius</i>	10	Canada, India, China, Russia




Fig. 1: Kimura 2-parameter distance Neighbour Joining (NJ) tree of 98 barcode sequences from 17 species belonging to family Mugilidae. COI sequence of *Lates calcarifer* collected from Parangipettai coastal waters was used as an out group. Specimen number denote the accession number of NCBI database and IOBML\* represents the specimens collected from Parangipettai coastal waters and the barcode sequences submitted at Barcode of Life Database (BOLD, www.barcodinglife.org)




Fig. 2: Phylogenetic hypothesis of Mugilidae based on COI data. The clade of congeneric species shows the genetic relatedness of *M. liza* and *M. platanus*. *Mugil cephalus* (IOBML37), *Liza tade* (IOBML2) and *Lates calcarifer* (IOBML45) was used as consecutive out groups to transparently reveal the genetic relationship between *M. liza* and *M. platanus*. IOBML\* represents the accession number assigned by Indian Ocean Barcoding of Marine Life at Barcoding of Life Database (BOLD, www.barcodinglife.org)

misplacements could be resolved by analyzing multiple mitochondrial gene sequencing techniques. Clade B contained the members of *Valamugil* sp., *Chelon* sp., *Aldrichetta* sp. and *Liza* sp. Within the family Mugilidae, *Liza aurata* was found to be the distant relative of *Mugil cephalus* as they were placed in two extremes of the clades of phylogenetic tree. However, each species of family Mugilidae strictly clustered with respective genus which explains the efficacy of COI in delineating the fishes to species level.

Occurrence of *Mugil liza* in the clades of *M. platanus* was an interesting observation which triggered the doubt of its close genetic relatedness and hence the second phylogram was constructed (Fig. 2). The phylogenetic tree was constructed using shuffled input sequences of *M. liza*, *M. platanus*, *M. cephalus* (different species within the same genus), *Liza tade* (species from another genus) and the same out group sequence of *Lates calcarifer* (a species from different family). The constructed phylogram proved the congeneric relatedness existing among the *Mugil liza* and *Mugil platanus*. The sequence of *Lates calcarifer* was clearly placed as an out group and the resolution of clade to genus and species level was apparent, as *Liza tade* was isolated from the members of *Mugil* sp., *Mugil cephalus* was transparently placed outside the clusters of *M. platanus*/*M.liza* complex in the clade.

Congeneric nature of *M. platanus* and *M. liza* was recently reported by multiple mitochondrial gene (16s rRNA, COI and Cyt b) sequencing technique (Fraga *et al.*, 2007). The confusion might have originated due to wide geographic distribution of the species. Robins *et al.* (1986), in his field guide to Atlantic coastal fishes, first reported the occurrence of *M. liza* in the coastal waters of Boston (USA) while Scorvo Filho *et al.* (1992) reported the occurrence of same species in Brazilian




Fig. 3: Intra-species variations among barcode sequences of species belonging to different genera in family Mugilidae

Table 2: Average inter-generic variations encoded in COI of family Mugilidae

	<i>Aldrichetta</i>	<i>Chelon</i>	<i>Crenimugil</i>	<i>Liza</i>	<i>Mugil</i>	<i>Valamugil</i>
<i>Aldrichetta</i>	*					
<i>Chelon</i>	0.235	0.050				
<i>Crenimugil</i>	0.238	0.263	0.178			
<i>Liza</i>	1.246	1.064	1.052	0.976		
<i>Mugil</i>	0.830	0.794	1.007	1.336	1.693	
<i>Valamugil</i>	0.263	0.230	0.254	1.123	0.702	0.069

\*Only one sequence of *Aldrichetta forsteri* is available

waters under the name *M. liza* which might have been the origin of the ambiguity regarding species nomenclature (Agapow *et al.*, 2004). Present study was well corresponded with this finding and constructed phylogram proved the congeneric nature of both the species and thus present conventional morphological taxonomy of these species (*M. platanus* and *M. liza*) should be reinvestigated.

**Genetic distance:** The genetic distance within and between the species and genera of family Mugilidae was calculated using pair-wise distance analysis via Maximum likelihood method (Table 2). The intra species genetic distance within the members of genus *Chelon* sp. was found low (0.05). It may be due to its recent divergence from other members of the family. The intra species variation of genus *Mugil* was found (1.693) higher (Fig. 3). This might be due to possible haplotype diversity existing within the species of *Mugil cephalus*, as reported earlier (Fraga *et al.*, 2007). The analysis showed that among the Mugilidae members, members of genus *Liza* were distantly related to the members of genus *Mugil* as they were grouped at both extreme ends of the phylogram. The overall genetic distance of Mugilidae family was found to be 1.078 which showed the richness of haplotype diversity existing within this family.

## CONCLUSION

The present analysis showed the close congeneric relationship between *M. liza* and *M. platanus* indicating high degree of gene flow within them and they do not support differentiation at species level. Until new phylogenetic groups are fully identified and implemented, the present species' status should be preserved to minimize risks of loss of important components of biodiversity. The COI sequence in the phylogram constructed clearly clustered the species of same genus in individual group, proving the efficacy of COI gene in delineating the members of Mugilidae to their



species level. Hence we conclude that COI sequence (DNA barcode) could be potentially used to identify the morphologically cryptic individual members of family Mugilidae.

#### **ACKNOWLEDGMENT**

We are thankful to Dr. T. Balasubramanian, Professor and Director of our centre for the encouragement and the authorities of our university for the facilities. We also extend our sincere thanks to Dr. Wafer, National Institute of Oceanography, Goa and Bioserve Biotechnologies (India) Pvt. Ltd. for their constant supporting throughout the research.

#### **REFERENCES**

- Agapow, P.M., O.R. Bininda-Emonds, K.A. Crandall, J.L. Gittleman, G.M. Mace, J.C. Marshall and A. Purvis, 2004. The impact of species concept on biodiversity studies. *Qual. Rev. Biol.*, 79: 161-179.
- Blasco-Costa, I., F.E. Montero, D.I. Gibson, J.A. Balbuena, J.A. Raga, L.S. Shvetsova and A. Kostadinova, 2008. A revision of the species of *Saturnius* Manter, 1969 (Digenea: Hemiuridae), parasites of mullets (Teleostei: Mugilidae). *Syst. Parasitol.*, 71: 53-74.
- Caldara, F., L. Bargelloni, L. Ostellari, E. Penzo, L. Colombo and T. Patarnello, 1996. Molecular phylogeny of grey mullets based on mitochondrial DNA sequence analysis: Evidence of differential rate of evolution at the intrafamily level. *Mol. Phylogenet. Evol.*, 6: 416-424.
- Cousseau, M.B., M.G. Castro, D.E. Figueroa and A.E. Gosztonyi, 2005. Does *Mugil liza* valenciennes 1836 (Teleostei: Mugiliformes) occur in argentinean waters. *Rev. Biol. Marina y Oceanografia*, 40: 127-131.
- Eiras-Stofella, D.R., P. Charvet-Almeida, E. Fanta and A.C. Vianna, 2001. Surface ultrastructure of the gills of the mullets *Mugil curema*, *M. liza* and *M. platanus* (Mugilidae, Pisces). *J. Morphol.*, 2: 122-133.
- Fraga, E., H. Schneider, M. Nirchio, E. Santa-Brigida, L.F. Rodrigues-Filho and I. Sampaio, 2007. Molecular phylogenetic analyses of mullets (Mugilidae, Mugiliformes) based on two mitochondrial genes. *J. Applied Ichthyol.*, 23: 598-604.
- Ghajariéh, H., M. Bruford, H.A. Dawah and C.F.C.S. Dodd, 2006. Mitochondrial phylogenetics of UK Eurytomids. *J. Entomol.*, 3: 167-179.
- Gilbert, C.R., 1993. Geographic distribution of the striped mullet in the Atlantic and eastern Pacific oceans. *Florida Sci.*, 56: 204-210.
- Hall, T.A., 1999. BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl. Acids. Symp. Ser.*, 41: 95-98.
- Harrison, I.J. and G.J. Howes, 1991. The pharyngobranchial organ of mugilid fishes: Its structure, variability, ontogeny, possible function and taxonomic utility. *Bull. Br. Mus. Nat. Hist. Zool.*, 57: 111-132.
- Heras, S., M.I. Roldan, C.M. Gonzalez and M.B. Cousseau, 2007. *Mugil cephalus*: Cosmopolitan species or species complex?. *Rapp. Comm. Int. Mer. Mediterr.*, 38: 499-499.
- Kamaruzzaman, B.Y., B.A. John, K. Zaleha and K.C.A. Jalal, 2011. Molecular phylogeny of horseshoe crab. *Asian J. Biotechnol.*, 3: 302-309.
- Khan, S.A., P.S. Lyla, B.A. John, C.P. Kuamr, S. Murugan and K.C.A. Jalal, 2010. DNA barcoding of *Stolephorus indicus*, *Stolephorus commersonii* and *Terapon jarbua* of parangipettai coastal waters. *Biotechnology*, 9: 373-377.

- Li, H., C. Zhao, H. Xu, B. Yu and C. Wu, 2011. A method to rapidly identify to what species unknown animals are closely related. *Asian J. Anim. Vet. Adv.*, 6: 362-370.
- Liu, W.Y. and C.J. Zhao, 2010. Comprehensive genetic analysis with mitochondrial DNA data reveals the population evolution relationship between chinese gamecocks and their neighboring native chicken breeds. *Asian J. Anim. Vet. Adv.*, 5: 388-401.
- Papasotiropoulos, V., E. Klossa-Kilia, G. Kiliias and S.N. Alahiotis, 2001. Genetic divergence and phylogenetic relationships in grey mullets (Teleostei: Mugilidae) using allozyme data. *Biochem. Genet.*, 39: 155-168.
- Papasotiropoulos, V., E. Klossa-Kilia, G. Kiliias and S.N. Alahiotis, 2002. Genetic divergence and phylogenetic relationships in grey mullets (Teleostei: Mugilidae) based on PCR-RFLP analysis of mtDNA segments. *Biochem. Genet.*, 40: 71-86.
- Papasotiropoulos, V., E. Klossa-Kilia, S.N. Alahiotis and G. Kiliias, 2007. Molecular phylogeny of grey mullets (Teleostei: Mugilidae) in Greece: Evidence from sequence analysis of mtDNA segments. *Biochem. Genet.*, 45: 623-636.
- Robins, C.R., G.C. Ray and J. Douglass, 1986. *Atlantic Coast Fishes*. Houghton Mifflin. Co. Boston, Massachusetts, pp: 354.
- Rossi, A.R., E. Gornung and D. Crosetti, 1997. Cytogenetic analysis of *Liza ramada* (Pisces, Perciformes) by different staining techniques and fluorescent *in situ* hybridization. *Heredity*, 79: 83-87.
- Rossi, A.R., E. Gornung, D. Crosetti, S. De Innocentiis and L. Sola, 2000. Cytogenetic analysis of *Oedalechilus labeo* (Pisces: Mugilidae), with report of NOR variability. *Mar. Biol.*, 136: 159-162.
- Rossi, A.R., M. Capula, D. Crosetti, D.E. Campton and L. Sola, 1998. Genetic divergence and phylogenetic inferences in five species of *Mugilidae* (Pisces: Perciformes). *Mar. Biol.*, 131: 213-218.
- Scorvo Filho, J.D., E.R. Almeida-Dias, L.M.S. Ayrosa and P.F. Colherinhas-Novato, 1992. Efeito da densidade sobre o desenvolvimento de alevinos de tainha listrada (*Mugilplatanus*) em agua doce. *Bol. Inst. Pesca.*, 19: 105-109.
- Tamura, K., J. Dudley, M. Nei and S. Kumar, 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24: 1596-1599.
- Thomson, J.M., 1997. The mugilidae of the world. *Mem. Queensland Mus.*, 41: 457-562.
- Ward, R.D., T.S. Zemlak, B.H. Innes, P.R. Last and P.D. Hebert, 2005. DNA barcoding Australias fish species. *Philos. Trans. Royal Soc. Lond. B Biol. Sci.*, 360: 1847-1857.